



Distribution of an Isopod Parasite *Mothocya renardi* in the Host *Strongylura leiura* from Arabian Sea Gujarat Coast, India

Ritesh V. Borichangar ^a, Mihir R. Patel ^b, Jeet N. Parmar ^{a*},
Jay B. Patel ^b, Sheetal K. Bharda ^c and Dhruvi P. Kotadiya ^a

^a Department of Fisheries Resource Management, College of Fisheries Science, Kamdhenu University Navsari, 396 450, India.

^b Department of Aquatic Animal Health Management, College of Fisheries Science, Kamdhenu University Navsari, 396 450, India.

^c Department of Fisheries Resource Management, ICAR- Central Institute of Fisheries Education Mumbai, 400 061, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajee/2024/v23i8580>

Open Peer Review History:
This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119433>

Original Research Article

Received: 02/05/2024

Accepted: 04/07/2024

Published: 08/07/2024

ABSTRACT

Crustacean isopod parasites, particularly in the marine environment, are prevalent around warmer seas and can significantly impact local fish populations. *Mothocya renardi*, a notable species of isopod parasites, commonly infests fishes in shallow, turbid coastal waters. This study aimed to

*Corresponding author: E-mail: jeetparmar97@gmail.com;

investigate the occurrence, taxonomy, and morphology of *M. renardi* infesting *Strongylura leiura* in the Arabian Sea around the northwest coast of India. Samples were collected from Dholai fishing harbor (Navsari District of Gujarat), during the post-monsoon months. Three specimens of *S. leiura* were found infected with pairs of male and female *M. renardi* in the brachial chambers, with all female parasites being brooded ovigerous. Detailed morphological analysis confirmed the presence of *M. renardi*, extending its known geographical range to the Arabian Sea.

Keywords: Isopod parasite; *Mothocya renardi*; *Strongylura leiura*; Arabian sea; India.

1. INTRODUCTION

Over 7000 species of crustacea are parasitic, exploiting an enormous range of invertebrate and vertebrate hosts in marine and freshwaters, and many more crustaceans exhibit other kinds of interspecific associations (symbioses) including commensalism, inquilinism, mutualism, and phoresis, with a similarly wide range of hosts [1]. Copepoda, Branchiura, Isopoda, Amphipoda, and Ostracoda, are known as the main groups comprising the parasitic crustacean, in addition, these groups are either ecto or endo parasites. In which the three major groups Copepoda, Isopoda, and Branchiura comprise the crustacean ecto-parasites for fish [2]. Crustacean ectoparasites on marine fish are diverse. Parasitic isopods are typically marine and usually inhabit the warmer seas they are mainly found in shallow waters along the coastline [3,4,5]. They are blood-feeding; several species settle in the buccal cavity of fish, while others live in the gill chamber or on the body surface including the fins. Isopods can cause morbidity and mortality in fish populations [6].

Isopods are typically dorsoventrally flattened peracaridans characterised by a relatively short head fused to the first pereon segment which bears the maxillipeds, a pereon of seven segments each bearing a pair of similar pereopods and the ventral brood pouch in mature females, and a pleon which usually exhibits some fusion between the posterior pleon segments and telson. Though free-living isopods tend to be detritivores while adult parasitic forms feed on host blood or host haemolymph. Their mouthparts form a cone with maxillipeds that tear at the flesh and tiny pointed mandibles that pierce into the tissue to penetrate blood vessels or blood sinuses [7].

There are three major groups of isopoda: Cymothoids, Epicaridians and Gnathiids. Cymothoids are parasites of fish, both as immature forms and as adults. Epicaridians are parasites of crustacea, again as immatures and adults. Gnathiids are larval parasites of fish, the

adults being free living and non-feeding. Cymothoid isopods are protandrous hermaphrodites. The first male to parasitize a fish changes into a female. Egg development apparently depends on the presence of a male, for each batch. Internal fertilisations occur immediately after the female has moulted, male deposits sperm into the female's genital opening. The sperm then fertilizes the eggs inside the female's body [8]. Gravid females release eggs into a brood pouch or 'marsupium' formed from their ventral oostegites. Here the eggs embryonate, hatch and undergo several moults to form the 'manca' stage. These are released from the brood pouch, sometimes more or less simultaneously as a result of contractions from the parent. The parent then moults, feeds, digests the meal, and eventually produces the next batch of eggs. Several batches may be produced during her lifespan. The mancae have only six pairs of legs (compared to seven in juveniles and adults), large compound eyes, and heavily setose pleopods with which they swim extremely rapidly. After a short free-swimming period, they are parasitic and need to find a fish to take their first meal within one to two days or they will die [7].

During the work to assess the ichthyofaunal biodiversity of Dholai fishing harbor, some fish specimens were caught with parasites attached to them. Dholai is a minor fishing harbor situated along the southwest coast of Gujarat, India; it is an important place for the fishery sector of the south Gujarat region. It spans over 23 hectares of land and has facilities such as a net mending shed, auction halls, small yard for minor boat repair and maintenance. The majority of the fishing vessels carry out operations from the Gulf of Cambay region in the north to the Ratnagiri coast in the south [9]. The objective of this study was to identify the host, and parasite and describe its morphology along with its taxonomy.

2. MATERIALS AND METHODS

During the monthly specimen collection of ichthyofaunal diversity from Dholai Fishing

Harbour on the southwest coast of Gujarat (Fig. 1), the specimens of *Strongylura leiura* were collected during post monsoon months of September to November 2023, and brought to the College of Fisheries Science, Navsari. Upon close inspection of the specimen, a pair of male and female ectoparasite was observed in all three specimens. It was identified as *Mothocya renardi*. Identification of the parasite was done using the Journal of Natural History [10]. It was preserved in 70% alcohol. Host (*Strongylura leiura*) identification was done through FAO species identification sheets [11]. Morphometrics and Meristics were recorded and preserved in 10% formalin. Both the specimens of the parasite (*M. renardi*) and host were submitted to the aquatic biodiversity museum at the College of Fisheries Science, Navsari (Accession No: A 5.1.1.2 for *S. leiura* and C 3.1.1.1 *M. renardi*).

3. RESULTS

The operculum of the *Strongylura leiura* was cut from the 7th branchiostegal rays upwards to remove the whole gill cover; this uncovered the whole gill filaments with the parasite still sitting in

its original place. It was carefully removed for observation. Upon close examination, it was determined that there was tissue damage to the internal brachial cavity and gill filaments of host.

For identification according to the standard identification guides total 13 parameters of parasite and 22 morphometric parameters and meristic characters of host were recorded and presented here in Tables 1 and 2, respectively.

***Mothocya renardi* (Gill ectoparasite)**

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea
 Superclass: Peracarida
 Class: Malacostraca
 Subclass: Eumalacostraca
 Superorder: Peracardia
 Order: Isopoda
 Suborder: Cymothoidea
 Superfamily: Cymothooidea
 Family: Cymothoidae
 Genus: *Mothocya*
 Species: *renardi*




Fig. 1. Location of Dholai Fishing Harbour (QGIS 3.32.0)

Family Cymothoidae has 39 genera, while 30 species come under Genus *Mothocya*. *Mothocya renardi* was originally described as *Livoneca renardi* by Bleeker, in 1857 which was revised as *Mothocya renardi*. Other synonyms which are currently categorized as unaccepted names include *Irona renardi*, (Schioedte & Meinert, 1884), *Irona melanosticta* (Barnard, 1914), and *Irona robusta*, (Nair, 1950).

Species description:

Female: The total length of the specimen is 32 mm; the body is widest at pereonite 3 which is 2.4 times as long as wide. Cephalon width surpasses its length and the first pereonite is crescent-shaped to house the cephalon. The eyes are based on the front lateral side of the cephalon; they are small, one-third the width of the cephalon. The width of pereonite from Pereonite 1 to Pereonite 3 increases, body is widest at pereonite 3. From Pereonite 4 to Pereonite 7 it gradually decreases. Pereopods are without spines, with their main mechanism being attaching to the host. Among all pereopods, the first pereopod is the shortest, while the seventh pereopod is the longest. Pleotelson is expanded horizontally, width is 1.2 times its length. Oostegites were visible in the ventral view of the brood pouch in the fresh specimen.

Colouration: Body uniformly creamy white without any pigmentation, including brood pouch while in fresh specimen. The hue of

the brood pouch darkens to a deep brown within a few days following the death of the parasite.

Male: Males are generally smaller than females. Around 18 mm in total length. Body twice in length than width. Small eyes set on broad cephalon. Pereonites increase in width from the first to the third pereonite body widest at the third pereonite (8 mm), and decrease from the fourth to the seventh pereonite, similar as seen in the female. The length and width of the Pleotelson are symmetrical.

Coloration: Body creamy white with a band of dark pigmentation running along the median line of pereonites and pleonites.

***Strongylura leiura*, Banded needlefish (Host):**

Taxonomy:

- Kingdom: Animalia
- Phylum: Chordata
- Subphylum: Vertebrata
- infraphylum: Ganthostomata
- Parvphylum: Osteichthyes
- Gigaclass: Actinopterygii
- Superclass: Actinopteri
- Class: Teleostei
- Order: Beloniformes
- Family: Belonidae
- Genus: *Strongylura*
- Species: *Strongylura leiura*

Table 1. Morphometric parameters of *Mothocya renardi*

Parameter	Female		Male	
	mm	% TL	mm	% TL
Total length (TL)	32	100.00	18	100.00
Width	13	40.63	8	44.44
Cephalon	3	9.38	3	16.67
Eye diameter	1	3.13	<1	-
Pereonite 1	3	9.38	6	33.33
Pereonite 2	9	28.13	7	38.89
Pereonite 3	13	40.63	8	44.44
Pereonite 4	12	37.50	7	38.89
Pereonite 5	11	34.38	6	33.33
Pereonite 6	10	31.25	5	27.78
Pereonite 7	8	25.00	4	22.22
Pleotelson width	11	34.38	5	27.78
Pleotelson length	9	28.13	4	22.22

Table 2. Morphometric parameters of host *Strongylura leiura*

Sr. No.	Parameter	% TL	Sr. No.	Parameter	% TL
1	Total length (TL)	100.00	12	Pectoral base	2.18
2	Standard length	90.10	13	Pre-pectoral length	26.14
3	Snout length	13.86	14	Pelvic fin length	5.54
4	Head length	25.54	15	Pelvic fin base	0.99
5	Eye diameter	2.18	16	Pre-pelvic length	55.84
6	Pre-orbital length	13.86	17	Anal fin length	17.62
7	Post-orbital length	16.63	18	Anal fin base	16.63
8	Dorsal fin length	15.05	19	Pre-anal base	71.29
9	Dorsal fin base	14.06	20	Body depth	5.94
10	Pre-dorsal length	73.27	21	Upper jaw length	13.07
11	Pectoral fin length	8.32	22	Lower jaw length	13.66

Coloration: The dorsal side of the whole body green colored. A silver-colored strip runs on the lateral side of the body, which expands posteriorly. Below the strip, the ventral side of the body is white in color.

4. DISCUSSION

Gill ectoparasite *Mothocya renardi* was observed from specimens of *Strongylura leiura* from Dholai fishing harbour, northwest coast of India. The banded needlefish resides in shallow coastal waters with high turbidity, around mangrove patches with sandy bottoms, and according to observations of Mertz [3] marine isopods reside

in shallow coastal waters, from these observations, it can be concluded that *M. renardi* have infested the host in shallow coastal waters. Out of all three specimens observed, all of them had infestations, with a pair of male and female present in the brachial cavity of host specimens. All three female were at the ovigerous stage in their lifecycle. The damage to the host was observed as necrosis of gill filaments, and damage of the branchial cavity by females (Fig. 2), this causes disturbance to the normal respiratory function, which could result in the failure of adequate oxygen supply and inorganic ions for normal metabolic activities of fish, which leads to morbidity and mortality of host.




Fig. 2. a) *Mothocya renardi* female dorsal view; b) Female lateral view; c) Female ventral view; d) *M. renardi* male dorsal view e) male lateral view f) male ventral view; g) *Strongylura leiura* position of parasite in gills lateral view h) host ventral view of brachial cavity i) host anterolateral view, damage done to brachial chamber by female parasite

Previous records confirmed the presence of *M. renardi* from India, northeast coast [12], southeast coast [13], and Malabar coast [14], but this is the first recorded instance of *M. renardi* infestation from the northwest coast of India.

5. CONCLUSION

This study observed *Mothocya renardi*, a gill ectoparasite, infesting specimens of the *Strongylura leiura* from Dholai fishing harbor on the northwest coast of India. The presence of these parasites in the brachial cavity of the host fish, particularly in shallow coastal waters, suggests a habitat preference for such environments. All observed specimens of the banded needlefish were found to be infested, with both male and female parasites present. While previous records have documented the presence of *M. renardi* along various coastal regions of India, this study marks the first recorded instance of infestation from the northwest coast of India.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Boxshall G, Hayes P. Biodiversity and taxonomy of the parasitic Crustacea. In Smit N, Bruce N, Hadfield K (Eds.), *Parasitic Crustacea*. Cham, Switzerland: Springer. 2019;73-134.
2. Ahmed F, Ali H, Bakheet Y, Ahmed Y. Fish's crustacean parasites: Types, prevalence, clinical signs, and control. *Sohag Journal of Science*. 2022;7(2):123-129.
3. Mertz LA. Grzimek's animal life encyclopedia (2nd ed.). Detroit, MI: Thomson-Gale; 2004.
4. Panakkool-Thamban A, Kappalli S, Kottarathil HA, Gopinathan A. *Mothocya renardi* (Bleeker, 1857) (Crustacea: Isopoda: Cymothoidae) parasitising *Strongylura leiura* (Bleeker)(Belonidae) off the Malabar coast of India: Redescription, occurrence and life-cycle. *Systematic Parasitology*. 2016, Jul;93:583-99.
5. Fujita H, Kawai K, Deville D, Umino T. Quatrefoil light traps for free-swimming stages of cymothoid parasitic isopods and seasonal variation in their species compositions in the Seto Inland Sea, Japan. *International Journal for Parasitology: Parasites and Wildlife*. 2023, Apr 1;20:12-9.
6. Ravichandran S, Rameshkumar G. Effect of parasitic isopods in commercial marine fishes. *Journal of Aquatic Biology and Fisheries*. 2014;2:574-579.
7. Boxshall G, Lester R, Grygier MJ, Hoeg JT, Glenner H, Shields JD, Lützen J. Crustacean parasites. In K. Rohde (Ed.), *Marine Parasitology*. 2005;123-169.
8. Shuster SM. Male alternative reproductive strategies in a marine isopod crustacean (*Paracerceis sculpta*): The use of genetic markers to measure differences in fertilization success among α -, β -, and γ -males. *Evolution*. 1989;43(8):1683-1698.
9. Borichangar RV, Parmar JN, Vyas UD, Tandell LV, Patel MR, Vala RB, Patel PP. First record of Grey Bamboo Shark, *Chiloscyllium griseum* (Muller & Henle 1838) from Dholai Port, Southwest Coast of Gujarat, India. *Indian Journal of Ecology*. 2023;50(6):2096-2101.
10. Bruce NL. Revision of the isopod crustacean genus *Mothocya* Costa, in Hope, 1851 (Cymothoidae: Flabellifera), parasitic on marine fishes. *Journal of Natural History*. 1986;20(5):1089-1192.
11. Fischer W, Bianchi G. (Eds.). *FAO species identification sheets for fishery purposes: Western Indian Ocean (Fishing Area 51)*. Rome: Food and Agricultural Organization of the United Nations; 1984.
12. Mohapatra SK, Mohanty SR, Behera RK, Seth JK, Mohapatra A. First record of *Mothocya renardi* and *Mothocya collettei* (Isopoda: Cymothoidae) from the northern part of East Coast of India and new host record of *Mothocya collettei*. *Journal of Parasitic Diseases*. 2021;45:1-4.
13. Ravichandran S, Sivasubramanian K, Rameshkumar G, Veerappan N. High prevalence and infestation of *Mothocya renardi* (Isopoda, Cymothoidae) in marine fish *Strongylura leiura* (Bleeker 1850). *Journal of Parasitic Diseases*. 2015; 40:1386-1391.

14. Panakkool-Thamban A, Kappalli S, Kottarathil HA, Gopinathan A. *Mothocya renardi* (Bleeker, 1857) (Crustacea: Isopoda: Cymothoidae) parasitising *Strongylura leiura* (Bleeker) (Belonidae) off the Malabar coast of India: Redescription, occurrence, and life-cycle. Systematic Parasitology. 2016;93:583-599.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/119433>