



Varietal Evaluation of Ber (*Zizyphus mauritiana* Lamk.) for Yield and Fruits Quality Attributes under Bundelkhand Region of Uttar Pradesh, India

Dharmendra Kumar ^{a*}, Om Prakash ^{a++}, Sarfaraj ^a,
Siddharth Kumar ^a, Vikas Kumar ^a and Shubham Yadav ^b

^a Department of Fruit Science, College of Horticulture, Banda University of Agriculture and Technology, Banda, U.P., India.

^b Department of Fruit Science, College of Horticulture, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i92400

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102442>

Original Research Article

Received: 08/05/2023

Accepted: 10/07/2023

Published: 22/07/2023

ABSTRACT

The experiment was conducted at Instruction farm, College of Horticulture, Banda University of Agriculture and Technology, Banda (U.P.) during the year 2021-2022. The trail consisting of ten treatments (T₁ -Thai Ber, T₂ -Umran, T₃ -Banarasi Karaka, T₄ -Seb, T₅ -Gola, T₆ -Chuhara, T₇ -Kaithali, T₈ -Mundia, T₉ -Elaichi and T₁₀ -Rashmi). The variability was recorded in fruit shape (ovate oblong, oblong, ovate and round), fruit colour (green, yellow green, golden yellow and green) and pulp colour (creamy white and creamy). The yield attributes variability was observed in fruit length (3.40-5.72 cm), fruit width (2.13-4.20 cm), fruit weight (20.27-48.33 g), fruit

**Professor;

*Corresponding author: E-mail: kkdharmendra4159@gmail.com;

volume (21.21-49.25 cc), stone length (1.59-2.80 cm), stone width (0.86-2.07 cm), stone weight (1.41-2.83 g), pulp weight (16.22-44.24 g), pulp and stone ratio (11.31-15.61) and specific gravity (0.95-0.98). The quality attributes parameter also showed wide range of variation in chemical properties i.e., TSS (10.61-16.62 °Brix), ascorbic acid (80.5-115.33 mg/100g), reducing sugars (4.05-7.11 %), non-reducing sugars (3.90-4.75 %), total sugars (7.95-11.86 %), acidity (0.32-0.75 %), protein (0.40-0.70 %) and pH (3.0-5.0). Therefore, on the basis of quality attribute among the different evaluation; the variety T₇ (Kaithali), T₂ (Umran) and T₃ (Banarasi Karaka) found to be superior. These promising varieties can be recommended for commercial cultivation at farmer's field.

Keywords: Ber; varietal evaluation; yield; fruit quality.

1. INTRODUCTION

Ber (*Zizyphus mauritiana* Lamk.) is a fruit native to India and China that belongs to the *Rhamnaceae* family. It is regarded as "King of Arid zone Fruits" and also known as "Poor man's Apple" (Nidhi, et al., 2019). Ber fruit has also been mentioned to holly book like Ramayana and Mahabharata [1]. According to Pareek [2], it can be found in both wild and cultivated forms from the warmer arid and semi-arid regions up to an altitude of 1500 meters above sea level. The centre of origin of Ber is Central Asia (India & China), where it is found growing under different climatic conditions, requirement of temperatures ranges from 39-42 °C and it can tolerate temperatures as high as 49-50 °C [3]. It is cultivated widely for its resistance to grow in drought and other diversified soil and climatic condition. It can be grown well in inferior soil with pH as high as 9.0 in arid and semi-arid regions [4]. The Ber leaves have also been tested by mixing in rations of goats in arid and semi-arid regions. Trees leaves and shrubs constitute about 60 % of the diet of goats and 40 % of sheep's [5]. Many wild Ber shrubs i. e. (*Zizyphus mauritiana*, *Z. rotundifolia* and *Z. nummularia*) were found growing on neglected lands, uncultivated lands in Uttar Pradesh (India). It also provides enough leaf biomass (pala) for fodder during the lean season and is suitable for hay and silage for goats due to its 11-13 % crude protein content [6]. Ber fruits are high in ascorbic acid and high in vitamins A, B and C as well as minerals like calcium, phosphorus and iron [7,8,9]. Ber fruit is high in proteins, fats, fibre, TSS, ascorbic acid, reducing, non-reducing and total sugars [10,11,12,13,4]. It is a drupe, globose to ovoid in shape, skin smooth or rough, glossy thin but tough, yellowish to reddish or blackish, flesh white, crisp and juicy, sub acid to sweet, becoming mealy in fully ripe fruit and colour of Ber fruit is changed from green to yellow to chocolate brown with the

maturity and ripening [14]. Ber fruit is normally eaten fresh and is a rich source of ascorbic acid, necessary minerals and carbohydrates [15].

Under Bundelkhand region, meagre information is available on evaluation of ber cultivars and there is no varietal recommendation. So, there is an urgent need to carry out research experiments for evaluation of ber varieties for the large-scale adoption of ber fruit crop for under Bundelkhand region of Uttar Pradesh.

2. MATERIALS AND METHODS

Experiment was conducted at Instruction farm, College of Horticulture Banda, while physico-chemical evaluation in the Post-Graduate Laboratory, Department of Basic Science, College of Horticulture, Banda University of Agriculture & Technology, Banda (U.P.) during the year 2021-2022. The experimental site is situated at 24° 53'-25° 55' N latitudes and 80° 07'-81° 34' E longitudes. The pH obtained the range from 6.65- 8.56 (acidic and alkaline soil). Average annual rainfall is 90 cm and most of it about 88-90% received during only in three months i. e. July, August and September. The annual temperature is comparatively high and it ranges in between 32-49.5 °C, 23-35 °C and 10-22 °C for summer, rainy and winters respectively. The experiment was carried out in randomized block design with three replications. The details of ten treatments were T₁ -Thai Ber, T₂ -Umran, T₃ -Banarasi Karaka, T₄ -Seb, T₅ -Gola, T₆ - Chhuhara, T₇ -Kaithali, T₈ -Mundia, T₉ -Elaiichi and T₁₀ -Rashmi. Statistical analysis of the data for physical parameters observation were recorded on Initial fruit set, Fruit drop, Fruit retention, Fruit shape, Fruit colour, Fruit length, Fruit width, Fruit weight, Fruit volume, Stone width, Stone length, Stone weight, Pulp colour, Pulp weight, Pulp and Stone ratio, Specific

gravity and Yield. Chemical parameters observations were recorded on TSS, Ascorbic acid, Reducing sugars, Non-reducing sugars, Total Sugars, Acidity, Protein and Determination of pH. The physical parameters like: Fruit weight, Stone weight and Pulp weight measure by electronic weighting machine and Fruit length, Fruit width, Stone width, Stone length was calculated with the help of Vernier Callipers. The total soluble solids were measured using an Erma hand refractometer (0-32 °Brix). Ascorbic acid was determined by using 2, 6-Dichlorophenol-indophenols visual titration method (Johnson and Dana, 1948). Reducing sugars were determined by Nelson method (1944). The percentage of non-reducing sugars was obtained by subtracting the values of reducing sugar from total sugars and multiplying by 0.95 (Somogyi, 1952). Total sugars were determined by Dubois et al. (1956) method. The titratable acidity of the pulp was determined by titrating 10 ml juice against 0.1 N NaOH using phenolphthalein as an indicator as per the procedure suggested by Sadashivam (1991). Lowry et al. (1951) used the Folin Ciocalteu's Phenol reagent to calculate protein content. A pH paper is a piece of paper used to find out if a solution is basic, acidic or neutral. The data obtained during experimentation was statistically analysed as per the method given by Panse and Sukhatme (1985) and results were evaluated at 5% level of significance.

3. RESULTS AND DISCUSSION

The different varieties of Ber differed significantly in initial fruit set, fruit drop and fruit retention. Significantly maximum initial fruit set 64.36 % was recorded by (T₇-Kaithali) and minimum fruit set 36.70 % was recorded in (T₉-Elaichi). The similar fruit set variability in ber were reported by Singh et al [16]. The minimum percentage of fruit set refers to the natural tendency of trees to shed their immature fruits. Highest fruit drop 71.10 % was recorded by (T₇-Kaithali) and the lowest fruit drop 57.00 % was recorded by (T₁-Thai Ber). Similar fruit drop variability was also reported by Tarai and Ghosh [17] and Singh et al [16]. Mostly fruit drop during early stage of fruit development may be due to quality or unsuccessful fertilization or ovule degeneration. Maximum fruit retention 43.00 % was observed by (T₁-Thai Ber) and the minimum fruit retention 28.90 % was observed by (T₇-Kaithali). Similar fruit retention variability was reported by Tarai and Ghosh [17]. The maximum fruit length, fruit width,

fruit weight, fruit volume, stone length, stone width, stone weight, Pulp weight and Pulp and Stone ratio were recorded in (T₁-Thai Ber) were resulting 5.72cm, 4.20cm, 48.33g, 49.25cc, 2.80cm, 2.07cm, 2.83g, 44.24g and 15.61 respectively. The minimum fruit length, fruit width, fruit weight, fruit volume, stone length, stone width, stone weight, pulp weight and pulp and stone ratio were recorded in (T₉-Elaichi) were resulting 3.40cm, 2.13cm, 20.27g, 21.21cc, 1.59cm, 0.87cm, 1.41g, 16.22g and 11.31 respectively. The similar results of fruits length, fruit width, fruit weight and fruit volume variability in ber were reported by Obeed et al [10], Shukla et al [18], Kumar et al [19] and Kumar et al [20]. Stone length, stone width, stone weight, pulp weight and pulp & stone ratio these results are in conformity with the previous to Shukla et al [18] and Kumar et al [20]. The maximum specific gravity 0.98 was observed in treatment (T₁-Thai Ber) and (T₃-Banarasi karaka). The lowest value 0.95 was observed (T₂-Umran) and (T₉-Elaichi). However, treatment (T₄-Seb) was found statistically at par with (T₅-Gola) and (T₇-Kaithali). The similar variations in specific gravity of ber studies were earlier Ghosh and Mathew [21], Obeed et al [10], Shukla et al [18]. The similar results of yield reported by Ghosh and Mathew [21], Rao and Subramaniyam [22], Tarai and Ghosh [17] and Singh et al [16]. Out of ten treatments, 5 are ovate-oblong (T₁-Thai Ber, T₂-Umran, T₄-Seb, T₆-Chhuhara and T₇-Kaithali), two are oblong (T₃-Banarasi Karaka and T₁₀-Rashmi), two are round (T₅-Gola and T₉-Elaichi) and one is ovate (T₈-Mundia). Among these, three treatments had light green (T₁-Thai Ber), (T₃-Banarasi karaka) and (T₇-Kaithali), three had yellow-green (T₂-Umran), (T₈-Mundia) and (T₉-Elaichi), two had golden yellow (T₅-Gola) and (T₁₀-Rashmi) and two had green (T₄-Seb) and (T₆-Chhuhara). Among them, seven treatments had creamy white (T₁-Thai Ber), (T₃-Banarasi karaka), (T₄-Seb), (T₆-Chhuhara), (T₈-Mundia), (T₉-Elaichi) and (T₁₀-Rashmi) and three treatments had creamy (T₂-Umran), (T₅-Gola) and (T₇-Kaithali).

The fruit quality attributes were recorded. The significantly maximum total soluble solid 16.62 °Brix was noticed in treatment (T₇-Kaithali) followed by treatment 16.22 °Brix (T₂-Umran) and 16.13 °Brix (T₃-Banarasi Karaka), while the minimum TSS value 10.61 °Brix was noted by treatment (T₁-Thai Ber). However, treatment (T₂-Umran) was found statistically at par with (T₃-Banarasi Karaka). Similar trend of total soluble

Table 1. Effect of different varietal evaluation substance on Initial fruit set, Fruit drop, fruit retention, fruit length, fruit width, fruit weight and fruit volume

| Treatments | Physical parameters | | | | | | |
|----------------------------------|-----------------------|----------------|---------------------|-------------------|------------------|-------------------|-------------------|
| | Initial fruit set (%) | Fruit drop (%) | Fruit retention (%) | Fruit length (cm) | Fruit width (cm) | Fruit weight (cm) | Fruit volume (cc) |
| T ₁ : Thai Ber | 44.50 | 57.00 | 43.00 | 5.72 | 4.20 | 48.33 | 49.25 |
| T ₂ : Umran | 61.87 | 69.87 | 32.83 | 4.80 | 3.37 | 32.29 | 33.27 |
| T ₃ : Banarasi Karaka | 58.00 | 67.17 | 30.13 | 4.70 | 3.22 | 31.40 | 32.75 |
| T ₄ : Seb | 54.70 | 62.77 | 37.23 | 5.20 | 3.73 | 35.92 | 36.95 |
| T ₅ : Gola | 51.67 | 61.63 | 38.37 | 4.58 | 3.10 | 30.75 | 31.40 |
| T ₆ : Chhuhara | 38.97 | 64.67 | 35.33 | 4.50 | 3.00 | 28.72 | 29.74 |
| T ₇ : Kaithali | 64.36 | 71.10 | 28.90 | 5.01 | 3.53 | 33.42 | 34.45 |
| T ₈ : Mundia | 47.43 | 59.67 | 40.33 | 4.30 | 2.80 | 26.25 | 27.20 |
| T ₉ : Elaichi | 36.70 | 65.63 | 34.37 | 3.40 | 2.13 | 20.27 | 21.21 |
| T ₁₀ : Rashmi | 40.97 | 58.70 | 41.30 | 3.70 | 2.25 | 23.60 | 24.60 |
| Mean(T) | 49.92 | 63.82 | 36.18 | 4.59 | 3.13 | 31.09 | 32.08 |
| SE(m)± | 0.29 | 0.22 | 0.23 | 0.12 | 0.05 | 0.36 | 0.36 |
| C.D. at 5% | 0.87 | 0.67 | 0.67 | 0.36 | 0.15 | 1.10 | 1.07 |

Table 2. Effect of different varietal evaluation substance on stone length, stone width, stone weight, pulp weight, pulp and stone ratio, specific gravity and yield

| Treatments | Physical parameters | | | | | | |
|----------------------------------|---------------------|------------------|------------------|-----------------|----------------------|------------------|------------------|
| | Stone length (cm) | Stone width (cm) | Stone weight (g) | Pulp weight (g) | Pulp and stone ratio | Specific gravity | Yield (Kg/plant) |
| T ₁ : Thai Ber | 2.80 | 2.07 | 2.83 | 44.25 | 15.61 | 0.98 | 41.38 |
| T ₂ : Umran | 2.44 | 1.60 | 2.50 | 28.32 | 11.31 | 0.95 | 56.46 |
| T ₃ : Banarasi Karaka | 2.34 | 1.52 | 2.42 | 27.88 | 11.50 | 0.98 | 53.23 |
| T ₄ : Seb | 2.60 | 1.81 | 2.63 | 31.25 | 11.88 | 0.97 | 50.17 |
| T ₅ : Gola | 2.25 | 1.45 | 2.36 | 27.88 | 11.15 | 0.97 | 47.22 |
| T ₆ : Chhuhara | 2.15 | 1.21 | 2.21 | 24.14 | 10.90 | 0.96 | 39.17 |
| T ₇ : Kaithali | 2.50 | 1.70 | 2.53 | 29.10 | 12.18 | 0.97 | 58.77 |
| T ₈ : Mundia | 2.00 | 1.11 | 1.92 | 21.13 | 10.98 | 0.96 | 44.80 |
| T ₉ : Elaichi | 1.59 | 0.86 | 1.41 | 16.22 | 11.47 | 0.95 | 34.77 |
| T ₁₀ : Rashmi | 1.84 | 1.02 | 1.83 | 19.12 | 10.47 | 0.96 | 37.21 |
| Mean(T) | 2.251 | 1.430 | 2.27 | 26.78 | 11.75 | 0.96 | 46.32 |
| Range | 1.59-2.80 | 0.86-2.07 | 1.41-2.83 | 16.22-44.24 | 11.31-15.61 | 0.95-0.98 | 34.77-58.77 |
| SE(m)± | 0.028 | 0.019 | 0.01 | 0.09 | 0.22 | 0.006 | 0.26 |
| C.D. at 5% | 0.085 | 0.058 | 0.04 | 0.27 | 0.67 | 0.017 | 0.79 |

solid was reported by Ghosh and Mathew [21], Ibrahim et al [23], Rao and Subramaniyam [22] and Gupta et al [24]. The significantly highest ascorbic acid (115.33 mg/100g) was recorded by the treatment (T₁ -Thai Ber) followed by 113.17 mg/100g (T₁₀ -Rashmi) and 112.50 mg/100g (T₈ -Mundia). While minimum ascorbic acid value recorded in (80.50 mg/100g) was noted by the variety (T₇ -Kaithali). Similar variation was observed Ghosh and Mathew [21], Obeed et al

[10] and Gupta et al [24]. The significantly highest reducing sugar 7.11 % was observed in (T₇ -Kaithali). However, followed by treatment 6.82 % (T₂ -Umran) and 6.55 % (T₃ -Banarasi Karaka). The minimum reducing sugars value 4.05 % was noted by the treatment (T₁ -Thai Ber). The significantly maximum non-reducing sugars 4.75 % content was observed in (T₇ -Kaithali), followed by treatment 4.62 % (T₂ -Umran) and 4.43 % (T₃ -Banarasi Karaka). The

minimum value 3.90 % was observed by the treatment (T₁-Thai Ber). However, treatment (T₂-Umran) was found statistically at par. The significantly maximum total sugars 11.86 % was recorded in treatment (T₇-Kaithali), followed by treatment 11.43 % (T₂-Umran) and 10.98 % (T₃-Banarasi Karaka). The minimum total sugars value 7.95 % was noted by the treatment (T₁-Thai Ber). The different treatment, significantly highest acidity 0.75 % was recorded in (T₁-Thai Ber) and followed by 0.70 % (T₁₀-Rashmi) and 0.60 % (T₈-Mundia). However, lowest acidity 0.32 % was recorded in (T₇-Kaithali). While treatment (T₁₀-Rashmi) statistically was recorded at par within (T₈-Mundia) and (T₆-Chuhara). Similar variation in terms of reducing sugars, non-reducing sugars, total sugars and acidity

were reported by Obeed et al. [10], Ibrahim et al. [23], Singh et al [25], Kumar et al [19] and Anjum et al [26]. Significantly maximum protein 0.70 % was observed in treatment (T₁-Thai Ber), followed by treatment 0.67 % (T₁₀-Rashmi) and 0.64 % (T₈-Mundia). In contrast, minimum protein 0.40 % among the treatment was observed in (T₇-Kaithali). However, treatment (T₁₀-Rashmi) statistically was observed at par within (T₈-Mundia), (T₅-Gola) and (T₄-Seb). Similar variation of ber was reported by Kumar et al [19]. The significantly highest (5.00) pH was recorded in Treatment (T₇-Kaithali) followed by 4.80 (T₈-Mundia) and 4.50 (T₃-Banarasi Karaka). The lowest pH value 3.00 was noted by (T₁-Thai Ber). Similar variation of ber was reported by Para [27,28].

Table 3. Effect of different varietal evaluation substance on fruit shape, fruit color and pulp color

| Treatments | Yield attributes | | |
|----------------------------------|------------------|-----------------|--------------|
| | Fruit shape | Fruit color | Pulp color |
| T ₁ : Thai Ber | Ovate Oblong | Light green | Creamy white |
| T ₂ : Umran | Ovate Oblong | Yellow green | Creamy |
| T ₃ : Banarasi Karaka | Oblong | Light green | Creamy white |
| T ₄ : Seb | Ovate Oblong | Green | Creamy white |
| T ₅ : Gola | Round | Golden yellow | Creamy |
| T ₆ : Chuhara | Ovate Oblong | Green | Creamy white |
| T ₇ : Kaithali | Ovate Oblong | Light green | Creamy |
| T ₈ : Mundia | Ovate | Greenish yellow | Creamy white |
| T ₉ : Elaichi | Round | Yellow green | Creamy white |
| T ₁₀ : Rashmi | Oblong | Golden yellow | Creamy white |

Table 4. Variability in chemical characteristics of Ber varieties under Bundelkhand region Uttar Pradesh

| Treatments | Chemical Characteristics | | | | | | | pH |
|-------------------------------------|--------------------------|--------------------------------|------------------------|-------------------------------|------------------------|----------------|----------------|---------------|
| | (TSS) °Brix | Ascorbic acid (mg/ 100pulp) | Reducing sugars (%) | Non reducing sugars (%) | Total sugars (%) | Acidity (%) | Protein (%) | |
| T ₁ : Thai Ber | 10.61 | 115.33 | 4.05 | 3.90 | 7.95 | 0.75 | 0.70 | 3.00 |
| T ₂ : Umran | 16.22 | 87.37 | 6.82 | 4.62 | 11.43 | 0.38 | 0.43 | 4.80 |
| T ₃ : Banarasi Karaka | 16.13 | 83.83 | 6.55 | 4.43 | 10.98 | 0.40 | 0.47 | 4.50 |
| T ₄ : Seb | 13.30 | 101.77 | 5.50 | 4.22 | 9.73 | 0.49 | 0.57 | 3.80 |
| T ₅ : Gola | 13.00 | 104.87 | 5.10 | 4.21 | 9.31 | 0.54 | 0.60 | 3.60 |
| T ₆ : Chuhara | 14.07 | 95.17 | 5.85 | 4.06 | 9.91 | 0.44 | 0.54 | 4.00 |
| T ₇ : Kaithali | 16.62 | 80.50 | 7.11 | 4.75 | 11.86 | 0.32 | 0.40 | 5.00 |
| T ₈ : Mundia | 12.50 | 112.50 | 4.80 | 4.09 | 8.89 | 0.60 | 0.64 | 3.40 |
| T ₉ : Elaichi | 15.60 | 91.00 | 6.21 | 4.38 | 10.59 | 0.42 | 0.50 | 4.20 |
| T ₁₀ : Rashmi | 11.72 | 113.17 | 4.35 | 4.12 | 8.47 | 0.70 | 0.67 | 3.20 |
| Mean(T) | 13.98 | 98.55 | 5.63 | 4.28 | 9.91 | 0.50 | 0.55 | 4.06 |
| Range | 10.61- 16.62 | 80.5-115.33 | 4.05-7.11 | 3.90-4.75 | 7.95- 11.86 | 0.32-0.75 | 0.40-0.70 | 3.00- 5.00 |
| SE(m)± | 0.16 | 0.48 | 0.02 | 0.092 | 0.09 | 0.08 | 0.05 | 0.04 |
| C.D. at 5% | 0.49 | 1.45 | 0.05 | 0.276 | 0.27 | 0.23 | 0.15 | 0.11 |

4. CONCLUSION

On the basis of the fruit quality attributes, the varieties (T₇ -Kaithali), (T₂ -Banarasi Karaka) and (T₃ -Umran) were screened as promising. These varieties can be further used for processing purpose and preparation of ber based products. These promising varieties can be recommended to the farmers for commercial cultivation. Significant variation was noted in all varieties of ber. The ber varieties are classified as early, mid and late on the basis of days taken to maturity. The following varieties of ber may be recommended for cultivation in Bundelkhand region of Uttar Pradesh. (T₄ -Seb and T₅ -Gola) as early season varieties, (T₇ -Kaithali and T₈ -Mundia) as mid-season varieties and (T₂ -Umran and T₉ -Elaichi) as late season varieties.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Pal R, Kumar A. Studies on effect feeding on fruiting and yield of ber (*Zizyphus mauritiana* Lamk.) under sodic soil. Int. J. Microbial. App. Sci. 2020;9(12):2991-2994.
2. Pareek OP. ber: International Centre for Underutilized crops, Southampton, U.K. 2001:248-266.
3. Candolle AD. Origin of cultivated plants. Bibliotheque Scientifiqu Internationale. 1886;43: 385.
4. Kumari S, Bhat DJ, Wali VK, Bakshi P, Jasrotia A. Physico-chemical studies of different ber (*Zizyphus mauritiana* Lamk.) germplasm under rainfed condition of Jammu. Bioscan. 2015;10(3):1427-1430.
5. Shinde BS, Barbind RP, Patil GR. Feeding value for Ber (*Zizyphus nummularia*) leaves for the growth performance of osmanabadi kids. Indian Journal of Small Ruminants. 2000;6(2):82-85.
6. Tewatia BS, Khirwar SS. Utilization of Ber (*Z. mauritina* L.) leaves hay and silage in goat. Indian J. Animal nutrition. 2002;19(4):329-333.
7. Shobha D, Bharathi P. Value addition to ber (*Zizyphus mauritiana* Lamk.) through preparation of pickle. Karnataka Journal of Agriculture Sciences. 2007;20(2):353-355.
8. Pareek S, Kitinoja L, Kaushik RA, Paliwal R. Post-harvest physiology and storage of ber. Stewart Post Harvest Rev. 2009;5(5):1-10.
9. Choi SH, Ahn JB, Kozukue N, Levin CE, Friedman M. Distribution of amino acids, flavonoids, total phenolics and anti-oxidative activities of jujube (*Zizyphus jujube*) fruits and seeds harvested from plants grown in Korea. J. Agric. Food Chem. 2011;59:6594-6604.
10. Obeed RS, Harhash MM, Mawgood AL. Fruit properties and genetic diversity of five ber (*Zizyphus mauritiana* Lamk.) cultivars. Pak. J. Biol. Sci. 2008;11(6):888-893.
11. Koley Kaur TK, Nagal C, Walia S, Jaggi S, Sarika SJ. Antioxidant activity and phenolic content in genotypes of Indian jujube (*Zizyphus mauritiana* Lamk.) Arab. J. Chem. 2011;4:321-324.
12. Krishna H, Parashar A. Phytochemical constituents and antioxidant activity and phenolics content in genotypes of Indian jujube. Arab. J. Chem. 2012;4:321-324.
13. Zozio S, Servent A, Hubert O, Hiol A, Pallet D, Mbeguie AD. Physico- chemical and biochemical characterization of ripening in jujube fruits from two accessions grown in Guadeloupe. Sci. Hortic. 2014;175:290-297.
14. Pareek S. Nutritional composition of jujube fruits. Emir. J. Food Agric. 2013;25(6):463-470.
15. Pareek S, Fageria MS, Dhaka RS. Performance of ber genotype under arid condition. Curr. Agric. 2002;26:63-65.
16. Singh VK, Shankar K, Tiwari NK, Rao OP. Effect of certain nutrients on fruit set, fruit retention, physical characters and yield of Ber fruits (*Zizyphus mauritiana* Lamk.) cv. Banarasi Karaka. International Journal of Bio- resource and stress management. 2016;7(4):648-652.
17. Tarai RK, Ghosh SN. Varietal evaluation for yield and yield parameters of ber under semi-arid region of West Bengal. J. Horti. Sci. 2010;5(1):17-20.
18. Shukla G, Singh R, Ram RB, Dwivedi DH. Genetic variability and correlation analysis in ber (*Zizyphus mauritiana* Lamk.) germplasm growth in Lucknow. Hort Flora Research Spectrum. 2012;1(2):122-126.
19. Kumar M, Singh S, Pathak DV, Gadara RK. Impact of natural ripening on physico-chemical characteristics of ber fruits. Agric international. 2016;3(2):12-18.
20. Kumar P, Srivastava AK, Prakash Om, Thakur N, Singh P. Germplasm survey, collection and evaluation of ber (*Zizyphus*

- mauritiana* Lamk.) under Bundelkhand Region of Uttar Pradesh. Ind. J. Pure App. Bio. Sci. 2021;9(1):259-266.
21. Ghosh SN, Mathew B. Performance of nine ber (*Zizyphus mauritiana* Lamk.) cultivars on top working in the semi-arid region of west Bengal. J. Appl. Hort. 2002;4(1):49-51.
 22. Rao KD, Subramaniyam K. Evaluation of yield performance of ber varieties under scarce rainfall zone. Agric. Sci. Digest. 2010;30(1):57-59.
 23. Ibrahim M, Shafique MZ, Helali OH, Rahman MM, Biswas, Islam MS. Studies on the physiological and biochemical composition of different ber (*Zizyphus mauritiana* Lamk.) cultivars at Rajshahi. Bangladesh J. Sci. Ind. Res. 2009;44(2): 229-232.
 24. Gupta N, Wali VK, Singh VB, Bakshi P, Jamwal M, Kumar V. Variability in physico-chemical characteristics of ber (*Zizyphus mauritiana* Lamk.) varieties under rainfed conditions of Shivalik foothills of Himalayas. Madras Agriculture J. 2012; 99(4-6):218-220.
 25. Singh B, Pathak S, Kulshreshtha SK, Ramprasad. Comparative study of physico-chemical attributes of ber (*Zizyphus mauritiana* Lamk.) fruits from different cultivars grown in eastern Uttar Pradesh. Environment & Ecology. 2015; 33(4):1539-1541.
 26. Anjum MA, Rauf A, Bashir MA, Ahmed R. The evaluation of biodiversity in some indigenous Indian jujube (*Zizyphus mauritiana*) germplasm through physico-chemical analysis. Acta Sci. Pol. Hortorum Cultus. 2018;17(4):39- 52.
 27. Para PA. Effect of Indian jujube pulp on physico-chemical and sensory characteristics of chicken sausages. Journal of meet Science and Technology. 2014;2(4):90-94.
 28. Dalal N, Neeraj, Vinita B. Value added products from ber (*Zizyphus mauritiana* Lamk.). Int. J. Curr. Microbiol. App. Sci. 2019;8(1):1603-1615.

© 2023 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/102442>