



Influence of Crop Establishment Methods and Weed Management Practices on Growth and Yield Attributes of Wheat

Deepak Kumar Yadav^{a*}, S. K. Verma^a, Chandrabhushan^a,
Nihal Chandra Mahajan^a and Prateek Kumar^a

^a Department of Agronomy, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi 221005, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i183412

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102769>

Original Research Article

Received: 22/05/2023
Accepted: 24/07/2023
Published: 03/08/2023

ABSTRACT

An experiment was conducted in *Rabi* season of 2019-20 and 2020-21; consist of six crop establishment methods *viz.*, Conventional till wheat - residue removed, Conventional till wheat - residue incorporation at 20 days before sowing (30 cm height), Conventional till wheat - mulching (6 t ha⁻¹), Zero till wheat without residue and Zero till wheat – residue as mulch (6 t ha⁻¹), zero till wheat- residue retention (30 cm) was assigned to main plots and weed management practices *viz.*, weed free (weeds were removed with the help of hand weeding during entire crop period), Weedy check, Pendimethalin (1 kg ha⁻¹, PE) *fb* sulfosulfuron + metsulfuron (30 + 2 g ha⁻¹, PoE) and Pendimethalin (1 kg ha⁻¹, PE) *fb* iodosulfuron + carfentrazone (2.5 + 20 g ha⁻¹, PoE) were allocated to sub plots. Among crop establishment methods, zero tillage with residue mulching @ 6 t ha⁻¹ and

*Corresponding author: E-mail: dkvnsydv@gmail.com;

among the weed management practices, Pendimethalin (1 kg ha⁻¹, PE) fb iodosulfuron + carfentrazone (2.5 + 20 g ha⁻¹, PoE) recorded maximum plant height and yield attributes of wheat over rest of the treatments.

Keywords: Residue; mulch; wheat yield; weed management.

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is the most extensively farmed cereal crop worldwide. It has an area of 220.41 million hectares and produces 734 million tonnes (FAO, 2019). With current growth rates, the global human population will total 2.4 billion people by 2050 [1]. "This generates an alarming situation in order to alleviate the hunger crisis. Wheat is the staple feed for 20% of the world's population [2] (Xuan, 2020), providing one-fifth of daily calorie intakes and 21% of dietary protein requirements worldwide" [3]. "Wheat production has a direct impact on global food security due to its widespread popularity, adaptation, and consumption. Smith, [1]. Wheat is second after rice in terms of acreage (45 million hectares) and production (107 million tonnes) in India. The indo-gangetic plains, which account for 15% of total geographic area and are a prominent area of the green revolution, are responsible for the production boom" [4]. "This region alone accounts for 50% of overall production and 40% of the country's food grain requirements. These country's plains are one of the world's largest, with deep and alluvium fertile soil suited for double and triple cropping" [5,6].

Farmers' practises of wheat sowing include repetitive tillage by cultivator and rotavator, which cannot be undertaken in a combine harvested paddy field, encouraging farmers to burn crop residue in the field to quickly remove crop residue. Burning agricultural residue is detrimental to the sustainability of any cropping system since it generates smokes that are bad for the environment, loses nutrients in straw, degrades soil health, and is economically unfeasible. In addition, residue burning in the field produced ash, which influenced the efficacy of pre-emergence herbicide treatment. These paddy wastes can be used in a variety of industries, including energy, packing material, and bedding material, as well as mulching, composting, feeding, and thatching [7]. To address this issue, zero tillage sowing practises clean the row zone, open the soil, place fertilisers and seed, and cover the seed in a single pass. It can also be used in residual field conditions to prevent residue burning. In this method, minimal

soil disturbance with residue improves soil organic carbon sequestration, saves US \$ 52 ha⁻¹ in ploughing fuel, and yields 5 to 7 percent higher [8]. Usman et al. [9] discovered that RT had the lowest weed density and dry weed biomass with the highest grain yield of wheat, which was statistically equal to grain yield under ZT when compared to CT.

Farmers did manual weeding, which is not economical due to salary increases [10] and also not practicable because weed eradication takes too long. Another challenge for timely weed management is migration of agricultural workforce from rural to urban areas. The issue of manual weed management has led farmers to seek alternate weed management solutions for sustainable crop production. Weed management with herbicide is the easiest and most viable method in this scenario [11], which considerably improves weed control in modern agriculture [12]. Using a single herbicide without rotating it, and using greater doses encourages herbicide resistance in weeds.

2. MATERIALS AND METHODS

The present study was conducted in Rabi season of 2019-20 and 2020-21. The soil of field was sandy clay loam in texture, with slightly alkaline (pH 7.38) in reaction. The soil was low in organic carbon (0.43%) available nitrogen (232.25 kg ha⁻¹) and phosphorus (18.63 kg ha⁻¹) and medium in available potassium (178.54 kg ha⁻¹). Average values for bulk density 1.42 g cm⁻³, particle density 2.62 g cm⁻³ and EC were 0.264 ds m⁻¹. The experiment was conducted in split plot design with twenty four treatment combinations. Treatments consisted of six crop establishment methods viz. Conventional till wheat - residue removed, Conventional till wheat - residue incorporation at 20 days before sowing (30 cm height), Conventional till wheat - mulching (6 t ha⁻¹), Zero till wheat without residue and Zero till wheat - residue as mulch (6 t ha⁻¹) zero till wheat- residue retention (30 cm) were assigned to main plots and weed management practices viz. weed free (weeds were removed with the help of hand weeding during entire crop period), Weedy check, Pendimethalin (1 kg ha⁻¹, PE) fb sulfosulfuron + metsulfuron (30 + 2 g ha⁻¹, PoE)

and Pendimethalin (1 kg ha⁻¹, PE) *fb* iodosulfuron + carfentrazone (2.5 + 20 g ha⁻¹, PoE) were allocated to sub plots. The experiment was replicated three times. Wheat variety 'MACS 6222' was sown under different crop establishment methods at row distance of 20 cm by opening slits with zero-till-drill machine. Application of Pendimethalin as pre-emergence (PE) were done at 2 DAS; and post emergence (PoE) herbicide as per treatment applied at 35 DAS of crop with the help of sprayer fitted with flat fan nozzle. The spray volume was 500 l ha⁻¹. Half amount of nitrogen and full dose of phosphorus and potash were applied as basal at the time of sowing, ¼ part of nitrogen was top dressed after first irrigation and remaining ¼ part of nitrogen was top dress at spike initiation stage. Four irrigations were given to crop as per need.

3. EXPERIMENTAL FINDINGS

3.1 Plant Height (cm)

In crop establishment practices, plant height were significantly influenced at all the stages except at 30 DAS. Among all crop establishment practices, zero till wheat with 100 % rice straw mulching (ZTW+SM₁₀₀) shows higher plant height at all stages during both the years of field observation. Conventional till wheat with 100 % mulching (CTW+SM₁₀₀) and zero tillage wheat with rice residue retention (30 cm) (ZTW+RR) was at par with the treatment ZTW+SM₁₀₀; whereas significantly higher than the conventional tillage practices with rice residue incorporation (30 cm) (CTW+RI), zero tillage wheat with no residue (ZTW-R) and conventional tillage wheat with no residue (CTW-R) during both the years, respectively. But at 30 DAS the difference of height was non-significant. Herbicide significantly influences the plant height at all the stage of field observation during both the years. "Application of pendimethalin @ 1 kg a.i. ha⁻¹ (PE) *fb* iodosulfuron + carfentrazone (PoE) @ 2.5 + 20 g a.i. ha⁻¹ recorded higher plant height and remains at par with pendimethalin @ 1 kg a.i. ha⁻¹ (PE) *fb* sulfosulfuron + metsulfuron (PoE) @ 30 + 2 g a.i. ha⁻¹ over the weedy check plot during both year of observation at all stage of observation but none of practices comparable with weed-free treatment" [13].

3.2 Effective Tillers (Per Running Meter)

In crop establishment practices, plant height was significantly influenced at all the stages except at 30 DAS. Among all crop establishment practices,

zero till wheat with rice straw mulching @ 6 ton ha⁻¹ (ZTW+SM) shows higher plant height at all stages during both the years of field observation. Conventional till wheat with rice straw mulching @ 6 ton ha⁻¹ (CTW+SM) and zero tillage wheat with rice residue retention (30 cm) (ZTW+RR) was at par with the treatment ZTW+SM; whereas significantly higher than the conventional tillage practices with rice residue incorporation (30 cm) (CTW+RI), zero tillage wheat with no residue (ZTW-R) and conventional tillage wheat with no residue (CTW-R) during both the years, respectively. But at 30 DAS the difference of height was non-significant. Herbicide significantly influences the plant height at all the stage of field observation during both the years. Application of pendimethalin @ 1 kg a.i. ha⁻¹ (PE) *fb* iodosulfuron + carfentrazone (PoE) @ 2.5 + 20 g a.i. ha⁻¹ recorded higher plant height and remains at par with pendimethalin @ 1 kg a.i. ha⁻¹ (PE) *fb* sulfosulfuron + metsulfuron (PoE) @ 30 + 2 g a.i. ha⁻¹ and significantly superior over the weedy check plot during both year of observation at all stage of observation but none of practices comparable with weed-free treatment.

3.3 Spike Length (cm)

Spike length were influenced by crop establishment practices parameter; zero till wheat with rice straw mulching @ 6 ton ha⁻¹ (ZTW+SM) shows slightly higher spike length compare to rest of crop establishment practices and lowest in conventional tillage wheat – no residue treatment; difference between the crop establishment practices were not reached upto significant variation during both the years of finding. Weed free treatment were recorded slightly higher spike length than the rest of practices and lowest in weedy check but the difference was remains non-significant during both the years.

3.4 Grains Spike⁻¹

Spike length were influenced by crop establishment practices parameter; zero till wheat with rice straw mulching @ 6 ton ha⁻¹ (ZTW+SM) shows slightly more spike length compare to rest of crop establishment practices and lowest in conventional tillage wheat – no residue treatment; difference between the crop establishment practices were not reached upto significant variation during both the years of finding. Among weed management practices in wheat crop weed free treatment were recorded slightly more spikelet per spike than the rest of practices and lowest in weedy check but the

difference was remains non-significant during both the years.

3.5 1000 Grain Weight (g)

Crop establishment practices parameter influence the 1000 grain weight in Table 2, which reveals that zero till wheat with rice straw mulching @ 6 ton ha⁻¹ (ZTW+SM) shows slightly higher grain weight compare to rest of crop establishment practices; whereas in weed management 1000 grains weight slightly more in weed free and lowest in weedy check but the difference in both establishment and weed management were remains non-significant during both the years.

4. DISCUSSION

The superiority of ZTW+SM in respect of plant height and crop growth parameters can be discussed in light of the fact that this treatment providing better suppression of weed spp. [14-17] by minimum soil disturbance for seed bed preparation, more availability of nutrients to crop with higher microbial activity which facilitate better crop performance in zero tillage; and at same time application residue as mulching

provide optimum growing environment by ensuring more availability of light, moisture and nutrients [18-20] than other crop establishment treatments. lower weed density and dry weight of weed in tank mix application of pendimethalin @ 1 kg a.i. ha⁻¹ (PE) fb iodosulfuron + carfentrazone (PoE) @ 2.5 + 20 g a.i. ha⁻¹ because less competition faced by crop which enhance the availability of soil moisture, nutrient, space and light so growth parameter were improved. Results are corroborated with the research findings of Upasani et al., [15]; Islam et al. [21] and Yadav et al. [22]. Increase in yield contributing factor due to vigorous crop growth parameter by obtaining favourable condition. More dry matter accumulation by crop were translocate to formation of better yield attributing factor. Pendimethalin @ 1 kg a.i. ha⁻¹ (PE) fb iodosulfuron + carfentrazone (PoE) @ 2.5 + 20 g a.i. ha⁻¹ recorded highest than the weedy check plot; whereas superiority in the weed free plot were recorded. The reason behind it may be due to the lesser crop-weed competition by broad spectrum herbicide effectively reduce the weed density which facilitate better growing. It confirms the conclusion drawn by Verma et al. [7]; Jaiswal, et al. [23]; Duary et al., [24] and Kumar et al. [25], [26].

Table 1. Effect of crop establishment and weed management practices on plant height (cm)

Treatments	30 DAS		60 DAS		90 DAS		At Harvest	
	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21
Crop establishment practices								
R ₁ : Zero till wheat residue as mulch	22.84	23.87	59.65	62.63	89.42	93.89	93.08	97.74
R ₂ : Conventional till wheat residue as mulch	23.75	24.94	57.42	60.01	86.08	89.95	89.61	93.64
R ₃ : Zero till wheat - residue retention at 30 cm height	22.03	22.89	55.43	57.59	83.08	86.32	86.49	89.86
R ₄ : Conventional till wheat - rice residue incorporation at 20 days before sowing	21.39	22.22	53.85	55.95	80.72	83.86	84.02	87.30
R ₅ : Zero till wheat residue removed	22.11	22.49	52.93	53.83	79.33	80.68	82.59	83.99
R ₆ : Conventional till wheat - residue removed	22.34	22.72	51.52	52.40	77.23	78.54	80.40	81.76
SEm±	0.64	0.67	1.32	1.38	1.98	2.06	2.06	2.15
LSD (p=0.05)	NS	NS	4.16	4.33	6.24	6.50	6.50	6.76
Weed management								
W ₁ : Weed free	24.40	25.25	60.02	62.13	89.97	93.13	93.66	96.94
W ₂ : Pendimethalin (1 kg ha ⁻¹ , PE) fb iodosulfuron + carfentrazone (2.5 + 20 g ha ⁻¹ , PoE)	23.13	23.93	56.89	58.89	85.28	88.28	88.78	91.89
W ₃ : Pendimethalin (1 kg ha ⁻¹ PE) fb sulfosulfuron + metsulfuron (30 + 2 g ha ⁻¹ , PoE)	22.13	22.91	54.45	56.37	81.62	84.50	84.97	87.96
W ₄ : Weedy check	19.99	20.67	49.17	50.88	73.70	76.27	76.72	79.40
SEm±	0.43	0.45	0.41	0.42	0.61	0.64	0.64	0.66
LSD (p=0.05)	1.25	1.29	1.17	1.22	1.76	1.82	1.83	1.90

Table 2. Effect of crop establishment and weed management practices on yield attributes

Treatments	No. of effective tillers (per running meter)		Spike length (cm)		No. of grains spike ⁻¹		1000 grain weight (g)	
	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21
Crop establishment practices								
R ₁ : Zero till wheat residue as mulch	96.21	101.05	11.07	11.62	37.10	38.96	37.80	39.70
R ₂ : Conventional till wheat residue as mulch	93.00	97.19	10.65	11.13	35.71	37.32	36.39	38.03
R ₃ : Zero till wheat - residue retention at 30 cm height	90.11	93.62	10.28	10.68	34.47	35.82	35.13	36.50
R ₄ : Conventional till wheat - rice residue incorporation at 20 days before sowing	87.81	91.24	9.99	10.38	33.49	34.80	34.13	35.46
R ₅ : Zero till wheat residue removed	86.48	88.81	9.82	10.08	32.92	33.81	33.54	34.45
R ₆ : Conventional till wheat - residue removed	84.44	86.72	9.56	9.82	32.04	32.91	32.65	33.53
SEm±	1.95	2.00	0.25	0.26	0.82	0.86	0.79	0.87
LSD (p=0.05)	6.14	6.29	0.78	0.80	2.59	2.70	2.48	2.75
Weed management								
W ₁ : Weed free	96.76	100.47	11.14	11.56	37.33	38.76	38.04	39.49
W ₂ : Pendimethalin (1 kg ha ⁻¹ , PE) fb iodosulfuron + carfentrazone (2.5 + 20 g ha ⁻¹ , PoE)	92.23	95.76	10.55	10.96	35.38	36.74	36.06	37.44
W ₃ : Pendimethalin (1 kg ha ⁻¹ PE) fb sulfosulfuron + metsulfuron (30 + 2 g ha ⁻¹ , PoE)	88.69	92.09	10.10	10.49	33.87	35.16	34.51	35.83
W ₄ : Weedy check	81.02	84.10	9.12	9.47	30.58	31.74	31.16	32.35
SEm±	0.75	0.62	0.09	0.08	0.29	0.26	0.29	0.27
LSD (p=0.05)	2.14	1.77	0.25	0.23	0.82	0.76	0.82	0.77

5. CONCLUSION

Among crop establishment methods, zero tillage with residue mulching @ 6 t ha⁻¹ and among the weed management practices, Pendimethalin (1 kg ha⁻¹, PE) fb iodosulfuron + carfentrazone (2.5 + 20 g ha⁻¹, PoE) recorded maximum plant height and yield attributes of wheat over rest of the treatments.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Smith P. Malthus is still wrong: We can feed a world of 9–10 billion, but only by reducing food demand. *Proceedings of the Nutrition Society*. 2015;74:187-190.
- McFall KL, Fowler ME. Overview of wheat classification and trade. *Wheat Science and Trade*; 2009.
- Shiferaw B, Smale M, Braun HJ, Duveiller E, Reynolds M, Muricho G, Crops that feed the world, past successes and future challenges to the role played by wheat in global food security. *Food Security*; 2013.
- Koshal AK. International changing current scenario of rice-wheat system in Indo-Gangetic plain region of India. *Journal of Scientific and Research Publications*. 2014;4(3):1-13.
- Gangwar KS, Singh KK, Sharma SK, Tomar OK. Alternative tillage and crop residue management in wheat after rice in sandy loam soils of Indo-Gangetic plains. *Soil and Tillage Research*. 2006;88:242-252.
- Pal DK, Bhattacharyya T, Srivastava P, Chandran P, Ray SK. Soils of the Indo-

- Gangetic plains: their historical perspective and management. *Current Science*. 2009;96(9):1193-1202.
7. Verma SK, Singh SB. Enhancing of wheat production through appropriate agronomic management. *Indian Farming*. 2008;58(5): 15-18.
 8. Erenstein O, Laxmi V. Zero tillage impacts in India's rice-wheat systems: A review. *Soil and Tillage Research*. 2008;100:1-14.
 9. Usman K, Khalil SK, Khan MA. Impact of tillage and herbicides on weed density and some physiological traits of wheat under rice-wheat cropping system. *Sarhad Journal of Agriculture*. 2010;26(4): 475-487.
 10. Abbas G, Ali MA, Abbas Z, Aslam M, Akram M. Impact of different herbicides on broadleaf weeds and yield of wheat. *Pakistan Journal of Weed Science Research*. 2009;15:1-10.
 11. Moss S. Integrated Weed Management (IWM): Why are farmers reluctant to adopt non-chemical alternatives to herbicides. *Pest Management Science*. 2019;75:1205-1211.
 12. Kraehmer H, Laber B, Rosinger C, Schulz A. Herbicides as weed control agents: State of the art: I. Weed control research and safener technology: The path to modern agriculture. *Plant Physiology*. 2014;166:1119-1131.
 13. Deepak Kumar Yadav, SK Verma, RK Singh, Chandra Bhushan, SK, Rajpoot, Anurag Upadhyay and Prateek Kumar. Effect of crop establishment and weed management practices on soil properties in wheat (*Triticum aestivum* L.). *The Pharma Innovation Journal* 2023;12(6):4607-4609.
 14. Shyam R, Singh R, Singh VK. Effect of tillage and weed management practices on weed dynamics, weed seed bank and grain yield of wheat in rice-wheat system. *Indian Journal of Weed Science*. 2014;46(4):322-325.
 15. Upasani RR, Sheela B, Singh MK. Tillage and weed management in direct-seeded rice -wheat cropping system. *Indian Journal Agronomy*. 2014;59(2):204- 208.
 16. Singh RP, Verma SK, Kumar S, Lakara K. Impact of tillage and herbicides on the dynamics of broad leaf weeds in wheat. *International Journal of Agriculture, Environment and Biotechnology*. 2017;10 (6):643-651.
 17. Rani SP, Duary B, Priyatam S. Effect of tillage and weed management practices on weed control and yield in wheat. *International Journal of Current Microbiology and Applied Sciences*. 2020; 9(6):2328-2335.
 18. Choudhary M, Jat HS, Datta A, Yadav AK, Sapkota TB, Mondal S, Meena RP, Sharma PC, Jat ML. Sustainable intensification influences soil quality, biota, and productivity in cereal-based agro-ecosystems. *Applied Soil Ecology*. 2018;126:189-198.
 19. Bera T, Sharma S, Thind HS, Sidhu HS, Jat ML. Soil biochemical changes at different wheat growth stages in response to conservation agriculture practices in a rice-wheat system of north-western India. *Soil Research*. 2017;56(1):91-104.
 20. Patel VK, Pathak RK, Kumar A, Singh A, Samiksha Patel A. Effect of tillage and weed management practices on soil physico-chemical characteristics and wheat economics. *International Journal of Current Microbiology and Applied Sciences*. 2020;9(4):2096-2102.
 21. Islam S, Gathala MK, Tiwari TP, Timsina J, Laing AM, Maharjan S, Chowdhury AK, Bhattacharya PM, Dhar T, Mitra B, Kumar S, Srivastwa PK, Dutta SK, Shrestha R, Manandhar S, Sherestha SR, et al. Conservation agriculture based sustainable intensification: Increasing yields and water productivity for smallholders of the eastern gangetic plains. *Field Crops Research*. 2019;238:1-7.
 22. Yadav P, Singh RS, Kumar P, Maurya NK, Pal RK, Verma H. Effect of weed management practices on weed flora of wheat crop. *The Pharma Innovation Journal*. 2022;11(4):320-324.
 23. Jaiswal D, Reddy MD, Pandey G, Kumar A. Evaluation of herbicides for control of weeds in wheat. *International Journal of Current Microbiology and Applied Sciences*. 2020;9(5):1157-1167.
 24. Duary B, Jaiswal DK, Dash S, Sar K, Patel N. Effect of tillage and pre-mix application of herbicides on weed growth and productivity of late-sown wheat. *Indian Journal of Weed Science*. 2021;53(2):188-190.
 25. Kumar B, Hasanain M, Raza MDB, Yadav RB, Singh RK, Mishra R, Ahmad G, Singh

- D. Effect of weed management practices on nutrient uptake and productivity of wheat. *The Indian Journal of Agricultural Sciences*. 2022;92(3):405–407.
26. Li J, Xuan, J, Cai R. Wheat a popular cereal crop. *Field Crop*. 2020;3.

© 2023 Yadav et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/102769>