



Sustainable Management of Root- Knot Nematode (*Meloidogyne javanica*) on Brinjal

**Kavita Choudhary^{a*}, S. P. Bishnoi^a, Raju Dhayal^a,
B. S. Chandrawat^b, Anil kulheri^c and Hemraj Gurjar^a**

^a Division of Nematology, Rajasthan Agriculture Research Institute, Durgapura, Jaipur, SKNAU, Jobner, Jaipur, Rajasthan-303329, India.

^b Department of Nematology, SKN College of Agriculture, SKNAU, Jobner, India.

^c Department of Genetics and plant breeding, SKN College of Agriculture, SKNAU, Jobner, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i102624

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/104621>

Original Research Article

Received: 05/06/2023

Accepted: 09/08/2023

Published: 12/08/2023

ABSTRACT

Brinjal (*Solanum melongena*) is an important *Solanaceae* family vegetable, which is grown all around the world. It is affected by various fungal, bacterial, viral, nematode diseases but it is a major host for root-knot nematode, *Meloidogyne javanica*. Hence, for effective management of root-knot nematode experiment was conducted in field condition using the culture filtrate (3gm/kg soil) of bio-agents (viz., *Trichoderma viride*, *Trichoderma harzianum*, *Purpureocillium lilacinum*, *Pochonia chlamydosporia* and *Pseudomonas fluorescens*). The findings of the experiments revealed that, when compared to the control, all of the tested bio-agents considerably improved plant development characteristics and decreased nematode populations. Among the tested bio-agents *T. viride* was found most effective treatment with improve plant growth characters as well as in reducing nematode population over other treatment in field condition.

*Corresponding author: E-mail: kavitajakhar29@gmail.com;

Keywords: Brinjal; root-knot nematode; *Meloidogyne* spp; management; bio-agent.

1. INTRODUCTION

Brinjal or aubergine (*Solanum melongena* L.) is a popular and widely consumed vegetable throughout the tropical and sub-tropical regions of the world. It is one of the important rainy and winter season vegetable crops cultivated in various states of India. It is a good source of minerals and vitamins [1,2]. Brinjal is infected by number of diseases caused by fungi, bacteria, viruses and nematodes and these diseases cause extensive yield losses [3]. Nematodes are hidden enemy of plant. Among the nematodes, root-knot nematode, *Meloidogyne* spp. is an important pest of vegetables all around the world and causes severe damage to brinjal crop [4] in India, the root-knot nematode was first reported through Barber in 1901 [5] on tea (*Thea sinensis*) roots from Devala territory of Tamil Nadu, South India. In Rajasthan, it was first time reported by Arya in 1957 [6] from Jodhpur, while Yadav and Naik [7] found *Meloidogyne* spp. and its distribution in the soils of Rajasthan. The nematode infected plants have reduced root system with less feeder roots [8]. Overall, plant-parasitic nematodes cause 21.3% crop losses amounting to Rs. 102040 million (1.58 billion USD) annually; the losses in 19 horticultural crops were assessed at Rs. 50,225 million, while for 11 field crops it was estimated at Rs. 51,815 million. Among the vegetable crops comparatively more losses recorded in tomato (Rs. 6035.2 million) and brinjal (Rs. 3499.12 million) [9]. Management of nematode is more difficult as compare to other pathogen because nematodes attack mainly underground parts of plants [10]. The nematodes control mostly depended on synthetic nematicides [11]. Although, nematicides are efficient and fast acting, yet they are currently being reappraised as there are relatively unaffordable to many small-scale farmers. The potential negative effect on environment after prolonged use have led to a total ban or restrict use of most chemical nematicides and an urgent need for safe and more effective alternatives [12]. Biological control of plant parasitic nematodes appears the alternative strategies for management in the soil [13,14]. Bio-products containing a microorganism (bacterium, fungus, virus, protozoan or alga) an active ingredient often referred as microbial pesticides, are host specific and are having potential with regard to integrated pest management [15]. Fungi and bacteria are the most dominant among microorganism they are

soil-borne groups in natural soil ecosystem and some of them have shown great potential effect as biological control agents for root-knot nematodes [16]. Keeping this in view, the present investigations were undertaken to assess the efficacy of bio-agents in brinjal against root-knot nematode *Meloidogyne* spp.

2. MATERIALS AND METHODS

The experiment conducted for management of root-knot nematode *M. javanica* in brinjal using of bio-agents in field.

2.1 Raising Nursery and Transplanting

Brinjal variety marudhar was used in all experiments. Brinjal seedlings were raised in the 1m x 1m size plots and seeds were sown in line. 4-5 weeks old seedlings used for transplanting in main field for various experiments.

2.2 Bed Preparation and Sampling from Beds

For the experiment, beds measuring one metre in length and one metre in width were created. Before the seedling was transplanted, the soil was sampled to ascertain the initial nematode population.

2.3 Preparation of Culture Filtrates of Fungal and Bacterial Bio-Agents

The potato dextrose agar (PDA) for fungal agents and nutrient agar (NA) for bacterial agents were prepared, inoculated with respective bio-agents in 100 ml conical flasks followed by incubation at 30 °C in a shaker for 48 hrs. The cultures were centrifuged at 6000 rpm for 20-30 minutes. The supernatant was kept as a stock solution of cent per cent concentration. One ml of sterilized double distilled water was added on fully grown fresh mother culture of bio-agents *Trichoderma viride*, *Trichoderma harzianum*, *Purpureocillium lilacinum*, *Pochonia chlamydosporia* and *Pseudomonas fluorescens* than scraped with a spade to produce slurry.

2.4 Management of Root-Knot Nematode on Brinjal Using Bio-Agents

The experiment conducted in pure culture plot infested with root-knot nematode, *M. javanica* (2 J₂/g of soil). Talc-based formulation of *Trichoderma viride*, *Trichoderma harzianum*,

Purpureocillium lilacinum, *Pochonia chlamydosporia* and *Pseudomonas fluorescens* were added to soil each @ 3g per kg soil on different time interval i.e., at the time of transplanting, 20 and 40 days after transplanting. Six plants of brinjal were maintained in each plot. Each treatment was replicated four times. Untreated check was also maintained for comparison. Plants were harvested after 60 days of transplanting and the observations on plant growth characters and nematode reproduction were recorded. Utmost care was taken right from sowing to till harvest of experiment for proper growth and development of plants.

2.5 Statistical Analysis

After completion of experiment, data were statically analyzed for interpretation of finding. The critical deference was calculated for comparison of treatment for significant at 5 % level of significance. Summary table along with $SE_{m\pm}$ and CD were worked out and presented in chapter "Experimental Results".

3. RESULTS AND DISCUSSION

3.1 Management of Root-Knot Nematode in Brinjal Using Bio-Agents

An experiment was conducted to study the effect of bio-agents as soil application on management of root-knot nematode, *M. javanica* on brinjal. Bio-agents *T. viride*, *T.harzianum*, *P. lilacinum*, *P. chlamydosporia* and *P. fluorescens* were used @ 3g/kg soil. Plants were uprooted 60 days after transplanting and observation were recorded on plant growth characters and nematode reproduction.

3.2 Plant Growth Character

3.2.1 Shoot length

Data presented in table showed that all bio-agents significantly increased the shoot length over untreated check. Among the bio-agents the highest shoot length was recorded with *Trichoderma viride* (58.00cm) followed by *Purpureocillium lilacinum* (42.75cm) and *Pseudomonas fluorescens* (38.00cm) @ 3g/kg soil. While, minimum shoot length (23.25cm) recorded with untreated check.

3.2.2 Shoot weight

Data recorded in table showed that among the tested bio-agents the highest shoot weight was recorded with *Trichoderma viride* (53.00gm)

followed by *Purpureocillium lilacinum* (41.15 gm) and *Pseudomonas fluorescens* (33.20 gm) @ 3g/kg soil. While, minimum shoot weight (8.60gm) was recorded with untreated check. Application of bio-agents significantly increased the shoot weight as compared to untreated check.

3.2.3 Root length

Result presented in table among the bio-agents the highest root length was recorded with *Trichoderma viride* (29.47cm) followed by *Purpureocillium lilacinum* (28.52cm) and *Pseudomonas fluorescens* (27.27 cm) @ 3g/kg soil. While the minimum root length (8.95cm) was recorded with untreated check. Application of bio-agents significantly increased the root length over untreated check.

3.2.4 Root weight

Data recorded in table showed that bio-agents significantly increased the root weight as compared to untreated check. Among the bio-agents the highest root weight was recorded with *Trichoderma viride* (5.25 gm) followed by *Purpureocillium lilacinum* (4.82 gm) and *Pseudomonas fluorescens* (4.45 gm) @ 3g/kg soil. While, minimum root weight (2.37) was recorded with untreated check.

3.3 Nematode Reproduction

3.3.1 Number of galls/plant

Experimental result revealed in table showed that among the bio-agents the minimum number of galls per plant were recorded with *Trichoderma viride* (153.75) followed by *Purpureocillium lilacinum* (238.00) and *Pseudomonas fluorescens* (284.75) @ 3g/kg soil. While, maximum number of galls (397.50) were recorded with untreated check. Application of bio-agents significantly reduced number of galls per plant over untreated check.

3.3.2 Number of egg masses/plant

Data presented in table showed that application of bio-agents significantly decreased number of egg masses per plant. Among the bio-agents *Trichoderma viride* was observed best treatment with minimum number of egg masses per plant (95.25) followed by *Purpureocillium lilacinum* (107.00) and *Pseudomonas fluorescens* (120.50) @ 3g/kg soil. However, untreated check was observed least treatment with maximum number of egg masses per plant (189.75).

Table 1. Effect of bio-agent on plant growth and nematode reproduction in open field

| Treatment | Shoot length (cm) | Shoot weight (gm) | Root length (cm) | Root weight (gm) | No. of galls/ plant | No. of egg masses/ Plant | Number of eggs and larvae / egg mass | Nematode juvenile/ 200cc soil | Final nematode population |
|-------------------------|--------------------------|--------------------------|-------------------------|-------------------------|----------------------------|---------------------------------|---|--------------------------------------|----------------------------------|
| <i>T. viride</i> | 58.00 | 53.00 | 29.47 | 5.25 | 153.75 | 95.25 | 200.00 | 422.75 | 19469 |
| <i>T.harzianum</i> | 25.75 | 17.05 | 14.62 | 2.15 | 378.75 | 185.25 | 281.75 | 741.75 | 52939.25 |
| <i>P.lilacinum</i> | 42.75 | 41.15 | 28.52 | 4.82 | 238.00 | 107.00 | 212.25 | 539.50 | 23252.00 |
| <i>P.chlamydosporia</i> | 29.25 | 18.48 | 16.25 | 4.17 | 315.00 | 139.50 | 249.75 | 679.75 | 35522.00 |
| <i>P.fluorescens</i> | 38.00 | 33.20 | 27.27 | 4.45 | 284.75 | 120.50 | 231.25 | 590.50 | 28454.75 |
| Control | 23.25 | 8.60 | 8.95 | 2.37 | 397.50 | 189.75 | 344.75 | 1233.75 | 66654.00 |
| SEm± | 0.96 | 1.24 | 0.71 | 0.28 | 1.60 | 1.34 | 1.29 | 3.14 | 479.93 |
| CD 5% | 2.89 | 3.74 | 2.14 | 0.85 | 4.82 | 4.04 | 3.89 | 9.47 | 1446.37 |
| CV % | 5.31 | 8.70 | 6.81 | 14.72 | 1.09 | 1.92 | 1.02 | 0.90 | 2.55 |

* Average of four replications

*Dose = @3 gm/ kg soil at the time of transplanting followed by 20 DAT and 40 DAT (Day After Transplanting)

3.3.3 Number of eggs/egg mass

Data recorded in table showed that among all treatments the minimum number of eggs per egg mass were recorded with *Trichoderma viride* (200.00) followed by *Purpureocillium lilacinum* (212.25) and *Pseudomonas fluorescens* (231.25) @ 3g/kg soil. Whereas, the maximum number of eggs per egg mass (344.75) were recorded with untreated check. Application of bio-agents significantly decreased number of eggs per egg mass over untreated check.

3.3.4 Nematode juvenile/200cc soil

Data presented in table showed that application of bio-agents significantly decreased nematode juvenile/200cc soil compared to untreated check. Among the bio-agents the minimum number of nematode juvenile/200cc soil were recorded with *Trichoderma viride* (422.75) followed by *Purpureocillium lilacinum* (539.50) and *Pseudomonas fluorescens* (590.50) @ 3g/kg soil. While, the maximum number of juveniles (1233.75) were recorded with untreated check.



Plate 1. Over view of bio-agent treated field of brinjal



Plate 2. Management of root-knot nematode, *Meloidogyne javanica* on brinjal through bio-agents





Fig. 1. Effect of bio-agents on plant growth characters of brinjal in field condition






Fig. 2. Effect of bio-agents on nematode reproduction on brinjal in field condition

3.3.5 Final nematode population

Application of bio-agents decreased the final nematode population over untreated check. Results presented in table showed that among the bio-agents the minimum nematode population were recorded with *Trichoderma viride* (19469.00) followed by *Purpureocillium lilacinum* (23252.00) and *Pseudomonas fluorescens* (28454.75) @ 3g/kg soil. While, maximum nematode population (66654.00) were recorded with untreated check.

The results of experiment on management of root-knot nematode, *M. javanica* on brinjal using bio-agents showed that all bio-agents significantly increased the plant growth characteristics (shoot length, shoot weight, root length and root weight) and decreased nematode reproduction (number of galls/plant, number of egg masses / plant, number of eggs/ egg mass, nematode juvenile/ 200cc soil and final nematode population) over untreated check. *Trichoderma viride* was the bio-agent with the highest plant growth characteristics and the lowest nematode reproduction, followed by *Purpureocillium lilacinum* and *Pseudomonas fluorescens* @ 3g/kg soil. While maximum nematode proliferation and little plant growth features were seen in untreated checks. Similar findings were observed with Meghwal and Baheti [17] revealed that okra plant growth significantly enhanced with seed treatment of bio-agents. Maximum plant growth and minimum root-knot nematode infection was recorded when seeds were treated with *T. viride* followed by *P. lilacinus* and *P. chlamydosporia* each at 2% over untreated check. Meena et al. [18] evaluated bio-agents efficacy as seed treatment @ 4 g/kg seed as well as added to soil @ 4 g/kg soil against wilt disease complex by *M. incognita* and *F. oxysporum f. sp. lycopersici*. Among bio-control agent, *Trichoderma viride* was found significantly superior in reduction of nematode population and per cent disease incidence along with increased

in plant growth parameters followed by *Paecilomyces lilacinus*.

Kumhar et al. [19] showed that *Pochonia chlamydosporia* at 4 per cent was found most effective followed by *P. chlamydosporia* at 2 per cent and *G. fasciculatum* at 4 per cent to enhance plant growth of maize and to reduce the infection of *Heterodera zea*e on maize. Bhati et al. [20] exhibited that the bio-agents {*Paecilomyces lilacinus* (now known as *Purpureocillium lilacinum*), *Trichoderma harzianum*, *Pochonia chlamydosporia*, *Pseudomonas fluorescens* and *Glomus fasciculatum* (VAM)} reduced nematode reproduction and enhanced the plant growth of cucumber over untreated check. *Trichoderma viride*, at 5.0g per plant above an untreated control on cucumber plants in a polyhouse, showed the greatest reduction in root galls, egg masses, egg mass contents, and nematode population. *Paecilomyces lilacinus* and *Trichoderma harzianum* came in second and third, respectively.

4. CONCLUSION

The growth characteristics of the plant (shoot length, shoot weight, root length, and root weight) were significantly improved by the bio-agents, while nematode reproduction (number of galls/plant, number of egg masses/plant, number of eggs/egg mass, nematode juvenile/200cc soil, and final nematode population) was significantly reduced. *Trichoderma viride* was the bio-agent with the highest plant growth characteristics and the lowest nematode reproduction, followed by *Purpureocillium lilacinum* and *Pseudomonas fluorescens* @ 3g/kg soil.

ACKNOWLEDGEMENT

The authors are grateful to Head Of Department for providing facilities and support for carrying out the research experiment in Division of

Nematology at RARI Durgapura and Department of Plant Pathology at SKNAU Jobner Jaipur.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Russo VM. Cultural methods and mineral content of eggplant (*Solanum melongena*) fruit. *Journal of the Science Food and Agriculture*. 1996;71(1):119-123.
2. Sadilova E, Stintzing FC, Carle R. Anthocyanins, colour and antioxidant properties of eggplant (*Solanum melongena* L.) and violet pepper (*Capsicum annuum* L.) peel extracts. *Zeitschrift für Naturforschung*. 2006;61(7-8):527-535.
3. Zeerak N, Iqbal Z, Kamran M, Iftikhar Y, Arshad M, Abbas M, Javed N, Bashir S, Rehman A. Root-knot nematodes associated with eggplant in different localities of district Sargodha-Pakistan and impact of pasteuria isolates on development of meloidogyne incognita. *International Journal of Biosciences*. 2017;11(4):107-115.
4. Kamran M, Anwar SA, Javed M, Khan SA, Sahi GH. Incidence of root-knot nematodes on tomato in Sargodha, Punjab, Pakistan. *Pakistan Journal of Nematology*. 2010;28:253-262.
5. Barber CA. A tea ell worm disease in South India. *Dept. Land Record, Madras Agricultural Branch 2. Bull. No. 45*; 1901.
6. Arya HC. Root-knot diseases of tomatoes in Jodhpur. *Science and Culture*. 1957;22:391-393.
7. Yadav BS, Naik SMP. Nematodes associated with economic plants of South East Plateau of Rajasthan. *Labdev Journal of Science and Technology*. 1966;4:184-186.
8. Anwar S, Mckenry MV. Incidence and reproduction of meloidogyne incognita on vegetable crop Genotype. *Pakistan Journal of Zoology*. 2010;42:135-141.
9. Kumar Vinod, Khan MR, Walia RK. Crop loss estimations due to plant-parasitic nematodes in major crops in India. *The National Academy of Sciences, India*; 2020.
10. Sikora RA, Fernández E. Nematode's parasites of vegetables. In: Liuc M, Sikora RA, Bridge J. (Eds.), *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*. CAB International, Wallingford (GBR). 2005:319–392.
11. Akhtar M, Malik A. Roles of organic soil amendments and soil organisms in the biological control of plant-parasitic nematodes. A review. *Bioresource Technology*. 2000;74:35-47.
12. Zukerman BM, Esnard J. Biological control of plant nematodes current status and hypothesis. *Japanese Journal of Nematology*. 1994;24:1–13.
13. Agyarko K, Asante JS. Nematode dynamics in soil amended with neem leaves and poultry manure. *Asian Journal of Plant Sciences*. 2005;4:426-428.
14. Collange B, Navarrete M, Peyre G, Mateille T, Tchamitchian M. Root-knot nematode (meloidogyne) management in vegetable crop production. The challenge of an agronomic system analysis. *Crop Protection*. 2011;30:1251–1262.
15. Arora R, Battu GS, Ramakrishnan N. Microbial pesticides. Current status and future outlook. In: Dhaliwal GS, Singh B. (Eds.), *Pesticides and Environment*. Commonwealth Publishers, New Delhi. 2000:344–395.
16. Kerry BR. Rhizosphere interactions and the exploitation of microbial agents for the biological control of plant-parasitic nematodes. *Annual Review of Phytopathology*. 2000;38:423–441.
17. Meghwal SK, Baheti BL. Management of root-knot nematode, meloidogyne incognita in okra (*Abelmoschus esculentus* L.) through bio-agents. *Indian Journal of Nematology*. 2017;47(2):192 - 196.
18. Meena P, Chandrawat BS, Ahir RR. Bio-agents for management of wilt complex in tomato by meloidogyne incognita and *Fusarium oxysporum* f. sp. *Lycopersici*. *Ann. Pl. Protec. Sci*. 2020;28(2):160-162.
19. Kumhar RN, Baheti BL, Chandrawat BS, Gurjar Om Prakash. Seed treatment with bio-agents: An environmental suitable approach for management of maize cyst nematode, heterodera zaeae on maize. *Current Nematology*. 2018;29(1,2): 1-5.

20. Bhati SS, Baheti BL, Singh Ishwar, Chandrawat BS. Impact of bio-agents against meloidogyne incognita infecting cucumber in poly-house. Biological Forum – An International Journal. 2022;14(2): 1481-1487.

© 2023 Choudhary et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104621>