



Advances in Research

16(6): 1-7, 2018; Article no.AIR.44315
ISSN: 2348-0394, NLM ID: 101666096

Evaluation of Growth Parameters and Yield Parameters of Three Cultivars of Potato Grown in Hamelmalo, Eritrea

Mulue Girmay¹, V. K. Sharma¹ and G. Sethumadhava Rao^{1*}

¹Department of Plant Protection, Hamelmalo Agricultural College, Keren, Eritrea.

Authors' contributions

This work was carried out in collaboration between all authors. The concept of this project designed and preformed by author MG. It was supervised by author VKS and the research work analysed and managed the literature review by author GSR. All the authors approved the final draft.

Article Information

DOI: 10.9734/AIR/2018/44315

Editor(s):

(1) Dr. Slawomir Borek, Department of Plant Physiology, Adam Mickiewicz University, ul. Umultowska, Poland.

Reviewers:

(1) S.S. Kushwah, College of Horticulture, RVS Krishi Vishwa Vidyalaya, India.

(2) Md. Mijanur Rahman, Banglades.

Complete Peer review History: <http://www.sciedomain.org/review-history/26684>

Original Research Article

Received 09 July 2018
Accepted 30 September 2018
Published 19 October 2018

ABSTRACT

The experiment was conducted at Hamelmalo Agricultural College farm station from September 2014 to January 2015. The design of the experiment was randomized complete block in triplicate. The growth parameters (plant height, stem number, leaf number at 30 and 60 days after planting) and yield variables were recorded in ton/ha. Among the cultivars; *Zafira* scored the highest number of plants per plot (32.6 plants/plot) as compared to *Picasso* (25.53 plants/plot) and *Ajiba* (22.07 plants/plot). The cultivars *Ajiba* and *Zafira* scored the highest vegetative growth. The highest growth and yield parameters were obtained from cultivars *Ajiba* (29.33 ton/ha) followed by *Zafira* (27.99 ton/ha) though both cultivars had the highest incidences of blights and collar rot diseases. The correlation analysis indicated there was no significant association among insect pests and yield ha⁻¹. However, the growth parameters showed significant correlations with yield ha⁻¹ at 0.01 level of significance.

Keywords: Stand count; plant height; stem number; leaf number; marketable and unmarketable potato production.

*Corresponding author: E-mail: sethumadhava.g@gmail.com;

1. INTRODUCTION

Potato (*Solanum tuberosum* L.) is nutritionally important vegetable due to its edible energy and edible protein. It produces more quantity of dry matter, edible energy and edible protein in lesser duration of time as compared to cereals like rice and wheat. According to Donnelly D and Kubow S [1], potato contains different nutrients like carbohydrates 10-30%, proteins 1-1.5% and lipids and dietary fibre 1-2% of the total fresh mass and also it is a source of different minerals, vitamins, and other phytochemicals. It is an important crop which is vegetatively propagated over the largest area in the world [2]. The produce of this solanaceous vegetable is supplementing meat and milk products by lowering energy intake and also by reducing food cost [3]. Though South America is known to be native of potato, it is grown in more than 100 countries globally. China ranks first, followed by Russia and India. Worldwide potato production in 2009 was harvested as 329.58 million tons from 18.65 million hectares. Highest production per hectare was found from the central Europe (43.94 ton/ha) followed by North America (42.43 ton/ha); whereas lowest production was from Africa 9.98 ton/ha [4].

Potato is one of the major vegetable crops after tomato and onion in Eritrea. Even though highland areas are suitable for potato cultivation and its total production is very low as compared to the demands of the consumers. Besides potato cultivars differ markedly in their yielding ability. Potato variety/cultivar experiments are important since potatoes are very sensitive to their growing environment and not all varieties/cultivars are adapted to all potato growing areas. Farmers are more interested in the cultivars that produce consistently high yields under their growing conditions. Total yield, number and weight of tubers per plant and average tuber weight and tuber quality such as specific gravity and starch content showed large variations among potato cultivars [5]. The germplasms well adapted to a particular set of conditions will be best suited for a locality for cultivation. A mixture of many locally grown varieties (*Carnetiom* and *Shashemene*) and cultivars (*Picasso*, *Condor*, *Ajaba*, *Zafira*, *Cosmos*, *Spunta*, etc) are used in Eritrea. *Picasso* and *Condor* are top yielding varieties however due to their pinkish skin colour; they are fetching lesser price and have less demand in the market as compared to *Zafira* and *Ajiba*

which are ranking third and fourth in production, and first and second in their market values [6].

There are no fixed development stages in potatoes as these are influenced by varieties, cultivars, stem size, soil fertility, weather conditions etc. Unlike another crops, phenological stages of potato overlap with each other making it difficult to distinguish between stages. For example, sometimes during the early growth stage, developing tubers have already begun to grow from the roots. According to Bohl et al. [7] of the seed piece size, physiologically old seeds generally produce more stems per seed piece than less-aged seed pieces. Potatoes planted from tuber seed have five stages sprout development, vegetative growth, tuber initiation, tuber bulking and tuber maturation stage [8].

In Eritrea; the total production area and yield of 2632 ha and 17.1 mt/ha respectively, whereas, the total production area and yield of potato in Zoba Anseba is 71 ha and 6 mt/ha respectively [9]. The crop is used as a food for consumption by the traditional farmers but in some cases it is considered as a source of income for the family. But its production is much lower as compared with other countries of the world. Therefore, this study was compulsive on the evaluation of growth performance and yielding potentiality of three cultivars grown in Hamelmalo of Zoba Anseba.

2. MATERIALS AND METHODS

2.1 Experimental Site

The experiment was conducted in the experimental fields of Hamelmalo Agricultural College in sub zoba Hamelmalo of Zoba Anseba which is located around 13 km from Keren to the north in Eritrea. It has the altitude 1280 m above sea level with 15° 55' 12.92" N latitude and 38° 27' 46.9"E longitudes. The study area receives an average annual rain fall of 459 mm and has an average temperature of 24°C. The area has sandy loam soil with pH of 6-7. Prior to the planting of the potato crop, soil samples from different areas of the experimental blocks were taken with the help of an auger at 0-30 cm depth, for the analysis which was carried out at National Agricultural Research Institute Halhale, Eritrea. Composite and representative air dried soil samples were analysed for its physico-chemical properties. The most prevalent component of

loamy soil is sand, but this is balanced with clay and silt. Loamy soil usually has the good characteristics of clay and sand; it has a friable nature, the ability to hold water well and is fertile. This soil is often found in mountainous regions or on the slopes of volcanoes [10].

2.2 Experimental Design, Field Preparation and Agronomical Practices

The experimental trial was conducted from 1st September 2014 to 15th January 2015 with three potato cultivars in RCBD experimental design under irrigation and all necessary cultural practices. The experimental plot size was 3m x 2.8m, with 0.7m space between rows, 0.3m between plants, 0.8m space between plots and 1m between replications. The temperature ranges from 16°C to 38°C with the hottest month being May and the coldest month in January. The total rainfall ranges from 465 to 673mm with a mean annual rainfall of 537.6 mm. The rain starts in June and lasts until the end of September. July and August are the months with the highest rainfall [11]. The experimental field was prepared with normal agronomic practices, FYM @ 50 quintals/ha was mixed in the soil before ploughing. In addition the fertilisers DAP and Urea were used before and after planting at the rate of 120 kg and 225 kg per hectare respectively. The doses of potassium (K₂O) 150 kg/ha were applied through band method at the time of planting to each experimental plot. The field was leveled and then the plots were made as per the requirement of the experimental design. Sowing of the potato tubers was done in the spacing of 30 cm and 70 cm between plants and rows respectively and the seed were sown in the depth of 10-15 cm. Earthing up was done two times at 30 and 45 days after emergence of the crop. The experimental units (plots) were irrigated and weeds were removed at an interval of 5 and 15 days respectively. All the above mentioned activities were done for the three cultivars uniformly.

2.3 Experimental Data of Growth Studies

2.3.1 Growth parameters

The growth parameters (stand count, plant height, stem number, leaf number and yield) deliberate and simple random sampling was used. Samples were taken from the two middle rows. In Stand count, the numbers of plants per

plot (3m x 2.8m) were taken from each of the experimental plots after the date of emergence of the crop. The available plants per plot were counted and an average number of plants per plot was calculated. Plant height (in centimeters) was measured with the help of meter scale from soil surface up to top growing point. Three plants from each plot were taken and measured with the help of meter scale and the average height of the plants was calculated. The data were recorded at the age of 30 and 60 days from the date of planting. The number of stems from each sample plant (three plants) was counted and the average stem number per plant was calculated for each plot. To determine the average leaf number a sample of ten stems from 5 sampled plants were taken from each plot. Total numbers of leaves from the sample stems were divided by the sample number and an average number of leaves per stem was calculated.

2.3.2 Yield and yield parameters

The yield was measured by weighing the tubers of 10 sample plants from each plot in kg using a weighing balance and divided in to the number of sampled plants to get the average tuber weight per plant [12]. Finally, the yield in kg per plant was converted to tons per hectare. In addition to that tubers from the sample, plants were graded as marketable and unmarketable and weighed separately. Data recorded on different parameters were analysed using statistical software Genstat and mean comparison were performed using the least significant difference (LSD) at 5% level of significance.

3. RESULTS AND DISCUSSION

3.1 Growth Studies

3.1.1 Stand count

Plant count was done for each of the three cultivars to determine the capacity of the cultivars to with stand the given environment. The table of analysis of this research showed a highly significant difference in the stand count of the plants per plot due to cultivars influence at 5% level of significance. Among the cultivars; *Zafira* scored the highest number of plants per plot (32.6 plants/plot) as compared to *Picasso* (25.53 plants/plot) and *Ajiba* (22.07 plants/plot) (Fig. 1).

The lowest value of the *Ajiba* cultivar was because of its much softened seeds of potatoes. Moreover, the highest sprouting accelerates the

early emergence of the plant, more rotting was observed once the seeds were inserted in the soil. In addition to that the lower number of plants per plot in *Ajiba* was due to the exhaustion of the stored food before sprouts emerge as the potato seed were sown in the depth 10-15cm. This result is supported by Chehaibi et al. [13] it was reported that shallow planting is preferred in wet and heavy soils because in such soils deep planting of the tubers may lead to exhaustion of stored food before the sprouts emerge above the soil. It is therefore, suggested that the cultivars which are highly sprouted before planting should be shallow planted.

3.1.2 Plant height at 30 days after planting

There was highly significant difference among the cultivars with regard to plant height at 5% level of significance. The cultivar *Ajiba* scored the maximum plant height 32.6cm followed by *Zafira* 31.67cm where as the lowest plant height value was obtained from *Picasso* cultivar 22.69cm. The cultivars *Ajiba* and *Zafira* were found at par but significantly different from the *Picasso* cultivar (Fig. 1).

3.1.3 Stem number per plant at 30 days after planting

The analysis of variance provided a highly significant difference among the cultivars at 5% level of significance because of the cultivar effect. The cultivars *Ajiba* and *Zafira* scored the highest number of stems per plant (3.8) where as the *Picasso* showed less number of stems per plant (2.78) as compared the above listed two cultivars (Fig. 1).

The more number of stems in the *Ajiba* and *Zafira* cultivars was due to the aged tubers (stored for long) as compared the third cultivar *Picasso* which had the lower number of stems per tubers. These findings are supported by the results of [7] who mentioned that regardless of the seed piece size and physiologically old seeds generally produced more stems per seed piece than less-aged seed pieces.

3.1.4 Leaf number per stem at 30 days after planting

Highly significant variation was observed in number of leaves per stem in the three different potato cultivars at 5% level of significance due to cultivar effects. The maximum number of leaves per stem was recorded from the *Zafira* cultivar

(14.69) where it showed a significantly higher number of leaves as compared to *Ajiba* (12.66) and *Picasso* (11.69) which was found at par with respect to the number of leaves per stem (Fig. 1).

3.1.5 Plant height at 60 days after planting

Plant height was recorded at 60 days after planting. The analysis showed that the cultivars had highly significant difference due to cultivar effect with regard to the plant height of the cultivars at 5% level of significance. The cultivars *Ajiba* and *Zafira* showed the maximum mean plant height 34.39 cm and 34.29 cm respectively which showed high variation from *Picasso* cultivar 25.34 cm (Fig. 2). The two cultivars (*Ajiba* and *Zafira*) were found at par, behave similarly as it was in case of plant height at 30 days after planting.

3.1.6 Stem number per plant at 60 days after planting

It was observed that all the cultivars differed significantly due the effect of variety with regard to the stem number at 5% level of significance. The cultivar *Ajiba* scored the highest stem number 4.84 per plant followed by *Zafira* 3.91 per plant. The lowest stem number was recorded from the cultivar *Picasso* 2.81 per plant.

3.1.7 Leaf number per stem at 60 days after planting

Highly significant variation was observed in the mean number of leaves per stem in the potato varieties/cultivars at 5% level of significance due to the effect of the variety on the number of leaves. The maximum numbers of leaves per stem were obtained from the *Zafira* cultivar (17.56) which showed a significant difference from the other two cultivars. This is because of its vigorous vegetative growth. *Ajiba* and *Picasso* scored the minimum number of leaves per stem (16.02) and (15.31) respectively.

The present finding is supported by the work of [14], where *Zafira* cultivar had maximum leaf number when planted in the month of September. Younger seed tubers produced higher foliage than older seed tubers. These results are in agreement with the findings of [2], but the finding is in opposite for *Picasso* cultivar as it had lower leaf number.




Fig. 1. Growth studies of three cultivars of potato grown in Hamelmalo at 30 days interval

3.2 Yield Parameters

3.2.1 The effect of variety on total yields

The effects of cultivar on total yield were analysed and the yield was recorded in ton per hectare to compare the response of the three cultivars. The cultivars showed highly significant difference at 5% level of significance in yield production. This result was in agreement with the report given by Santerre et al. [15]. Maximum yield was obtained from the *Ajiba* (29.33 ton/ha) followed by *Zafira* (27.99 ton/ha). More or less similar results from *Zafira* cultivar (26 ton/ha) were reported from [16]. Based on the field observations it was noticed that the *Ajiba* and *Zafira* cultivars maintained vigorous vegetative growth. Both the cultivars found at par but showed significantly higher yields to the *Picasso* cultivar which had minimum production (19.49ton/ha). This was probably because of the late emergence and poor performance of the *Picasso* cultivar in the study area. Even though *Ajiba* cultivar gave the high production, the production was found lower as compared to the result reported by Belay T. [14] which was 73.5 ton /ha (Fig. 2).

3.2.2 The effect of variety on marketable yields

The yield was recorded in tons per hectare to know the response of the three cultivars and

analysed to assess the effects of cultivar on total marketable yield. The cultivars showed highly significant difference due to cultivar effects at 5% level of significance. The maximum marketable yield was obtained from the *Ajiba* (26.61 ton/ha) followed by *Zafira* (26.16 ton/ha). Both the cultivars behaved equally and showed a significant difference from the *Picasso* cultivar which gave minimum production (17.77 ton/ha). In spite of the some disease incidences which were noticed during the growth period, the *Ajiba* and *Zafira* cultivars gave maximum production as compared the *Picasso* cultivar. This might be due to the high vegetative growth of these cultivars in the field condition and their early emergence to *Picasso* which helped them to become resistant to biotic and abiotic conditions.

3.2.3 The effect of variety on unmarketable yields

The results showed that there was significant difference in unmarketable yield of the cultivars at 5% level of significance. The maximum unmarketable yield was recorded from the *Ajiba* cultivar (2.72 ton/ha) which was significantly different from the cultivars *Zafira* (1.84 ton/ha) and *Picasso* (1.724 ton/ha). The high unmarketable produce from *Ajiba* cultivar was due to the high percentage incidences of collar rots and blights which were occurred in the crop (Fig. 2).




Fig. 2. Yield parameters of three cultivars of potato grown in Hamelmalo

4. CONCLUSIONS

During the experimental period, the cultivars showed satisfactory vegetative growth, but their productions were not as much as anticipated. This might probably be due to the variation in the ecological conditions such as soil types, soil condition and time of planting (photoperiod). The cultivars *Ajiba* and *Zafira* scored the highest vegetative growth (plant height and stem number) at 30 days after planting where the maximum leaf number was recorded from *Zafira* cultivar at 30 and 60 days after planting. Both the *Ajiba* and *Zafira* cultivars gave the highest plant height, whereas maximum stem number was obtained from *Ajiba* cultivar at 60 days after planting. The cultivars showed high variation in yield production.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Donnelly D, Kubow S. Role of potato in human health; 2011. Available:http://www.agrireseau.qc.ca/pdt/documents/Donnelly_Kubow.pdf [Accessed 25th of January, 2014]
2. Struik PC, Wiersema SG. Seed potato technology Wageningen Perts, The Netherlands; 1999.
3. Bhajantri S. Production, processing and marketing of potato in Karnataka-an economic analysis; 2011. Available:<http://www.ageconsearch.umn.edu/bitstream/113945/2/my%20thesis.pdf> [Accessed 26th of June, 2014]
4. Faulkner G. Essential trends in World Potato Markets; 2012. Available:http://www.europatatscongress.eu/docs/Taormina/Europatat_Congress_Guy_Faulkner.pdf [Accessed 28th of June, 2014]
5. Samih A, Azmi A, Ayed A, Yasin A, Nazeir H. Impact of cultivar and growing season on potato under center pivot irrigation system. World Journal of Agricultural Sciences. 2011;7(6):718-721.
6. Anonymous. Vegetable crops research program. Annual report of National Agricultural Research Institute. Halhale, Eritrea; 2009.
7. Bohl WH, Nolte P, Kleinkopf GE, Thornton MK. Potato seed management, seed size and age; 2000. Available:<http://www.cals.uidaho.edu/edcomm/pdf/CIS/CIS1031.pdf> [Accessed 20th of December, 2014]
8. Rana MK. Olericulture: Kalyani: Ludhiana, India; 2008.
9. MoA, Potato production. Head office Asmara, Eritrea; 2012.
10. MoA, Statistical report. Asmara, Eritrea; 2005.

11. Anonymous. Annual average rainfall report zoba Anseba branch, Ministry of Agriculture; 1997.
12. Tantowijoyo W, van de Fliert E. All about potatoes, an Ecological Guide to Potato Integrated Crop Management; 2006.
Available:http://sresearch.cip.cgiar.orgtype3webfileadminicmtoolboxICMToolboxIntegrated_crop_management_All_about_potatoes_complete_EN_0602.pdf. [Accessed 1st of December, 2014]
13. Chehaibi S, Hamdi W, Abrougui K. Effects of planting depth on Agronomic Performance of two potato varieties grown in the Sahel region of Tunisia. Development and Agricultural Economics Journal. 2013;5(7):272-276.
14. Belay T. Effect of planting dates and varieties on growth and yield of potato (*Solanum tuberosum* L.) under hamelmalo agro-climatic conditions. MSc thesis, Hamelmalo Agricultural College, Eritrea; 2014.
15. Santerre C, Cash J, Chase R. Influence of cultivar, harvest-date and soil N on sucrose, specific gravity and storage stability of potatoes grown in Michiga; 1986.
Available:[http:// asrc. am/ uploads/ media/m-4.pdf](http://asrc.am/uploads/media/m-4.pdf) [Accessed 14 of November, 2014]
16. NARI, The potato variety, *Zafira*. Halhale, Eritrea; 2011.

© 2018 Girmay et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/26684>