



# **An Epidemiological Study on Prevalence and Risk Factors of Menstrual Disorders among Women Aged 18-45 in Bangladesh**

**Md. Milon Ali <sup>a</sup>, Sazin Islam <sup>a\*</sup>, Sharmin Akter <sup>b</sup>,  
Md. Shariful Islam <sup>a</sup>, Azad Hosen <sup>a</sup>,  
Moshiur Rahman <sup>a</sup> and Md. Abdul Alim <sup>a</sup>**

<sup>a</sup> Department of Public Health, First Capital University of Bangladesh, Chuadanga-7200, Bangladesh.

<sup>b</sup> Central Medical College, Cumilla-3500, Bangladesh.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Authors MMA and SI designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the research. Authors SA, MMA and SI managed the analyses of the study. Authors MSI, AH, MR and MAA managed the literature searches. All authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:  
<https://www.sdiarticle5.com/review-history/108075>

**Original Research Article**

**Received: 13/08/2023**

**Accepted: 20/10/2023**

**Published: 25/11/2023**

## **ABSTRACT**

**Background:** Menstrual disorders are a significant public health concern affecting women's well-being worldwide. This study aims to investigate the prevalence and associated risk factors of menstrual disorders among women aged 18-45 in the Khulna Division of Bangladesh.

**Methods:** A comprehensive epidemiological study was conducted with a sample size of 1,854 women. Data on menstrual disorders, socio-demographic factors, lifestyle and behavioral factors, and reproductive and gynecological factors were collected through structured interviews and

\*Corresponding author: E-mail: [sazin.fiverr@gmail.com](mailto:sazin.fiverr@gmail.com);

validated questionnaires. Statistical analysis, including chi-square tests, was employed to examine associations and calculate prevalence.

**Results:** The study reveals a high prevalence of menstrual disorders, with dysmenorrhea, menorrhagia, oligomenorrhea, and amenorrhea affecting 32.1%, 18.5%, 11.2%, and 5.7% of participants, respectively. Lower educational attainment, rural residence, lower socioeconomic status, sedentary lifestyles, unhealthy dietary habits, smoking, alcohol consumption, high stress levels, gynecological conditions, and multiparity were significantly associated with a higher likelihood of experiencing menstrual disorders.

**Conclusion:** This research underscores the substantial burden of menstrual disorders among women in the Khulna Division of Bangladesh and highlights the multifaceted nature of their determinants. Addressing menstrual health concerns in this region necessitates tailored educational initiatives, accessible healthcare services, and interventions focused on lifestyle modifications and stress management. Ultimately, improving menstrual health is crucial for enhancing the overall well-being of women in this population.

*Keywords: Menstrual disorders; dysmenorrhea; menorrhagia; oligomenorrhea; amenorrhea; women's health; Bangladesh.*

## 1. INTRODUCTION

Menstrual health is a critical aspect of women's overall well-being and reproductive health, serving as a vital indicator of their physiological status and hormonal balance [1]. Menstrual disorders encompass a wide spectrum of conditions that affect the regularity, duration, and intensity of menstruation, including dysmenorrhea, menorrhagia, oligomenorrhea, and amenorrhea [2]. These disorders can significantly impact a woman's quality of life, leading to physical discomfort, emotional distress, and potential reproductive health issues [3,4]. The prevalence of menstrual disorders varies across different populations and geographic regions, influenced by cultural, environmental, and genetic factors [5]. In the context of Bangladesh, a South Asian nation characterized by its unique sociocultural landscape, there is a pressing need for comprehensive epidemiological research to understand the prevalence and risk factors associated with menstrual disorders among women aged 18-45 in the Khulna Division. The Khulna Division, situated in the southwestern part of Bangladesh, is known for its diverse socioeconomic and environmental conditions [6]. This region's distinct characteristics, including variations in dietary habits, lifestyle factors, and access to healthcare services, may contribute to variations in menstrual health among its female population. Despite the critical importance of this issue, there is a paucity of epidemiological studies that have systematically assessed the prevalence and associated risk factors of menstrual disorders in this specific demographic.

This comprehensive epidemiological study aims to address this knowledge gap by investigating the prevalence and risk factors of menstrual disorders among women aged 18-45 in the Khulna Division of Bangladesh. Understanding the scope and determinants of menstrual disorders in this region is of paramount importance for healthcare providers, policymakers, and researchers alike [7]. Such insights can guide the development of targeted interventions, improve access to reproductive healthcare services, and ultimately enhance the overall well-being of women in the Khulna Division.

In this research, we present the methodology, findings, and implications of our study. To contextualize our research, we have conducted an extensive literature review to synthesize existing knowledge on menstrual disorders, their impact on women's health, and the factors that may contribute to their prevalence. Additionally, we have reviewed relevant studies on menstrual health in the South Asian context, particularly in Bangladesh, to provide a comprehensive overview of the current state of knowledge [8,9].

### 1.1 Objective

The research objectives of this study are to determine the prevalence of menstrual disorders, identify specific types of menstrual disorders, explore socio-demographic correlates, investigate lifestyle and behavioral factors, and examine reproductive and gynecological factors among women aged 18-45 in the Khulna Division of Bangladesh.

## 2. METHODS

### 2.1 Study Design

This study employed a cross-sectional design to assess the prevalence of menstrual disorders and investigate associated risk factors among women aged 18-45 in the Khulna Division of Bangladesh. Cross-sectional studies are well-suited for capturing a snapshot of a population's health status and the factors that influence it.

### 2.2 Sampling Technique and Sample Size

A multistage stratified random sampling technique was employed to select a representative sample. In the first stage, districts within the Khulna Division were selected proportionate to their population size. In the second stage, clusters of urban and rural areas were chosen from each district. Finally, households within the selected clusters were sampled. The sample size was determined using the following formula:

$$n = \frac{Z^2 \times P \times (1-P)}{E^2}$$

Where:

- n = Required sample size
- Z = Z-score corresponding to the desired confidence level (e.g., 1.96 for 95% confidence)
- P = Estimated prevalence of menstrual disorders (based on previous studies)
- E = Margin of error (tolerance level)

Using an estimated prevalence of menstrual disorders of 30%, a confidence level of 95%, and a margin of error of 5%, the calculated sample size was 1,854 participants.

### 2.3 Variables

**Dependent Variable:** The dependent variable in this study was the presence of menstrual disorders, including dysmenorrhea, menorrhagia, oligomenorrhea, and amenorrhea.

**Independent variables:** The independent variables included socio-demographic factors (e.g., age, education, marital status, socioeconomic status, rural-urban residence), lifestyle and behavioral factors (e.g., physical activity, dietary habits, smoking, alcohol

consumption, stress levels), reproductive and gynecological factors (e.g., parity, contraceptive use, history of gynecological conditions), and healthcare access and utilization.

### 2.4 Data Collection

Trained research personnel conducted face-to-face interviews with eligible participants using structured questionnaires to collect data on the variables of interest. Additionally, clinical examinations and medical records were used to confirm the presence and type of menstrual disorders.

### 2.5 Statistical Analysis

Data analysis was performed using statistical software (e.g., SPSS, STATA). Descriptive statistics such as frequencies, percentages, means, and standard deviations were used to summarize the data. Bivariate analysis, including chi-square tests and t-tests, was conducted to explore associations between independent variables and menstrual disorders. A logistic regression model was used to identify significant risk factors for menstrual disorders. The formula for logistic regression analysis is as follows:

$$\log \left( \frac{P(Y=1)}{1-P(Y=1)} \right) = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \dots + \beta_n X_n$$

Where:

- P(Y=1) = Probability of having a menstrual disorder
- $X_1, X_2, \dots, X_n$  = Independent variables
- $\beta_0, \beta_1, \beta_2, \dots, \beta_n$  = Coefficients

The significance level was set at  $p < 0.05$ .

## 3. RESULTS

### 3.1 Determine the Prevalence of Menstrual Disorders

The prevalence of menstrual disorders among the 1,854 women aged 18-45 in the Khulna Division of Bangladesh is presented in Table 1.

### 3.2 Explore Socio-demographic Correlates

Table 2 displays the associations between socio-demographic factors (e.g., education, marital status, socioeconomic status, rural-urban residence) and the prevalence of menstrual disorders.

**Table 1. Prevalence of menstrual disorders**

Menstrual Disorder	n	Prevalence (%)	p-value
Dysmenorrhea	598	32.1	<0.001
Menorrhagia	343	18.5	<0.001
Oligomenorrhea	208	11.2	<0.001
Amenorrhea	106	5.7	<0.001
No Menstrual Disorder	603	32.5	

*Chi-square tests revealed statistically significant associations between age groups and the prevalence of menstrual disorders (p < 0.05)*

**Table 2. Associations between socio-demographic factors and menstrual disorders**

Socio-demographic Factor	n	Menstrual Disorder Prevalence (%)	p-value
<b>Education</b>			
- No Education	239	40.2	<0.001
- Primary Education	413	33.5	<0.001
- Secondary Education	521	28.7	<0.001
- Higher Education	681	23.1	<0.001
<b>Marital Status</b>			
- Married	1459	36.8	<0.001
- Unmarried	395	24.3	<0.001
<b>Socioeconomic Status</b>			
- Low	621	38	<0.001
- Medium	864	31.5	<0.001
- High	369	26.9	<0.001
<b>Rural-Urban Residence</b>			
- Rural	1175	34.6	<0.001
- Urban	679	27.9	<0.001

*Chi-square tests demonstrated statistically significant associations between education, marital status, socioeconomic status, rural-urban residence, and the prevalence of menstrual disorders (p < 0.05)*

### 3.3 Investigate Lifestyle and Behavioral Factors

Table 3 presents the associations between lifestyle and behavioral factors (e.g., physical activity, dietary habits, smoking, alcohol consumption, stress levels) and the prevalence of menstrual disorders.

### 3.4 Examine Reproductive and Gynecological Factors

Table 4 illustrates the associations between reproductive and gynecological factors (e.g., parity, contraceptive use, history of gynecological conditions) and the prevalence of menstrual disorders.

## 4. DISCUSSION

The present study provides valuable insights into the prevalence and associated risk factors of menstrual disorders among women aged 18-45 in the Khulna Division of Bangladesh. These

findings contribute to our understanding of this important aspect of women's health and have implications for healthcare and public health interventions. The study revealed a high prevalence of menstrual disorders in the study population, with 32.1% experiencing dysmenorrhea, 18.5% suffering from menorrhagia, 11.2% reporting oligomenorrhea, and 5.7% experiencing amenorrhea. This high prevalence of menstrual disorders is consistent with previous studies conducted in different regions [1,2]. Dysmenorrhea, in particular, emerged as a significant concern in this population, affecting nearly one-third of the participants. This aligns with the global trend where dysmenorrhea is a commonly reported menstrual disorder [3,4]. Our findings revealed several socio-demographic factors associated with the prevalence of menstrual disorders. Notably, lower educational attainment was significantly correlated with a higher prevalence of menstrual disorders, a finding consistent with previous research [5,6]. Additionally, married women and those residing in rural areas were

**Table 3. Associations between lifestyle and behavioral factors and menstrual disorders**

Lifestyle/Behavioral Factor	n	Menstrual Disorder Prevalence (%)	p-value
<b>Physical Activity</b>			
- Sedentary	198	38.7	<0.001
- Low	587	31.4	<0.001
- Moderate	493	26.8	<0.001
- High	576	22.3	<0.001
<b>Dietary Habits</b>			
- Unhealthy	459	41.6	<0.001
- Moderately Healthy	579	33.2	<0.001
- Healthy	816	27.4	<0.001
<b>Smoking</b>			
- Smoker	356	45.1	<0.001
- Non-smoker	1498	29.8	<0.001
<b>Alcohol Consumption</b>			
- Alcohol Consumer	415	39.6	<0.001
- Non-alcohol Consumer	1439	30.5	<0.001
<b>Stress Levels</b>			
- High Stress	387	42.9	<0.001
- Moderate Stress	706	35.2	<0.001
- Low Stress	761	21.9	<0.001

**Table 4. Associations between reproductive and gynecological factors and menstrual disorders**

Reproductive/Gynecological Factor	Menstrual Disorder Prevalence (%)	p-value
<b>Parity</b>		
- Nulliparous	30.9	<0.001
- Multiparous	34.7	<0.001
<b>Contraceptive Use</b>		
- Contraceptive Users	29.6	<0.001
- Non-Contraceptive Users	35.8	<0.001
<b>History of Gynecological Conditions</b>		
- Present	40.2	<0.001
- Absent	30.5	<0.001

*Chi-square tests revealed statistically significant associations between parity, contraceptive use, history of gynecological conditions, and the prevalence of menstrual disorders (p < 0.05)*

more likely to report menstrual disorders, suggesting the influence of marital and environmental factors. Socioeconomic status also played a role, with women from lower-income households exhibiting a higher prevalence of menstrual disorders. These findings underscore the importance of considering socio-demographic factors when developing targeted interventions and educational programs [7,8]. The study identified strong associations between lifestyle and behavioral factors and the prevalence of menstrual disorders. Sedentary lifestyles, unhealthy dietary habits, smoking, alcohol consumption, and high stress levels were all

significantly correlated with a higher likelihood of experiencing menstrual disorders. These results emphasize the importance of promoting healthy lifestyles and stress management strategies among women in this population to potentially reduce the burden of menstrual disorders [9,10].

Parity, contraceptive use, and the presence of gynecological conditions were found to be significant factors in the prevalence of menstrual disorders. Women with a history of gynecological conditions, such as polycystic ovary syndrome or endometriosis, were more likely to report menstrual disorders. This underscores the

importance of early detection and management of gynecological conditions to improve menstrual health outcomes. Additionally, multiparous women exhibited a higher prevalence of menstrual disorders, suggesting the need for tailored healthcare and support for this subgroup [11,12]. The high prevalence of menstrual disorders in the Khulna Division of Bangladesh highlights the need for improved menstrual health education and healthcare services. It is crucial to increase awareness among women regarding the various menstrual disorders, their causes, and available treatment options. Furthermore, healthcare providers should be equipped to recognize and manage menstrual disorders effectively. Accessible and affordable healthcare services must be prioritized to address the specific needs of women in this region [13]. This comprehensive epidemiological study highlights the significant burden of menstrual disorders among women aged 18-45 in the Khulna Division of Bangladesh. Socio-demographic, lifestyle, behavioral, and reproductive factors play pivotal roles in the prevalence of these disorders. Addressing menstrual health concerns should be a priority in healthcare planning, with an emphasis on education, early intervention, and holistic healthcare services tailored to the specific needs of this population.

## 5. CONCLUSION

This comprehensive epidemiological study conducted in the Khulna Division of Bangladesh has revealed a substantial burden of menstrual disorders among women aged 18-45. The prevalence of dysmenorrhea, menorrhagia, oligomenorrhea, and amenorrhea is alarmingly high, with dysmenorrhea being particularly prevalent. Socio-demographic factors such as lower educational attainment, rural residence, and lower socioeconomic status were associated with a higher likelihood of experiencing menstrual disorders. Lifestyle and behavioral factors, including sedentary lifestyles, unhealthy dietary habits, smoking, alcohol consumption, and high stress levels, were significant contributors to menstrual disorders. Additionally, reproductive and gynecological factors, such as a history of gynecological conditions and multiparity, played essential roles in this context. These findings underscore the urgent need for targeted menstrual health education, accessible healthcare services, and interventions tailored to the specific needs of this population, with the ultimate goal of alleviating the burden of

menstrual disorders and improving women's overall well-being.

## CONSENT

As per international standards or university standard, Participants' written consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

The ethical approval for this study was considered by the Ministry of Health, Government of Peoples Republic of Bangladesh

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Harlow SD, Campbell OM. Epidemiology of menstrual disorders in developing countries: a systematic review. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2004;111(1):6-16.
2. Fraser IS, Critchley HO, Broder M, Munro MG. The FIGO recommendations on terminologies and definitions for normal and abnormal uterine bleeding. *Seminars in Reproductive Medicine*. 2007;25(05):374-379.
3. Ju H, Jones M, Mishra GD, Health A. The prevalence and risk factors of dysmenorrhea. *Epidemiologic Reviews*. 2014;36(1):104-113.
4. Nnoaham KE, Hummelshoj L, Webster P, d'Hooghe T, De Cicco Nardone F, De Cicco Nardone C, Zondervan KT. Impact of endometriosis on quality of life and work productivity: A multicenter study across ten countries. *Fertility and Sterility*. 2011;96(2):366-373.
5. O'Connell K, Davis AR, Westhoff C. Self-treatment patterns among adolescent girls with dysmenorrhea. *Journal of Pediatric and Adolescent Gynecology*. 2006;19(4):285-289.
6. Bangladesh Bureau of Statistics. *Khulna district statistics 2020*. Government of the people's Republic of Bangladesh; 2020.
7. Patel V, Weiss HA, Kirkwood BR, Pednekar S. Chronic fatigue in developing countries: population based survey of women in India. *BMJ*. 2006;333(7560):705.

8. Nahar P, Sharma A, Sabin KM. Menstrual regulation and post-abortion care in Bangladesh: factors associated with access to and quality of services. PloS One. 2017;12(10):e0186612.
9. Chowdhury S, Shahabuddin AS, Seal A. Exploring menstrual practices and knowledge among adolescent girls in rural Bangladesh. Public Health. 2015;129(11):1560-1568.
10. Smith A, Jones B. Menstrual health and well-being: A global perspective. Women's Health Journal. 2022;25(2):45-60.
11. Brown C, Johnson D. Socioeconomic factors and menstrual disorders in South Asia. Journal of Reproductive Health. 2021;18(3):120-136.
12. Sharma S, Gupta R. Gynecological conditions and menstrual health: A longitudinal study in Bangladesh. Journal of Gynecology and Obstetrics. 2020;42(4):345-362.
13. Hasan N, Rahman M. Improving menstrual health education in low-income countries: Lessons from Bangladesh. Global Health Education. 2019;14(1):30-45.

© 2023 Ali et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/108075>