



# Effect of Storage Conditions and Duration of Storage on Seed Germination of *Tetrapleura tetraptera* (Schum and Thonn) Taub and (*Annona muricata*) Linn

C. A. Dada <sup>a\*</sup> and S. Arowosegbe <sup>b</sup>

<sup>a</sup> Department of Science Laboratory Technology (Biotechnology Option), Ekiti State University, Ado-Ekiti, Nigeria.

<sup>b</sup> Department of Plant Science and Biotechnology, Ekiti State University, Ado-Ekiti, Nigeria.

## Authors' contributions

This work was carried out in collaboration between both authors. Author CAD designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author SA manage the analyses and literature searches of the study. Both authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/aprj/2024/v12i4256>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117431>

Original Research Article

Received: 20/03/2024

Accepted: 23/05/2024

Published: 29/05/2024

## ABSTRACT

*Tetrapleura tetraptera* and *Annona muricata* are medicinal plant species with a great medicinal values. Despite their usefulness, preservation of their seed from harvest time to the time when they are required for planting has not been given adequate attention. Therefore, this study investigated the effect of storage conditions and duration of storage on seed germination of *T. tetraptera* and *Annona muricata*. Fifty (50) seeds each of *Tetrapleura tetraptera* and *Annona muricata* were planted in top soil to determine their germination percentages before subjected to storage conditions, while *Tetrapleura tetraptera* has 94% germination, *Annona muricata* has 72%

\*Corresponding author: E-mail: [comfort.dada@eksu.edu.ng](mailto:comfort.dada@eksu.edu.ng);

germination. Four storage conditions viz: cold room, refrigerator, room air conditioned and laboratory were used and the experiment was carried out in Ekiti State University, Ado – Ekiti. Seeds stored at different temperatures were collected and planted in polythene nylons filled with top-soil and watered daily, the experiments were replicated four times to test for the germinability on monthly basis for six months. In *Tetrapleura tetraptera*, highest percentage germination (50%) was recorded in seeds stored in the air conditioned room at 1 month. This was followed by seeds store in the refrigerator with 47% at 2 month. Seeds stored in the room air conditioned had percentage germination of 39% at 2 months, 30.50% at 3 months and 10.70% at 4 month of the experiment. No further germination in seeds under this treatment was observed at the last two months of the experiment. While seeds stored in refrigerator recorded 5.20% at 1 months of the experiment. Seeds stored in cold room and those in laboratory recorded no emergence throughout the period of the experiment. Also in *Annona muricata*, Highest percentage germination (58.00%) was observed in seeds stored in the room air conditioned at 1 month of the experiment, this was followed by the seeds in this same temperature at two month of the experiment with mean plumule emergence value of 39.00%, seeds under this temperature (air condition) did not respond to germination at 5<sup>th</sup> and 6<sup>th</sup> months of the experiment. Seeds of *Annona muricata* stored in the cold room responded to germination with unappreciable value of 2.05% only at third month of the experiment. Only 1.6% of the seeds stored in the laboratory emerged at 5<sup>th</sup> month of experimental time, seeds stored in the refrigerator did not respond for the first two months, last fifth and sixth month of the experiment but had 15.53% and 35.00% plumule emergence at third and fourth month of the experiment respectively. Thus, storage of seeds of *T. tetraptera* and *A. muricata* in air conditioned room with temperature of -20°C might enhance germination in these species.

**Keywords:** Medicinal plant; storage conditions, duration of storage; *Tetrapleura tetraptera*; *Annona muricata*.

## 1. INTRODUCTION

*Tetrapleura tetraptera* (Schum and Thonn.) Taub. Commonly known as Aridan in South Western Nigeria is a medicinal plant of the family Mimosaceae. The plants is locally known as Uyayak in Ibiobio; Edeminang in Efik; Osakirisa or Oshosho in Igbo and Dawo in Hausa (all in Nigeria) [1]. Its fruit is used for the management of convulsion, leprosy, inflammation, and rheumatism [2,1,3].

*Tetrapleura tetraptera* is one of the highly valued medicinal plants in Nigeria. The plant was reported to have both cardiovascular and neuromuscular activities. Pharmacologically, *T. tetraptera* is found to be anti-hypotensive, anti-ulcerative, anti-inflammatory and anti-microbial [1].

*Annona muricata*, called soursop due to the sweet and sour flavor of the large fruit, is a lowland tropical fruit-bearing tree in the Annonaceae family. Related species include cherimoya (*A. cherimola*) and sugar-apple (*A. squamosa*). Other common names include graviola and guanábana (sometimes shortened to guanába). The genus name, 'Annona' is from the Latin word 'anon', meaning 'yearly produce', referring to the production of fruits of the various species in this genus [4]. *Annona muricata* is a

small, upright, evergreen tree. Its young branches are hairy.

Based on chemotaxonomy approach, some plants of the Annonaceae family have anticancer activity [5]. *Annona montana* contains monotetrahydrofuranic acetogenins which have toxicity to liver cancer [6]. *Annona squamosa*, containing ribosome-inactivating protein (RIP), an immunotoxin for the treatment of cancer [5]. Based on this chemotaxonomy studies, plant that have close kinship for example, *Annona muricata* are likely to contain similar compounds [6].

The post-harvest process of medicinal plants have great importance in the productive chain, its direct influence on the quality and quantity of the active principles. Among the post-harvest processes, drying and storage are essential to maintain the product with the physical and chemical characteristics closer to those found in the fresh plant, and failures in any of these steps influence the quality of the final product. Drying, if not performed properly, can compromise the content of active ingredients while, incorrect storage can lead to loss of material, whether for physical or biological reasons [7,8]. Medicinal plants are often stored for long periods, before being used as raw material for the manufacture of various products [9]. However, improper

storage can result in physical, chemical and microbiological changes [10,11].

Seed viability is the ability of the embryo to germinate and it is affected by a number of factors including temperature, light, oxygen, water and species type. Germinability which is determined by germination percentage is the proportion of seeds that germinate from seeds subjected to the right conditions for growth while the germination rate is the speed with which the seeds germinate and is affected by seed viability, dormancy and environmental effects that impact on the seed and seedling (Zamora 2014). This is given as a percentage of germination over a certain amount of time, for example 90% germination within 20 days. Germination energy is the proportion of seeds in a given sample which germinate within the time of peak germination, generally taken as the highest number of germinations in a 24-hour period [12]. The interest in germination energy is based on a theory that only those seeds which germinate rapidly and vigorously under favourable conditions are likely to produce vigorous seedlings in field conditions, whereas weak seeds or delayed germination is often fatal [13].

Simple techniques have been adopted to maintain the seed viability in both domesticated and wild sources [14], (Vertucci and Roos 1991). Seed deterioration and loss of viability, is a natural phenomenon occurring during storage [15,16] and inappropriate storage medium such as room temperature storage [17,18] has often resulted in low seed germination. Several factors, including temperature, nature of the seeds, seed moisture content, relative humidity influence the seed longevity during storage [19,20]. Proper storage conditions, may effectively retain substantial viability in seeds over a considerable storage period [21,22,20].

A long-term storage of seeds, especially under unfavourable conditions, leads to loss of viability. The nature of this physiological damage is variable, e.g. short-term deterioration in the field is different from long term deterioration during storage, which in turn is different from mechanical damage [12]. All parts of the seed such as the seed coat, which is of maternal origin and acts as a physical and chemical barrier, and the embryo are susceptible to physiological damage. Seeds are grouped based on their storage characteristics as orthodox and recalcitrant. Orthodox seeds are those that can be dried to moisture contents of 10% or less, in

this condition they can successfully be stored at subfreezing temperatures [19]. Recalcitrant seeds are seeds that do not survive drying and freezing during exsitu conservation and cannot resist the effects of drying or temperatures less than 10°C; thus, they cannot be stored for long periods like the orthodox seeds because they can lose their viability. Plants that produce recalcitrant seeds include avocado, mango, mangosteen, lychee, cocoa, rubber tree, some horticultural trees, and several plants used in traditional medicine such as species of *Viola* and *Pentaclethra* [20].

Seeds of different plant species lose viability at varying degrees. For example, onion seed is very difficult to store while barley seed maintains good germination under a variety of storage conditions (Zamora 2014). Seeds and especially oilseeds including *E. bussei* contain large quantities of carbon stored as carbohydrate, lipid and protein [23]. The chemical composition of oilseeds which are rich in lipids causes specific oxidative processes to occur during storage and these are said to cause limited longevity depending on their specific chemical composition [24].

Seed of these two plant species have been reported by the local farmers in Ekiti State, Nigeria to have a negative challenge after storage. Hence, the reason for this study.

## 2. MATERIALS AND METHODS

### 2.1 Seed Source

Fresh fruits of *Tetrapleura tetraptera* were obtained from Oba's market, Ado-Ekiti, Ekiti State, Nigeria and were taken to the herbarium of the Department of Plant Science and Biotechnology, AdoEkiti, Nigeria for authentication. While the mature fruits of *Annona muricata* were collected from the parent plant from its natural habitat in Ipoti-Ekiti located in Ijero Local Government Area of Ekiti State.

### 2.2 Viability Test

Seeds were subjected to viability test by using 2,3,5-Triphenyl Tetrasolium chloride at 1% according to Kruse et al., (2004).

### 2.3 Procedure

#### 2.3.1 First step

Fifty seeds each of *Tetrapleura tetraptera* and *Annona muricata* were divided into five (5) parts and sown into black polythene bags separately filled with top soil and subjected to natural

condition favourable for germination. Germination initiation was observed for two (2) months when the total emergence was counted and recorded.

### 2.3.2 Second step

Two hundred seeds of *Tetrapleura tetraptera* and *Annona muricata* were divided into four groups consisting of 50 seeds each. Each batch were stored at four temperatures viz: cold room (22°C), refrigerator (5°C), air conditioner (-20°C) and laboratory/room temperature (2°C) as control. 20 seeds each of *Tetrapleura tetraptera* and *Annona muricata* from the four batches, stored at different temperatures were collected and planted to test for the germinability on monthly basis for six months. Planting was done in polythene nylons filled with top-soil and watered daily. The experiments were replicated four times. The number of seeds that germinate in each treatment was recorded and the percentage germination was calculated.

## 3. RESULTS

### 3.1 Percentage Germination of Seeds Before Subjected to Storage Conditions

Table 1 shows the percentage germination of both *T. tetraptera* and *A. muricata* before subjected to storage conditions. In *T.*

*tetraptera*, 94% germination was observed while 72% germination was recorded in *A. muricata*.

**Table 1. Percentage germination of seeds of *T. tetraptera* and *A. muricata***

Plants	Percentage germination (%)
<i>T. tetraptera</i>	94
<i>A. muricata</i>	72

### 3.2 Effect of Storage Conditions and Duration of Storage on Seed Germination of *T. tetraptera*

Table 2 shows that various storage condition and duration had different effect on seed germination of *T. tetraptera*. Results obtained revealed that seeds stored in cold room and those stored in the laboratory did not germinate throughout the period of experiment. Seeds stored in the refrigerator had mean germination of 5.20% and 47.00% for the first and second months experiment respectively and shows no further germination at 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> months of the experiment. Highest seed germination (50%) was recorded in seeds under air condition at the first month of the experiment, mean germination of 39%, 30.50% and 10.70% were observed at 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup> month of the experiment respectively with no further germination at 5<sup>th</sup> and 6<sup>th</sup> month in this same (air condition) treatment.

**Table 2. Percentage Germination of *T. tetraptera* seeds stored under different storage conditions**

Months	Percentage Germination (%)			
	Cold room	Refrigerator	Air conditioned	Room Laboratory
1 <sup>st</sup>	0	5.20	50.00	0
2 <sup>nd</sup>	0	47.00	39.00	0
3 <sup>rd</sup>	0	0	30.50	0
4 <sup>th</sup>	0	0	10.70	0
5 <sup>th</sup>	0	0	0	0
6 <sup>th</sup>	0	0	0	0

**Table 3. Percentage Germination of *A. muricata* seeds stored under different storage conditions**

Months	Percentage Plumule Emergence (%)			
	Cold room	Refrigerator	Air conditioned	Room Laboratory
1 <sup>st</sup>	0	0	58.00	0
2 <sup>nd</sup>	0	0	39.00	0
3 <sup>rd</sup>	2.05	15.53	10.50	0
4 <sup>th</sup>	0	35.00	9.82	0
5 <sup>th</sup>	0	0	0	1.6
6 <sup>th</sup>	0	0	0	0

### 3.3 Effects of Storage Conditions and Duration of Storage on Seed Germination of *A. muricata*

Table 3 shows that seeds of *Annona muricata* stored in the cold room responded to germination with unappreciable value of 2.05% only at third month of the experiment, while no emergence was observed at first two months and the last three months of the experiment in this treatment. Only 1.6% of the seeds stored in the laboratory emerged at 5<sup>th</sup> month of experimental time, seeds stored in the refrigerator did not respond for the first two months, last fifth and sixth month of the experiment but had 15.53% and 35.00% plumule emergence at third and fourth month of the experiment respectively.

Highest percentage germination (58.00%) was observed in seeds stored in the room air conditioned at one month of the experiment, this was followed by the seeds in this same temperature at two month of the experiment with mean plumule emergence value of 39.00%, a decrease in the percentage plumule emergence in seeds under this same temperature was observed at 3<sup>rd</sup> and 4<sup>th</sup> months of the experiment with 10.50% and 9.82% respectively. While the seeds under this temperature (air condition) did not respond to germination at 5<sup>th</sup> and 6<sup>th</sup> months of the experiment.

### 4. DISCUSSION

Results from this study emphasized the fact that appropriate storage conditions and duration of storage are of great necessity in preserving the viability of seeds from the time of collection to the time when they are needed for sowing. In fact, Genes and Nasreen [25] observed that the use of seed is one of the propagation methods that can be used to multiply plants but the seeds need to be capable of germinating in the first place., hence the need for knowledge of its methods of germination. Since most plants produce seeds only once a year, knowledge of methods of seeds preservation for next season's generation was deemed important and so was the determination of how soon the seeds would lose its viability.

The results obtained in this study revealed that seeds of *T. tetraptera* and *A. muricata* stored in air conditioned room had the best percentage germination when compared with seeds in other storage conditions. Reason for this could be that temperature in other storage conditions might be

too low or high to promote seed germination in this plant or that they were too high to enhance proper germination of this species. This finding agreed with previous observation of Lars [26] that seeds germinate faster with increasing air temperatures up to a point. Extreme heat will slow germination and also increase moisture loss. The temperatures for optimal germination vary with the type of plant, some annual flowers and vegetables are extremely sensitive to cold, extremely hot or cold temperatures can hamper plant growth, as well as affect seed germination Lars [26]. Gamaene et al., [27] ascertained that under no storage condition would recalcitrant seed be kept for long as this tends to reduce its viability [28-30].

### 5. CONCLUSION

Storage conditions with extremely hot or cold temperatures does not encourage seed germination of these plant species. Therefore, storage of seeds of *T. tetraptera* and *A. muricata* in air conditioned room with temperature of -20°C to keep its viability intact is recommended

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Ojewole JA, Adewunmi CO. Anti-inflammatory and hypoglycemic effects of *Tetrapleura tetraptera* (Taub) (Fabaceae) fruit aqueous extract in rats. J. Ethnopharmacol. 2004;95(23):177-182.
2. Okwu DE. The potentials of *Ocimum gratissimum*, *Pergularia extensa* and *Tetrapleura tetraptera* fruit as spice and flavouring agents. Nig. Agric J. 2003;35: 143-148.
3. Jamwal KS, Sharma ML, Chandhoke NR, Chatak BJ. Pharmacological action of 6,7-Dimethoxycoumarin (Scoparone) isolated from *Artemisia scoparia* Waldst & Kit. Indian J. Med. Res. 1972;60(5): 763.
4. Natural resources Conservation Service. Plants profile *Annona squamosa* L. United States Department of Agriculture. 2008;4-17.
5. Sismindari Hussana A, Mubarika S. Squensing of DNA Supercoil Double Helix with *In vitro* method from *Annona squamosa* L. 2008;32:142- 147

6. Roesker R, Catharino RR, Malta LG. Antioxidant activity of *Annona crassiflora*: Characterization of major components by electrospray ionization mass spectrometry. Food Chemistry. 2007;104: 1048-1054.
7. Martinazzo AP. Secagem, armazenamento e qualidade de folhas de *Cymbopogon citratus* (D.C.) Stapf. 156 f. Tese (Doutorado em Engenharia Agrícola) – Departamento de Engenharia Agrícola, Universidade Federal de Vicosa, Minas Gerais; 2006.
8. Martinazzo AP, Correa PC, Melo EC, Barbosa FF. Difusividade efetiva em folhas de *Cymbopogon citratus* (DC.) Stapf submetidas a secagem com diferentes comprimentos de corte e temperaturas do ar. Revista Brasileira de Plantas Mediciniais. 2007;9(1):68-72.
9. Masand S, Madan S, Balian SK. Modern concept of storage and packaging of raw herbs used in Ayurveda. International Journal of Research in Ayurveda and Pharmacy. 2014;5(2):242-245.
10. Peter KV. Handbook of Herbs and Spices. 3. New York, USA; 2006.
11. Mayuoni-Kirshinbaum L, Daus A, Porat R. Changes in sensory quality and aroma volatile composition during prolonged storage of 'Wonderful' pomegranate fruit. International Journal of Food Science and Technology. 2013;48:1569-1578.
12. Mrđa J, Crnobarac J, Dušanić N, Jocić S, Miklić V. Germination energy as a parameter of seed quality in different sunflower genotypes – Genetika. 2011; 43(3):427-436.
13. Aldhous JR. Nursery practice. Forestry Comm. Bull. No. 43, London; 1972.
14. Ellis RH, Hong TD, Roberts EH. The low-moisture-content limit to the negative logarithmic relation between seed longevity and moisture content in three sub species of rice. Annal. Bot. 1992;69(1):53–58.
15. Nasreen S, Khan BR, Mohmad AS. The effect of storage temperature, storage period and moisture content on seed viability of soyabean. Pakistan J. of Biol. Sci. 12/2000;3:12.
16. Schmidt L. Guide to handling of Tropical and Subtropical forest seeds Denmark: DANIDA Forest Seed Centre, Borch Tyrk A/S; 2002.
17. Hezewijk MJ, Van BAP, Verleij JAC, Pieterse AH. Germination of *Orobanche crenata* seeds as influenced by conditioning temperature and period. Can. J. of Bot. 1993;71(6):786-792.
18. Muller E, Cooper EJ, Alsos IG. Germinability of arctic plants is high in perceived optimal conditions but low in the field. Botany. 2011;89(5):337-348.
19. Onyekwelua JC, Fayose OJ. Effect of storage methods on the germination and proximate composition of *Treulia africana* seeds. In: Proceedings of the International Conference on Agricultural Research for Development; University of KasselWitzenhausen and University of Göttingen, Tropentag; 2007.
20. Pradhan BK, Badola HK. Seed germination response of populations of *Swertia chirayita* following periodical storage. Seed Technol. 2008;30(1):63–69.
21. Butalo JS, Badola HK. Effect of pre-sowing treatment on seed germination and seedling vigour in *Angelica glauca*, a threatened medicinal herb. Cur. Sci. 2004;87(6):796-799.
22. Chen SY, Kuo SR, Chien CT. Storage behaviour of seeds of *Cinnamomum osmophloem* and *Neolisea aciculate var. var. billianna* (*Jauracaceae*). Seed Sci. Technol. 2007;35(1):237-243.
23. Bewley JD, Black M. Seed physiology of development and germination. Plenum Press, New York; 1985.
24. Balešević S, Tatic M, Miladinovic J, Pucarevic M. Changes of fatty acids content and vigor of sunflower seed during natural aging. Helia. 2007;30(47): 6168.
25. Nasreen S. Effects of storage period and temperature on seed viability of Wheat. Park J. Bio Sci. 1995;2:1492 – 1493.
26. Lars S. Guide to handling of tropical and subtropical forest seed. Centre, Humlebaek, Denmark; 2000.
27. Gamaene SHW, Pritchard C, Harris A. *Vitellaria paradoxa*. Handling and storage of recalcitrant and intermediate tropical forest tree seeds, Newsletter. Danida Forest Seed Centre, Humlebaek, Denmark. 1999;5:15-16.
28. Genes F, Nyomora AMS. Effect of storage time and temperature on germination ability of *Escoecaria bussei*. Tanzania Journal of Science. 2018;44(1):123-133.

29. Vertucci CW, Roos EE. Theoretical basis of protocols for seed storage. *Plant Physiol.* 1990; 94(3):1019–1023.
30. Zamora FJH. Effect of moisture on lentil seed germination and growth; 2013. Available:<http://en.wikipedia.org/wiki/Germination>.

---

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/117431>