



# Contribution to Phytochemical, Antioxidant and Antibacterial Studies of Extracts of Crushed, Sieved and Rejected organs of Two Medicinal Plants from Côte d'Ivoire

N'guessan Patrick Audrey <sup>a\*</sup>, N'guessan Djagna Michael <sup>a</sup>,  
Konan Dibi Jacques <sup>a</sup>,  
Mamyrbékova-Békro Janat Akhanovna <sup>a</sup>  
and Békro Yves-Alain <sup>a</sup>

<sup>a</sup> *Laboratoire de Chimie Bio-Organique et de Substances Naturelles, UFR-SFA, Université Nangui Abrogoua, Abidjan, Côte d'Ivoire.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/ijbcrr/2024/v33i6901>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/122838>

**Original Research Article**

**Received: 27/06/2024**

**Accepted: 02/09/2024**

**Published: 05/09/2024**

## **ABSTRACT**

**Aims:** This research work was carried out on the Broyat, Tamisat and Refus of the whole plant *Heterotis rotundifolia* and the leaves of *Manihot esculenta*, two medicinal plants from Côte d'Ivoire. This study aims to evaluate the influence of the size of particles from plant matrices on the extraction of organic phytochemicals and their biological activity.

\*Corresponding author: E-mail: [patrickaudrey2014@gmail.com](mailto:patrickaudrey2014@gmail.com);

**Cite as:** Audrey, N'guessan Patrick, N'guessan Djagna Michael, Konan Dibi Jacques, Mamyrbékova-Békro Janat Akhanovna, and Békro Yves-Alain. 2024. "Contribution to Phytochemical, Antioxidant and Antibacterial Studies of Extracts of Crushed, Sieved and Rejected Organs of Two Medicinal Plants from Côte d'Ivoire". *International Journal of Biochemistry Research & Review* 33 (6):185-95. <https://doi.org/10.9734/ijbcrr/2024/v33i6901>.

**Methodology:** For this purpose, the crude extracts obtained by extraction (Soxhlet/maceration) and selective by liquid-liquid contact were prepared. Concerning the phytochemical screening, the selective extracts were used to carry out the TLC while the hydroethanolic crude extracts were used for identification tests by colored reactions (in tube), the dosage of total polyphenols, total flavonoids and condensed tannins as well as for their antioxidant potential and their antibacterial action. The antioxidant activity was evaluated by spectrophotometry using the DPPH radical while the antibacterial activity was estimated against the strains *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa*.

**Results:** Phytochemical screening results show that plant extracts contain polyphenols, flavonoids, tannins, terpenes, sterols and sugars. The Broyat yields a greater number of extractable phytocompounds than the Tamisat and the Refus, regardless of the type of extraction solvent and plant species used. The Tamisat contains the highest number of compounds of interest, and therefore displays greater antioxidant potential against DPPH and antibacterial activity against *Staphylococcus aureus*.

**Conclusion:** At the end of this study, it emerged that it would be preferable to use the Broyat, for phytochemical studies, and the Tamisat for biological studies.

**Keywords:** *Heterotis rotundifolia*; *Manihot esculenta*; antioxidant; antibacterial; studies.

## 1. INTRODUCTION

Plant extracts have long been used in various medicinal and therapeutic applications. The growing emphasis on natural compounds and their health-promoting properties has led to a significant increase in studies aimed at better understanding their chemical composition and biological activities [1]. A crucial step in the analysis of organic compounds present in plants is extraction. Different extraction methods and techniques have been developed to obtain extracts rich in bioactive compounds. Extraction is influenced by the type of solvent, temperature, liquid-solid ratio, pH, pressure, number of extraction steps and particle size [2,3]. Indeed, particle size refers to the size of the particles in a material, and in this specific case, it is the size of the particles of crushed and sieved plant organs used in the extraction. When plant organs are ground, they can be reduced into different particle sizes which can then be sieved. In order to optimize the yield of the extraction of secondary metabolites from plant organs, some experimenters prefer to use Broyat to carry out the extraction while others opt instead for Tamisat. Therefore, we were interested in knowing which of the plant meshes (Broyat, Tamisat, Refus) would be likely to give the best results both in terms of the extraction of phytocompounds and the biological profile. It is in this context that this study was initiated. The present work aims to evaluate the influence of the size of particles from plant matrices on the extraction of organic phytocompounds and their biological activity. To achieve this, it was specifically necessary to extract the

phytoconstituents from Broyat, Tamisat and Refus from the study plants; to carry out qualitative and quantitative phytochemical screening then to evaluate the antioxidant and biological potential of the plant mesh.

## 2. MATERIALS AND METHODS

### 2.1 Materials

#### 2.1.1 Plant material

The plant material consists of the leaves of *Manihot esculenta* and the whole plant *Heterotis rotundifolia*. The harvest was made on the site of Nangui ABROGOUA University (UNA) (Abidjan / Ivory Coast). The plant material collected was identified by referring to the species in the herbarium of the National Floristic Center (CNF) of the Félix HOUPHOUËT-BOIGNY University (Cocody / Abidjan), then authenticated by Professor MALAN Djah François (Systematic botanist at the UNA). The plant material was cleaned and dried for 2 days away from the sun in a ventilated room then in a room with air conditioning (18°C) for 7 days and in an oven (45°C) for 3 days.

#### 2.1.2 Laboratory equipment

The laboratory equipment consists of usual laboratory glassware, an oven, a UV-254/366 nm lamp (VL-6.LC), an electronic balance (XY6002C), a rotary evaporator brand (BÜCHI type EL-131), a refrigerator, a dryer, a UV-visible spectrophotometer (Spectro AL 800), a hood, an

electric grinder (Nutri Blitzer GS 158- V1457). To carry out Thin Layer Chromatography (TLC), chromatoplates (silica gel 60 F254, aluminum support, 20 × 20) (Merck) were used.

### 2.1.3 Chemicals

The chemicals used were purchased from Ryca Pharma, Polychemistry and Chimtec. The organic solvents used have a purity of 96%.

### 2.1.4 Biological material

The bacterial strains (*Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*) were provided by the biobank of the Pasteur Institute of Côte d'Ivoire (IPCI). They were selected because of their recurrent involvement in a good number of human pathologies.

## 2.2 Methods

### 2.2.1 Acquisition of powders of different particle sizes

After drying, the plant material was pulverized using an electric grinder (nutri blitzer GS 158-V1457) to provide powders called "Broyat". Subsequently, the Broyat was passed through a sieve with a diameter of less than 1 mm to obtain on the one hand, "Tamisat", and on the other hand "Refus" which constitutes the powder having been retained in the sieve. The vegetable meshes (Broyat, Tamisat, Refus) were used to prepare the different extracts.

### 2.2.2 Soxhlet extraction

A Whatman cartridge containing 5 g of vegetable powder is introduced into the extraction chamber, connected at the top to a condenser, and at the bottom to a ground-neck flask containing 300 mL of solvent. Petroleum ether, ethyl acetate and ethanol were used as extraction solvents, respectively. This operation was repeated 3 times. After vacuum filtration, the crude ethereal, ethyl acetate and ethanolic extracts are concentrated using a rotary evaporator at 40°C, then stored in vials protected from light. The extracts obtained by Soxhlet were used to calculate only the yields. This preliminary study was carried out in order to observe the behavior of solvents of different polarity with respect to the powders of the Broyat, the Tamisat and the Refus from the study plants.

### 2.2.3 Preparation of hydroethanolic and selective extracts

Five (5) g of powder from the Broyat, the Tamisat and the Refus of each plant were macerated in 60 mL of an 80% hydroethanolic solution for 24 hours. The operation was repeated 3 times with the same marcs. After filtration on Büchner, the macerates obtained were combined, then concentrated at 40 ° C using a rotary evaporator. The crude hydroethanolic extracts obtained were used on the one hand, for the preparation of selective extracts and on the other hand, for phytochemical screening by color reactions, quantification of phenolic compounds (polyphenols, flavonoids and total tannins), and for the evaluation of antioxidant and antibacterial potential.

A volume of 15 mL of the crude hydroethanolic extract was exhausted by successive fractionations with (3 × 10 mL) hexane, (3 × 10 mL) dichloromethane and (3 × 10 mL) ethyl acetate. The different selective organic filtrates are concentrated in the oven then stored in the refrigerator at 4°C. They were used for phytochemical screening by TLC.

### 2.2.4 Qualitative tests in tube and on TLC plate

To carry out colored tube tests on crude hydroethanolic extracts, the analytical techniques used are those described in the literature [4-6].

The TLC screening of the selective extracts was carried out following the methods described in the literature [7].

### 2.2.5 Quantitative analysis

#### 2.2.5.1 Dosage of total phytophenols

Total polyphenol contents were determined using the Folin-Ciocalteu colorimetric method [8]. To 1 mL of each extract are added 1.5 mL of sodium carbonate (Na<sub>2</sub>CO<sub>3</sub>) (17%, m/v) and 0.5 mL of Folin-Ciocalteu reagent (0.5N). The whole is incubated at 37°C for 30 min and the absorbance is read at 760 nm on the spectrophotometer against a blank without plant extract taken as a reference. The quantification of total phytophenols is carried out according to a linear calibration line ( $y = ax + b$ ) carried out by a standard extract of gallic acid at different concentrations (0 to 1000 µg/mL) under the same conditions that the sample.

### 2.2.5.2 Dosage of total flavonoids

This assay was carried out according to the method used by Arvouet-Grand and colleagues [9]. 500  $\mu$ L of 2% aluminum chloride ( $AlCl_3$ ) in methanol was added to an equal volume of plant extract. After 10 min of incubation, the reading was taken with a spectrophotometer at 415 nm. Quercetin (0-100  $\mu$ g/L), used as a standard, allowed the establishment of the calibration curve. Three (3) tests were carried out for each plant extract, and the result given is an average of the 3 readings.

### 2.2.5.3 Dosage of condensed tannins

The determination of condensed tannins in the different plant extracts was carried out according to the method described by Heimler and colleagues [10]. For 400  $\mu$ L of each plant or standard sample, 3 mL of a 4% methanolic vanillin solution and 1.5 mL of concentrated HCl are added. The mixture is incubated for 15 min and the absorbance is read at 500 nm on the spectrophotometer. Condensed tannin contents are deduced from the calibration curve established with catechin.

## 2.2.6 Evaluation of antioxidant potential by the DPPH test

The method used is that of Brand-Williams and colleagues [11] with slight modifications in terms of concentrations. DPPH is solubilized in absolute ethanol to obtain a solution with a concentration of 0.03 mg/mL. Different concentration ranges of each plant extract are prepared with the same solvent (1 mg/mL, 0.5 mg/mL, 0.25 mg/mL; 0.125 mg/mL; 0.0625 mg/mL; 0.03125 mg /mL; 0.015625 mg/mL; 0.007812 mg/mL; ; 0.003906 mg/mL and 0.001953 mg/mL). In dry and sterile tubes, 1 mL of plant extract solution to be analyzed and 2 mL of DPPH solution are introduced. After shaking, the tubes are placed away from light for 30 min. The absorbance of the mixture is measured at 517 nm with a spectrophotometer against a blank (2 mL of DPPH solution + 1 mL of absolute ethanol). The positive reference control used is ascorbic acid (vitamin C) prepared under the same conditions as the samples.

## 2.2.7 Biological study

### 2.2.7.1 Preparation of the inoculum for the efficacy test

Preparation of the bacterial inoculum began with the collection of a well-isolated bacterial colony

18 to 24 hours later. It was homogenized in 2 mL of sterile physiological water in order to obtain an optical density of 0.5 Mc Farland, with a bacterial population of approximately 10<sup>6</sup> CFU/mL. This inoculum was used to inoculate the surface of the Mueller Hinton agar plates in tight streaks using a swab to carry out the tests [12].

### 2.2.7.2 Testing the effectiveness of extracts

On the agar previously inoculated, the extracts to be tested with a concentration of 100 mg/mL were placed in the previously prepared wells. The antibiotics (Cefotaxime (COX), Ceftazidime (CZD) and Cefoxitin (FOX)) involved in the treatment of pathologies linked to the bacterial strains to be tested served as positive controls. They were also placed on the agar. Sterile distilled water constituted the negative control [13,14].

### 2.2.7.3 Determination of the Minimum Inhibitory Concentration (CMI)

One hundred microliter (100  $\mu$ L) of bacterial inoculum is distributed into the microplate wells. To these quantities are added 100  $\mu$ L of the different concentrations ( $C_1$  to  $C_6$ ) of the extracts to be tested, from the highest concentration to the lowest with the exception of the wells of the growth control ( $T_c$ ) in which only 100  $\mu$ L of sterile distilled water (the total volume of each well being 200  $\mu$ L). Concerning the wells reserved for sterility witnesses ( $T_s$ ), only 200  $\mu$ L of distilled water are added. The microplate is subsequently incubated at 37°C for 24h. The minimum inhibitory concentration (CMI) corresponds to the concentration of the first experimental well from which no disorder is observed with the naked eye. This operation was repeated 3 times in a row [15].

### 2.2.7.4 Determination of the minimum bactericidal concentration (CMB)

First, 4 successive dilutions at 1/10 from 10<sup>-1</sup> to 10<sup>-4</sup> are made from the starting inoculum. These dilutions and the inoculum are inoculated in 5 cm streaks on the different MH agar plates, then incubated at 37°C for 24 h. These Petri dishes constituted the A dishes [14]. Then, the wells of the microplate in which no disorder is observed with the naked eye are seeded in 5 cm streaks on the MH agar plates, starting with the CMI tube. They are then incubated at 37°C for 24 hours. These boxes constituted boxes B. The CMB corresponds to the lowest concentration of

extract for which there is at most 0.01% of surviving bacteria. It is read by comparing box B to box A [15].

### 3. RESULTS AND DISCUSSION

#### 3.1 Extraction Yields

The phytochemicals present in the different powders studied were extracted using a Soxhlet with 3 types of solvents (petroleum ether, ethyl acetate and ethanol). The yields of these extractions are recorded in Tables 1, 2 and 3.

The Soxhlet extractions carried out in this work made it possible to obtain variable extract yields depending on the botanical nature of the plant species, the type of solvent and the type of powder (Broyat, Tamisat or Refus) used for the different extractions. Considering the results of the yields obtained, ethanol is the solvent which best extracts the phytochemicals unlike ethyl acetate and petroleum ether. The extractable power of a solvent being linked to its polarity, these results are therefore due to the fact that ethanol is more polar than ethyl acetate and itself more polar than petroleum ether. We note that whatever the plant species and the type of solvent used, the extraction yields of Broyat are higher than those of Tamisat and Refus. Indeed, grinding allows better accessibility to compounds by increasing their bioavailability while sieving induces a differential distribution of active compounds depending on particle size [16].

#### 3.2 Qualitative Phytochemical Profiles

The results of the detection tests by color tube reactions and by TLC of the secondary metabolites of the extracts of *M. esculenta* and *H. rotundifolia* powders are recorded in Table 4.

Taking into account the results obtained at the end of the qualitative study, we note that the extracts of *M. esculenta* and *H. rotundifolia* contain polyphenols, flavonoids, coumarins, tannins, sterols, terpenes and proteins. Phenolic acids were observed only in *H. rotundifolia*. Furthermore, the qualitative analysis carried out by Etekpo [17] on the plant species *H. rotundifolia* indicated the presence of sterols, steroids, terpenes, triterpenes, 1,8-diantracene, coumarins, anthocyanins, flavonoids, tannins, phenolic acids, saponins and alkaloids.

In addition, the Broyat, Tamisat and Refus of the study extracts are made up of the same phytoconstituents. This qualitative study therefore does not make it possible to predict which Broyat, Tamisat or Refus contains the most phytochemicals.

#### 3.3 Contents of Phytophenols, Total Flavonoids and Condensed Tannins

Tables 5-7 present varied contents of phytophenols, total flavonoids and condensed tannins of the hydroethanolic extracts obtained from Broyat, Tamisat and Refus from the leaves of *Manihot esculenta* and the whole plant *Heterotis rotundifolia*.

**Table 1. Percentages obtained from petroleum ether extracts**

Plant Species	Yield (%)		
	Broyat	Tamisat	Refus
<i>M. esculenta</i>	17.40 ± 0.00	16.00 ± 0.00	17.20 ± 0.09
<i>H. rotundifolia</i>	15.73 ± 0.31	7.87 ± 0.09	11.80 ± 0.13

**Table 2. Percentages obtained from ethyl acetate extracts**

Plant Species	Yield (%)		
	Broyat	Tamisat	Refus
<i>M. esculenta</i>	13.80 ± 0.00	13.40 ± 0.27	11.80 ± 0.00
<i>H. rotundifolia</i>	16.07 ± 0.22	10.60 ± 0.27	10.40 ± 0.27

**Table 3. Percentages obtained from ethanol extracts**

Plant Species	Yield (%)		
	Broyat	Tamisat	Refus
<i>M. esculenta</i>	20.47 ± 0.36	14.67 ± 0.22	18.53 ± 0.22
<i>H. rotundifolia</i>	17.60 ± 0.27	10.60 ± 0.40	13.53 ± 0.09

**Table 4. Summary of qualitative analyzes in tube and by TLC**

	PP	Flav	Coum	Tan	Ste/Ter	Sap	Suc	Cp. Red	Ac. Ph	Pro
BMe	+	+	+	+	+	-	+	-	-	+
TMe	+	+	+	+	+	-	+	-	-	+
RMe	+	+	+	+	+	-	+	-	-	+
BHr	+	+	+	+	+	-	+	-	+	+
THr	+	+	+	+	+	-	+	-	+	+
RHr	+	+	+	+	+	-	+	-	+	+

PP : Polyphenols; Flav: Flavonoids; Coum: Coumarins; Tan: Tannins; Ste/Ter: Sterols/Terpenes; Sap: Saponins; Suc: Sugars; Cp. Red : Reducing compounds ; Ac. Ph : Phenolic acids; Pro: Protein; BMe: Broyat of *M. esculenta*; TMe: Tamisat of *M. esculenta*; RMe: Refus of *M. esculenta*; BHr: Broyat of *H. rotundifolia*; THr: Tamisat of *H. rotundifolia*; RHr: Refus of *H. rotundifolia*

**Table 5. Total phytophenol contents of hydroethanolic extracts**

Extracts	Total phytophenol contents (mg EAG/g DM)
BMe	146.24 ± 3.67
TMe	84.10 ± 2.23
RMe	92.85 ± 0.12
BHr	53.29 ± 0.60
THr	43.98 ± 0.44
RHr	57.13 ± 1.31

**Table 6. Total flavonoid contents of hydroethanolic extracts**

Extracts	Total flavonoid contents (mg EQ/g DM)
BMe	20.77 ± 1.74
TMe	23.49 ± 1.15
RMe	7.68 ± 0.62
BHr	3.76 ± 0.47
THr	1.42 ± 0.04
RHr	4.45 ± 0.16

The quantification of phenolic compounds in the leaves of *M. esculenta* revealed that the Broyat presents a better content of total polyphenols with a value of 146.24±3.67 mg EAG/g DM followed by the Refus with 92.85±0.12 mg EAG/g DM while Tamisat had the lowest content 84.10±2.23 mg EAG/g DM. Furthermore, The Refus signed the highest content of total polyphenols (57.13±1.31 mg EAG/g DM) in the whole plant of *H. rotundifolia* followed by Broyat (53.29±0.60 mg EAG/ g DM), as for Tamisat, it presented the lowest content (43.98±0.44 mg EAG/g DM); It appears that following the quantitative analysis of phenolic compounds, Tamisat presented the lowest content of total polyphenols regardless of the plant species used. Regarding the quantification of total flavonoids and condensed tannins in the leaves of *M. esculenta*, The Tamisat manifested the highest contents respectively (23.49±1.15 mg

EQ/g DM and 66.78±3.16 mg EC/g DM) followed by Broyat with respective values of 20.77±1.74 mg EQ/g DM and 51.09±0.98 mg EC/g DM. Note that the lowest contents were observed with Refus with respective values of 7.68±0.62 mg EQ/g DM and 20.82±1.34 mg EC/g DM. On the other hand, regarding the determination of the quantities of total flavonoids and condensed tannins in the whole plant *H. rotundifolia*, the Refus presented the highest contents (4.45±0.16 mg EQ/g DM and 14.26±0.61 mg EC/g DM). The Broyat for their part, showed total flavonoid contents of the order of 3.76±0.47 mg EQ/g DM and those of condensed tannins worth 9.59±0.39 mg EC/g DM. The Tamisat presented the lowest contents of respective values 1.42±0.04 mg EQ/g DM and 3.18±0.24 mg EC/g DM.

It appears from this quantitative analysis of total flavonoids and condensed tannins that the Tamisat of *M. esculenta* had the highest contents compared to the Broyat and Refus coming from the same plant matrix. On the other hand, in the case of the plant species *H. rotundifolia*, it was rather the Refus which showed the highest contents of total flavonoids and condensed tannins.

**Table 7. Condensed tannin contents of hydroethanolic extracts**

Extracts	Condensed tannin contents (mg EC/g DM)
BMe	51.09 ± 0.98
TMe	66.78 ± 3.16
RMe	20.82 ± 1.34
BHr	9.59 ± 0.39
THr	3.18 ± 0.24
RHr	14.26 ± 0.61

BMe: Broyat of *M. esculenta*; TMe: Tamisat of *M. esculenta*; RMe: Refus of *M. esculenta*; BHr: Broyat of *H. rotundifolia*; THr: Tamisat of *H. rotundifolia*; RHr: Refus of *H. rotundifolia*

### 3.4 Antioxidant profiles with respect to DPPH

#### 3.4.1 Percentage Reduction (PR)

Figs. 1 and 2 present the PR values obtained from *M. esculenta* leaf powders with concentrations ranging from 0.031 mg/mL to 0.500 mg/mL and those from the whole *H. rotundifolia* plant for concentrations between 0.016 mg/mL and 0.130 mg/mL.

The hydroethanolic extracts of the powders of the two plant species show an antioxidant potential whatever the concentration of the

extract. The synergistic action of the reducing secondary metabolites present in the study extracts could be the origin of their antioxidant activity when analyzing the results of the quantitative tests. Indeed, certain secondary metabolites such as flavonoids, terpenes, etc. are known for their ability to reduce free radicals [18,19].

Furthermore, the highest PR values are observed in Tamisat for the two plant matrices. For a better assessment of the effectiveness of said extracts, their median concentration (CR<sub>50</sub>) [20] was determined graphically using the EXCELL software.




Fig. 1. Percentage reduction in hydroethanolic extracts from *M. esculenta* leaves




Fig. 2. Percentage reduction in hydroethanolic extracts from the whole *H. rotundifolia* plant

### 3.4.2 Median Reduction concentration (CR<sub>50</sub>)

DPPH radical using ascorbic acid (vitamin C) as a reference antioxidant. The CR<sub>50</sub> of the plant extracts studied and vitamin C were determined graphically using EXCELL software. Each CR<sub>50</sub> is calculated by solving the equation  $y= ax+b$  of each trend curve with:  $y=50$ . The average CR<sub>50</sub> values obtained are shown in Table 8.

The antioxidant activities of the different plant extracts were evaluated based on their CR<sub>50</sub> in comparison with that of vitamin C. Thus, the lower the value of the CR<sub>50</sub>, the greater the antioxidant power of the extract [21]. The evaluation of the antioxidant activity by spectrophotometry which is intended to be quantitative showed that vitamin C presents the best antioxidant power with a CR<sub>50</sub> equal to 0.002 mg/mL. However, the antioxidant activity of the different plant extracts is not negligible compared to that of the reference molecule (vitamin C). This is because vitamin C is a pure molecule while plant extracts are complex in terms of chemical composition. On the other hand, the Tamisat extracts showed greater antioxidant activity with respect to DPPH than the Broyat and Tamisat. In addition, Broyat have a greater activity than Refus. In the case of *M. esculenta* leaf extracts, the sieved particles are richer in total flavonoids and condensed tannins. This would justify the fact that Tamisat have

greater antioxidant activity than those of Broyat and Refus. Indeed, flavonoids and tannins, more precisely proanthocyanidins, are powerful antioxidants [22]. Furthermore, with regard to the whole plant *H. rotundifolia*, the highest antioxidant activity observed in Tamisat could be due to its phytochemical composition, in particular to the presence of phenolic acids detected only in said plant species.

### 3.5 Antibacterial Profile

#### 3.5.1 Antibacterial activity in solid media

The antibacterial activity in a solid medium made it possible to research the sensitivity of bacteria to hydroethanolic extracts of powders (Broyat, Tamisat, Refus) from the leaves of *M. esculenta* and the whole plant of *H. rotundifolia*. It also made it possible to check their behavior in the face of the different antibiotics used. The choice of antibiotics was made according to their specificity in relation to the different groups of bacteria used for the tests. Thus, cefotaxime (COX) was used for *Escherichia Coli*, cefoxitin (FOX) was used for *Staphylococcus aureus* and finally ceftazidime (CZD) was tested on *Pseudomonas aeruginosa*. These antibiotics played the role of positive controls and sterile distilled water (EDS) that of negative control. The results obtained are recorded in Table 9.

**Table 8. CR<sub>50</sub> values of *M. esculenta* leaf powders, whole *H. rotundifolia* plant and vitamin C**

	BMe	TMe	RMe	BHr	THr	RHr	Vit. C
CR <sub>50</sub> (mg/mL)	0.229	0.215	0.403	0.059	0.02	0.088	0.002

BMe: Broyat of *M. esculenta*; TMe: Tamisat of *M. esculenta*; RMe: Refus of *M. esculenta*; BHr: Broyat of *H. rotundifolia*; THr: Tamisat of *H. rotundifolia*; RHr : Refus of *H. rotundifolia*; Vit.C : Vitamine C

**Table 9. Diameters of inhibition zones of bacterial strains**

	Diameters of inhibition zones (mm)									
	BMe	TMe	RMe	BHr	THr	RHr	COX	FOX	CZD	EDS
<i>E. Coli</i> ATCC	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	18.00 ± 0.00	-	-	06.00 ± 0.00
<i>S. aureus</i> ATCC	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	11.00 ± 0.00	13.00 ± 0.00	11.00 ± 0.00	-	22.00 ± 0.00	-	06.00 ± 0.00
<i>S. aureus</i> 068	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	10.00 ± 0.00	11.00 ± 0.00	10.00 ± 0.00	-	18.00 ± 0.00	-	06.00 ± 0.00
<i>P. aeruginosa</i> ATCC	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	-	-	18.00 ± 0.00	06.00 ± 0.00
<i>P. aeruginosa</i> 363	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	06.00 ± 0.00	-	-	06.00 ± 0.00	06.00 ± 0.00

BMe: Broyat of *M. esculenta*; TMe: Tamisat of *M. esculenta*; RMe : Refus of *M. esculenta*; BHr: Broyat of *H. rotundifolia*; THr: Tamisat of *H. rotundifolia*; RHr : Refus of *H. rotundifolia*; COX : cefotaxime; FOX : cefoxitin; CZD : ceftazidime; EDS : sterile distilled water



**Table 10. Antibacterial parameters of plant extracts from *H. rotundifolia* against *Staphylococcus aureus***

Strains	Extracts	CMI (mg/mL)	CMB (mg/mL)	CMB/CMI	Interpretation
<i>Staphylococcus aureus</i> ATCC	BHr	50	>100	>2	Bacteriostatic
	THr	6.25	12.5	2	Bactericidal
	RHr	50	>100	>2	Bacteriostatic
<i>Staphylococcus aureus</i> 068	BHr	50	>100	>2	Bacteriostatic
	THr	6.25	12.5	2	Bactericidal
	RHr	50	>100	>2	Bacteriostatic

The diameters of the zones of inhibition of bacterial strains by plant extracts vary from one strain to another depending on the extract used. Some powders have shown antimicrobial activity, however, others have not. In fact, the strain is said to be resistant to the substance when the diameter of its inhibition zone is less than 8 mm, sensitive when it is between 9 and 14 mm, very sensitive when it is between 15 and 19 mm, and extremely sensitive when it is greater than 20 mm [23]. Analyzing the values of the diameters of the inhibition zones, it appears that the strains of *Staphylococcus aureus* ATCC and *Staphylococcus aureus* 0.68 are sensitive to the hydroethanolic extracts of *H. rotundifolia* powders. On the other hand, said extracts showed no activity against *Escherichia Coli* ATCC, *Pseudomonas aeruginosa* ATCC and *Pseudomonas aeruginosa* 363. In addition, the antibacterial activity manifested by the powders of *H. rotundifolia* against *Staphylococcus aureus* ATCC and *Staphylococcus aureus* 0.68 would be linked to the presence of flavonoids, tannins, coumarins, sterols and terpenes. Indeed, these above-mentioned compounds are recognized for their biological anti-radical and antibacterial activities [24].

### 3.5.2 Antibacterial activity in liquid medium

The antibacterial activity in liquid medium made it possible to determine the antibacterial parameters CMI, CMB of the hydroethanolic extracts of *H. rotundifolia* powders. The CMI, CMB and the CMB/CMI ratio of the extracts active against *Staphylococcus aureus* are recorded in Table 10.

The results in Table 10 present the quotient of the minimum bactericidal concentration (CMB) by the minimum inhibitory concentration (CMI) of the extracts of *H. rotundifolia* on the strains *Staphylococcus aureus* ATCC and *Staphylococcus aureus* 068. Indeed, the extracts for which this quotient is equal to 2, are called

bactericidal while those for which it is greater than 2 are called bacteriostatic. Thus, for the plant active on these two strains, it turns out that the sieved particles (Tamisat) have a bactericidal effect while the Broyat and Refus are both limited only to a bacteriostatic effect. The bactericidal nature of Tamisat from *H. rotundifolia* on the two strains of *Staphylococcus aureus* would be linked to its higher antioxidant potential. In fact, Tamisat demonstrated better antioxidant character than all other extracts studied. Also, Chamayou and colleagues [25] showed that after grinding, significant modifications in texture, structure, shape and surface can occur. The increase in the contact surface due to fragmentation increases the physical, chemical or enzymatic reactivity of the particles. Certain nutritional components can also see their availability increase following the destruction of cellular structures due to the increase in temperature due to the breakdown energy of molecules [26]. This is what would explain the significant biological activity of the finer particles and therefore of Tamisat compared to the Broyat and Refus.

## 4. CONCLUSION

The present work focused on a comparative study of powders of *Manihot esculenta* and *Heterotis rotundifolia* divided into three categories, namely: Broyat, Tamisat and Refus, the extracts of which were obtained by Soxhlet extraction, maceration and liquid-liquid contact. The phytochemical composition, the antioxidant activity with respect to the DPPH radical as well as the antibacterial activity of each of these categories of powders were highlighted. The results of the phytochemical screening carried out by color reactions and by means of TLC highlighted the presence of flavonoids, tannins, coumarins, sterols, terpenes and proteins in all the study extracts. Phenolic acids were observed only in *H. rotundifolia*. In addition, Broyat gives a greater number of extractable phytochemicals compared to Tamisat and Refus regardless of

the type of extraction solvent and the plant species used. The Tamisat contains the most compounds of interest and therefore exhibits greater antioxidant potential against DPPH and greater antibacterial activity against *Staphylococcus aureus*.

It appears at the end of this study that it would be preferable to directly use the powder obtained after grinding, i.e. the Broyat, in the context of a phytochemical study and the Tamisat for biological studies.

As a perspective, we plan to carry out this study on a larger number of plant species; use a better grinding system and several sieves of different mesh sizes in order to have Tamisat of different particle sizes as well as different extraction methods; characterize and quantify the identified active compounds using LC-MS and GC-MS analyses.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Chabrier JY. Medicinal plants and forms of use in phytotherapy. Doctoral thesis. Henri Poincaré University, Nancy 1 (France). 2010;165.
2. Azmir J, Zaidul ISM, Rahman MM, Sharif KM, Mohamed A, Shaena F. Techniques for extraction of bioactive compounds from plant materials: A review. *Journal of food Engineering*. 2013;117 (4):426-436.
3. Chirinos R, Rogez H, Campos D, Pedreschi R, Larondelle Y. Optimization of extraction conditions of antioxidant phenolic compounds from mashua (*Tropaeolum tuberosum* Ruis & Pavon) tubers. *Separation and purification technology*. 2007;55:217-225.
4. Ladiguina EY, Saffronitch LN, Otriachenkova VE, Balandina IA, Grinkevitch NI. Chemical analysis of medicinal plants. Edition Moskva, Vischaya Chkola. 1983;347.
5. Dohou N, Yamni K, Tahrouch S, Idrissi HLM, Badoc A, Gmira N. Phytochemical screening of an endemic Ibero-Moroccan *Thymelaea lithroides*. *Bull Soc Pharm Bordeaux*. 2003;142:61-78.
6. Békro YA, Mamyrbekova-Békro JA, Boua BB, Tra Bi FH, Ehilé EE. Ethnobotanical study and phytochemical screening of *Caesalpinia benthamiana* (Baill.) Herend. And Zarucchi (Caesalpinaceae). *Science and Nature*. 2007;4(2):217-225.
7. Mamyrbékova-Békro JA, Konan MK, Békro YA, Djié BMG, Zomi BTJ, Mambo V. Phytocompounds of the extracts of four medicinal plants of Côte d'Ivoire and assessment of their antioxidant potential by thin layer chromatography. *European Journal of Scientific Research*. 2008;24 (2):219-228.
8. Singleton VL, Ortofer R, Lamuela-Raventos RM. Analysis of total phenol and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Packed L(ed) Methods in enzymology*. Orlando Academic Press. 1999;152-178.
9. Arvouet-Grand, Colleagues. Standardization of propolis extract and identification of the main constituents. *Belgian pharmacy journal*. 1994;49(6):462-468.
10. Heilmer D, Vignadini P, Dini MG, Vincieri FF, Romani A. Antiradical activity and polyphenol composition of local Brassicaceae edible varieties. *Food chemistry*. 2006;99:464-469.
11. Brand-Williams W, Cuvelier ME, Berset C. Use of free radical method to evaluate antioxidant activity. *LWT -Lebensmittel-Wissenschaft und Technologie*. 1995; 28(1):25-30.
12. CASFM. European Society of Clinical Microbiology and Infectious Diseases Recommendation. 2020;181.
13. Wiegand I, Hilpert K, Hancock REW. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nature protocols*. 2007;3:163-175.
14. Guessennd N, Oussou KR, Koffi K, Dosso M. Determination of the antibacterial activity of natural substances from plants in the pharmacopoeia of Côte d'Ivoire, Technical sheet (2). *Pasteur Institute of Côte d'Ivoire*. 2005 ;18.

15. Dosso M, Faye-kette H. Quality control of the antibiogram in current practice: Experience of the bacteriology laboratory of the Pasteur Institute of Côte d'Ivoire. *The international bacteriologist*. 2000; special issue:53.
16. Zaiter A. Study of the phytochemistry of 12 plants from the Lorraine region according to the particle size of superfine powders. Doctoral thesis, University of Lorraine. 2017;147.
17. Etekpou DS. Phytochemical, antioxidant and some biological applications of *Heterotis rotundifolia* (Sm.) Jacq.-Fél. (Melastomataceae) from Côte d'Ivoire. Single PhD thesis, Nangui Abrogoua University, Abidjan. 2024;141.
18. Sivapriya M, Srinivas L. Isolation and purification of a novel antioxidant protein from the water extract of Sundakai (*Solanum torvum*) seeds. *Food Chemistry*. 2007;104:510-517.
19. Kolak U, Kabouche A, Öztürk M, Kabouche Z, Topçu G, Ulubelen A. Antioxidant diterpenoids from the roots of *Salvia barrelieri*. *Phytochemical Analysis*. 2009;20:320–327.
20. Tanoh SK, N'Gaman-Kouassi CC, Boa D, Mamyrbekova-Békro JA, Békro YA. Antioxidant activity of crude hydroethanolic and hydroacetic extracts of the organs of four medicinal plants from Côte d'Ivoire, *Nature and Technology Review*. 2019; 11(2):28-34.
21. Falleh H, Ksouri R, Abdelly C. Antioxidant activity and polyphenol content in the different organs of the wild artichoke *Cynara cardunculus*. *Institute of Arid Regions, Arid Regions Review, Tunisia*. 2006;344.
22. Rösch D, Krumbein A, Kroh LW. Antioxidant gallicoligomers, dimeric and trimeric proanthocyanidins from sea buckthorn (*Hippophaë rhamnoides*) pomace. *European Food Research and Technology*. 2004;219(6) :605 -613.
23. Ponce AG, Fritz R, Del AC, Roura SI. Antimicrobial activity of essential oil on the native microflora of organic Swiss chard. *Lebensmittel- Wissenschaft und Technolog*. 2003;36:679-684.
24. Koudoro YA, Bogninou G, Sophie R, Bossou AF, Arlette D, Agbangnan D et al. Secondary metabolites, antibacterial and antiradical activities of extracts of the trunk bark of *Acacia polyacantha* harvested in Benin. *International Journal of Advanced Research*. 2019;7(10): 1087-1092.
25. Chamayou A, Fages J. Broyage dans les industries agroalimentaires. *Technologie des pulvérulents dans les IAA, Lavoisier*. 2033;375-406.
26. Maaroufi C, Melcion JP, De Monredon F. Fractionation of pea flour with pilot scale sieving. I. physical and chemical characteristics of pea seed fractions. *Anim Feed Sci Technol*. 2000;85:61–78.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/122838>