



Effect of Level of Sulphur and Phosphorus on Yield and Quality of Soybean Grown in Inceptisol

Rohini Rajendra Madane ^{a++*}, Archana Yallappa Mahajan ^{b#}
and Pratima Tanaji Otari ^{c†}

^a Agriculture Polytechnic College, Phaltan, India.

^b PAHCA, Halgaon, India.

^c Shriram College of Agriculture, Paniv, India.

Authors' contributions

This work was carried out in collaboration among all authors. The research was carried out during the M.Sc. degree of the author RRM under Soil Science and Agriculture Chemistry Department at RCSM College of Agriculture, Kolhapur. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajsspn/2024/v10i4411>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124390>

Original Research Article

Received: 24/08/2024
Accepted: 27/10/2024
Published: 04/11/2024

ABSTRACT

The field experiment was conducted during *Kharif* 2017 at Post Graduate Research Farm, Rajarshree Chhatrapati Shahu Maharaj, College of Agriculture Kolhapur, with the view to study the effect of levels of sulphur and phosphorus on yield and quality of soybean grown in inceptisol. The soil of the experimental site was slightly alkaline in reaction, very low in available nitrogen, medium in available phosphorus and medium in available potassium and deficient in available sulphur. The field experiment was carried out in Factorial Randomized Block Design with three replications and sixteen treatments comprising four levels of sulphur (0, 20, 40, and 60 kg ha⁻¹) through elemental sulphur and four levels of phosphorus (0, 75, 100, and 125 kg ha⁻¹) through DAP. The result indicated that the significantly highest grain yield (27.20 q ha⁻¹, 25.04 q ha⁻¹ respectively) was

⁺⁺ Assistant Professor of Soil SSAC;

[#] Assistant Professor of Agronomy;

[†] Assistant Professor of Agricultural Microbiology;

*Corresponding author: Email: rohinimadane30@gmail.com;

recorded by application of sulphur and phosphorus at 60 kg ha⁻¹ (S₄) and 100 kg ha⁻¹ (P₃) than rest of the sulphur and the phosphorus levels. The application of (S₄, P₃) at 60 kg ha⁻¹ and 100 kg P ha⁻¹ showed significantly highest straw yield (38.85 q ha⁻¹, 35.50 q ha⁻¹ respectively) than rest of the combination of sulphur and Phosphorus levels. The quality parameter records highest percent of 1000 grain weight, oil content and oil yield were found in 60 kg S along with 100 kg P ha⁻¹ while the lowest seed yield, 1000 grain weight and oil content were found in control. Therefore, application of 60 kg S with 100 kg P appeared as the promising practice for obtaining higher seed yield and better quality of Soybean.

Keywords: Soybean; DAP; phosphorus.

1. INTRODUCTION

Soybean (*Glycine max* L. Merrill) is a leguminous crop and belongs to the sub-family papilionaceae of the family Fabaceae. Soybean is native to China. It was introduced into northern India in 1000 AD from Central China. In India, the states of Madhya Pradesh and Maharashtra are major producers of soybean, accounting for 49.08 lakh hectares and 65.83 lakh tonnes of the production. Soybean is nature's versatile plant; it supplies an abundant amount of protein (38-42 %), oil (18-22 %), and is enriched in unsaturated fatty acids and essential amino acids across a wide range of environmental condition. It gives 2-3 times more protein yield per hectare than other legumes or oilseed crops.

Kumar et. al. [1] reported that "both seed and stover yield of soybean increased significantly due to individual as well as combined application of phosphorus and sulphur. The combined application of sulphur with phosphorus produced the highest seed and stover yield of soybean".

Dhage et al. [2] indicated that "grain and straw yield, as well as the uptake of phosphorus and sulphur, increased with the rate of application of phosphorus and Sulphur individually and in various combinations. The application of various levels of phosphorus and sulphur influenced the quality parameters of soybean, including protein content and test weight. Available phosphorus in the soil increased with increasing levels of phosphorus. Similarly, available sulphur in the soil increased with increasing levels of sulphur. Sulphur is recognized as the fourth major nutrient after nitrogen, phosphorus and potassium. Sulphur plays an important role in many physiological functions. Soybean is a source of sulphur containing amino acids such as cysteine, cystine, methionine, which are essential for protein synthesis and also promote oil and chlorophyll synthesis. Soybean seed yield was increased by sulphur application".

Chausaria et al. [3] recorded "an improvement in protein and oil content due to application of phosphorus and sulphur in soybean crop. Sulphur improves the nutrient use efficiency of other plant nutrients, particularly nitrogen (N) and phosphorus (P). Sulphur provides direct nutritive value".

Phosphorus is essential for the growth and development of root nodules, their multiplication, and the effectiveness of root nodule bacteria. Phosphorus is the second major plant nutrient and is an indispensable element that plays a unique role in several plant metabolic and energy transformation processes. It hastens maturity and improves the quality of grain. Phosphorus acts as an energy source (ATP) for plants.

2. MATERIALS AND METHODS

The experiment was conducted during the kharif season in 2017 in an inceptisol at the Post Graduate Research Farm, Rajarshree Chhatrapati Shahu Maharaj College of Agriculture, Kolhapur, Maharashtra, India. The chemical composition indicates that the soil at the experiment site was medium deep (inceptisol) and deficient in available sulphur (7.42 kg ha⁻¹). Additionally, the soil was alkaline in reaction, very low in available nitrogen (137.98 kg ha⁻¹), medium in available phosphorus (15.33 kg ha⁻¹), and available potassium (181.30 kg ha⁻¹). The experiment was laid out in a factorial randomized block design with sixteen treatments and three replications. The treatments consisted of four levels of sulphur (0, 20, 40, 60 kg S ha⁻¹) and four levels of phosphorus (0, 75, 100, 125 kg P ha⁻¹). Sulphur and phosphorus were applied through elemental sulphur and diammonium phosphate (DAP). Plot-wise soil samples and treatment-wise plant samples were collected and analyzed as per standard procedures.

2.1 Methods are Used for Soil Analysis

Table 1. Methods are used for soil analysis

Sr. No.	Parameters	Methods
1.	pH (1:2.5)	Potentiometry
2.	EC	Conductivity
3.	Organic Carbon (%)	Walkley and Black Wet Oxidation
4.	Available N (Kg ha-1)	Alkaline permanganate (Distillation)
5.	Available P (Kg ha-1)	0.5 M NaHCO ₃ extract at pH 8.5 (Colorimetry)
6.	Available K (Kg ha-1)	Neutral normal ammonium acetate using (Flame photometer)
7.	Available S (Kg ha-1)	Turbidimetric

2.2 Methods are Used for Plant Analysis and Quality Traits

Table 2. Methods are used for plant analysis and quality traits

Sr. No.	Parameters	Methods
1.	Total N	Micro Kjeldahl (Diacid H ₂ SO ₄ :H ₂ O ₂ digestion)
2.	Total P	Vanadomolybdate yellow color method (triacid H ₂ SO ₄ :HNO ₃ :HClO ₄ digestion)
3.	Total K	Flame photometer
4.	Total S	Turbidimetric
5.	Protein content	Micro Kjeldahl (Diacid H ₂ SO ₄ :H ₂ O ₂ digestion)
6.	Oil content	Soxhlet extractor using ether as a solvent
7.	Germination Count	Quadratic method
8.	Chlorophyll	Spectrophotometer

3. RESULTS AND DISCUSSION

3.1 Grain Yield

The data in relation to effect of levels of Sulphur and Phosphorus on grain yield is presented in Table 3. Sulphur application to soybean up to 60 kg S ha⁻¹ increases the grain yield of soybean. Phosphorus application also successively and significantly increased grain yield of the crop over preceding levels up to 100 kg P ha⁻¹; however, the response at the higher level of 125 kg P ha⁻¹ was found to be almost at par with 100 kg P ha⁻¹. The significantly highest grain yield (27.20 q ha⁻¹) was recorded with the application of sulphur at 60 kg ha⁻¹ (S₄) compared to the other sulphur levels. The different levels of phosphorus application did not show much variation in grain yield. The application of Phosphorus at 100 kg ha⁻¹ (P₃) resulted in the significantly highest grain yield (25.04 q ha⁻¹). Decrease in yield (24.51 q ha⁻¹) at the P₄ level might be due to the higher concentration of phosphorus, which may cause toxic effect and nutrients imbalance.

The highest grain yield was recorded by treatment T₁₅ (S₄P₃) (29.65 q ha⁻¹). Yield attributes of the plant due to sulphur application are well known, considering the physiological role of sulphur in the plant body. Sulphur

enhances cell multiplication, elongation, expansion, and imparts a deep green colour to leaves due to better chlorophyll synthesis, resulting in increased food supply, essential replacement of amino acids, and relatively greater amount of dry matter accumulation. The supply of phosphorus to the soil might have accelerated cell division and enlargement, carbohydrate, fat metabolism, and respiration in the plant favouring increased growth and yield. The similar results were also recorded by Kumar et.al. [1], Dhage et al. [2], Mahmoodi et al. [4]. The yield components and seed yield of soybean were significantly influenced by phosphorus and sulphur application, as noted by Barman et. al. [5].

3.2 Straw Yield

The results indicated that the increase in levels of 'S' significantly increased the straw yield of soybean. The application of S at 60 kg ha⁻¹ (S₄) showed a significantly higher straw yield (38.85 q ha⁻¹) than the rest of the sulphur levels. Among the different levels of phosphorus, the highest straw yield was observed with P₃, i.e 100 kg P ha⁻¹ yielding 35.50 q ha⁻¹.

Considering the interaction effects of levels of S & P, it was observed that the straw yield of soybean was significantly influenced by the

interaction effect of S & P levels. Treatment T₁₅ (S₄P₃) and T₁₄ (S₄P₂) recorded the highest straw yield (42.00 and 41.96 q ha⁻¹ respectively) and were found to be statistically at par with each other and significantly superior to the over rest of the treatments. Application of sulphur might have increased the availability of nutrient to the soybean plant due to an improved nutritional environment, which in turn favourably influenced energy transformation, activation of enzymes, chlorophyll synthesis, and increased carbohydrate metabolism. Similar results were also obtained by Kumar et al. [1] Dhage et al. [2] and Mahmoodi et al. [4].

3.3 Effect of S and P on Quality Parameters of Soybean

1000 Grain Weight: The different levels of S & P and their interaction showed a significant effect on thousand grain weight. Application of sulphur at 60 kg ha⁻¹ recorded highest thousand grain weight (150.58 g), which was significantly superior to rest of the S levels. The application of different levels of phosphorus i.e. P₂, (146.12g) P₃ (147.54g) and P₄ (145.84g) did not show much variation in thousand grain weight and were found statistically at par with each other, but significantly superior to P₁ (140.64g) i.e. 0 k ha⁻¹ P.

Table 3. Effect of sulphur and phosphorus and their interaction on yield of soybean (q ha⁻¹)

Levels of sulphur (kg ha ⁻¹)	Levels of phosphorus (kg ha ⁻¹)				Mean	Treatments		
	P ₁ (0)	P ₂ (75)	P ₃ (100)	P ₄ (125)		S.E±	C.D. at 5%	
Grain								
S ₁ (0)	15.68	18.09	17.73	18.27	17.44			
S ₂ (20)	16.97	24.50	25.09	24.66	22.81	S	0.37	1.07
S ₃ (40)	20.24	27.26	27.68	27.24	25.61	P	0.37	1.07
S ₄ (60)	22.67	28.60	29.65	27.87	27.20	SxP	0.74	2.14
Mean	18.89	24.61	25.04	24.51	23.26			
Straw								
S ₁ (0)	18.90	24.10	25.00	26.32	23.58			
S ₂ (20)	24.13	35.96	36.00	35.27	32.84	S	0.49	1.43
S ₃ (40)	27.73	36.99	39.00	36.62	35.09	P	0.49	1.43
S ₄ (60)	32.32	41.96	42.00	39.11	38.85	SxP	0.99	2.87
Mean	25.77	34.75	35.50	34.33	32.59			




Fig. 1. Effect of different levels of sulphur and phosphorus interaction on grain yield of soybean




Fig. 2. Effect of different levels of sulphur and phosphorus interaction on straw yield of soybean

Table 4. Effect of sulphur and phosphorus on quality parameter of soybean.

Treatment	1000 grain weight (g)	Oil Per- cent (%)	Oil yield Kg ha ⁻¹	Protein content (%)	Chlorophyll content mg g ⁻¹	Germination count (%)
S ₁	140.52	17.79	310.99	39.91	0.133	97.17
S ₂	143.58	19.62	448.15	40.02	0.170	97.67
S ₃	145.47	20.19	517.53	40.05	.0210	97.25
S ₄	150.58	21.10	574.66	40.57	0.195	96.92
SE ±	0.62	0.08	8.17	0.20	0.002	0.8
CD at 5%	1.81	0.23	23.61	NS	0.005	NS
P ₁	140.64	19.10	365.79	40.27	0.168	97.33
P ₂	146.12	19.85	493.27	40.11	0.169	97.25
P ₃	147.54	19.96	504.69	40.33	0.180	97.08
P ₄	145.84	19.79	487.59	39.85	0.191	97.33
SE ±	0.62	0.08	8.17	0.20	0.002	0.8
CD at 5%	1.81	0.23	23.61	NS	0.005	NS

The interaction effect of sulphur and phosphorus indicated that the highest thousand grain weight of soybean was observed in treatment T₁₅ (S₄P₃) (155.00g) which was significantly superior to the rest of the treatment except T₁₄ (S₄P₂) and T₁₆ (S₄P₄), which recorded 151.51g and 153.08g, respectively. Sulphur plays an important role in growth by attributing metabolic and synthesis activities of oil, starch, and protein. The increase in thousand grain weight may be due to these result activities. Similar findings were reported by Mahmoodi et al. [4] and Dhage et al. [2].

Oil Per cent: The oil percentage in soybean grain increased with higher levels of sulphur. However, the highest oil percentage was observed at the in S₄ level (21.10%), i.e.

application of sulphur at 60 kg ha⁻¹, which was significantly superior to the other sulphur levels. The application of phosphorus at 100 kg ha⁻¹ (P₃) recorded the highest oil percentage (19.96 %) in soybean and was significantly superior to the control (P₁) (19.10 %) i.e. 0 kg ha⁻¹ P. But, P₂ (19.85 %) and P₄ (19.79 %) levels are statistically at par with the P₃ level. Further, oil per centage decreased beyond 100 kg ha⁻¹ P application, i.e. in P₄ level, 125 kg ha⁻¹ P (19.79 %).

This might be due to the higher concentration of P causing nutrient imbalance. Application of 60 kg S and 100 kg P ha⁻¹ recorded the highest oil percentage (21.41%) over all other treatments except treatment T₁₄ i.e. (21.35%), i.e.

application of 60 kg S and 75 kg P ha⁻¹. There was an improvement in quality parameters due to S and P application. Similar findings were reported by Harendra, Kumar and Das [6]. Furthermore, the interaction between P and S was significant. All the S levels increased oil content significantly at every level of P. The maximum oil content was recorded with a treatment combination of P2O5 and S. by Jarupula et al. [7]. The oil content of soybean was significantly influenced by phosphorus and sulphur application. As reported by Barman et al. [5]

Oil Yield: The oil yield of soybean was enhanced due to different levels of sulphur and phosphorus. The highest oil yield (574.66 kg ha⁻¹) was obtained at 60 kg sulphur per hectare, which was significantly superior to the other sulphur levels (0, 20, and 40 levels). The application of phosphorus at 100 kg ha⁻¹ showed highest oil yield (504.69 kg ha⁻¹), significantly superior to the control i.e. P₁ (365.79 kg ha⁻¹) 0 kg P ha⁻¹. However, the application of phosphorus at 75 kg ha⁻¹ (P₂) (493.27 kg ha⁻¹) and 125 kg ha⁻¹ (P₄) (487.59 kg ha⁻¹) was statistically at par with 100 kg ha⁻¹ phosphorus (P₃). The interaction effect of different sulphur and phosphorus levels found to be significant among the various treatment combinations. The treatment T₁₅ which involved the application of sulphur at 60 kg ha⁻¹ with 100 kg ha⁻¹ phosphorus, recorded the highest oil yield (634.80 kg ha⁻¹), followed by treatment T₁₄ (S₄P₂), which involved 60 kg S ha⁻¹ with 75 kg ha⁻¹ P (610.60 kg ha⁻¹) these treatments were statistically at par with each other and significantly superior to the other treatments.

This improvement in oil yield is attributed to the enhancement in quality parameters (1000 grain weight, protein and oil content) due to application of sulphur and phosphorus. The improvement of sulphur and phosphorus through the growing media to the soybean crop is also noted. Additionally, the enzyme acetic thiokinase, which is involved in the conversion of acetyl Co-A to malonyl Co-A and this activity of thiokinase is governed by sulphur, potentially resulting in increased oil yield. Similar findings were reported by Harendra Kumar and Das, [6] The maximum oil content, oil yield were recorded with a treatment combination of P2O5 and S, as reported by Jarupula et al. [7].

Protein Content (%): “The different levels of sulphur, phosphorus and their interactions did

not showed much more variation in protein content and were found non significant in protein per cent. Similar results were obtained” by Dhage et al. [2].

Total Chlorophyll Content: The different levels of S and P and their interaction showed a significant effect on chlorophyll content. The level of S₃ i.e. application of sulphur at 40 kg ha⁻¹ recorded the highest chlorophyll content (0.210 mg g⁻¹) and was found to be significantly superior over the rest of the sulphur levels. The application of phosphorus (P₄) at 125 kg ha⁻¹ showed the highest chlorophyll content (0.191 mg g⁻¹) and was found to be statistically at par with P₃ (0.180 mg g⁻¹) and significantly superior over the control i.e. P₁ 0 kg P ha⁻¹ [4,8].

The interaction effect of sulphur and phosphorus indicated that the highest chlorophyll content of soybean was observed in treatment T₁₂ (S₃P₄) (0.236 mg g⁻¹), which was significantly superior over the rest of the treatments. The chlorophyll content of the leaf is influenced by both N and S nutrition of the crop. Application of 40 kg S ha⁻¹ significantly increased the chlorophyll content of the leaves. Higher rates of applied sulphur did not further increase the chlorophyll content of the leaves. Similar results were obtained by Reddy and Ganeshamurthy [9].

Germination Per cent: The different levels of sulphur, phosphorus and their interactions were found non significant in germination per cent.

4. CONCLUSION

The application of sulphur at 60 kg ha⁻¹ and phosphorus at 100 kg ha⁻¹ showed a significant effect in increasing the grain yield (27.20 q ha⁻¹) and straw yield (25.04 q ha⁻¹) of soybean. The quality parameter records highest percent of 1000 grain weight, oil content and oil yield were found in 60 kg S along with 100 kg P ha⁻¹ while the lowest seed yield, 1000 grain weight and oil content were found in control. Therefore, application of 60 kg S with 100 kg P appeared as the promising practice for obtaining higher seed yield and better quality of Soybean.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar S, Wani JA, Lone BA, Singh P, Dar ZA, Qayoom S, Fayaz A. Effect of different levels of phosphorus and sulphur on seed and stover yield of soybean [*Glycine max* (L.) Merrill] under 'Eutrochrepts'. Asian Research Journal of Agriculture. 2017; 5(1):1-7.
2. Dhage SJ, Patil VD, Ptange MJ. Effect of various levels of phosphorus and sulphur on yield, plant nutrient content, uptake and availability of nutrients at harvest stages of soybean. International Journal of Current Microbiological and Applied Sciences. 2014;3:833-844.
3. Chaurasia, Amit Kumar GP. Richharia, Shridha Chaurasia. Response of Soybean (*Glycine max*) to doses and sources of sulphur. Indian J. of Agricultural Sciences. 2009;79(5):356-358.
4. Mahmoodi B, Amir A, Mosavi Morteza, Daliri S, Namdari M. The evaluation of different values of phosphorus and sulfur application in yield, yield components and seed quality characteristics of soybean (*Glycin max* L.) Advances in Environmental Biology. 2013;7(1):170-176.
5. Barman DK, Sarkar SK, Salam MA, Paul SK. Response of Yield and Quality of Soybean (cv. BARI Soybean-6) to Phosphorus and Sulphur Fertilization under Old Brahmaputra Floodplain Soil of Bangladesh. Agricultural Science Digest. 2022;1. DOI: 10.18805/ag.DF-408 (1-5).
6. Kumar and Das K Yadav. Effect of phosphorus and sulphur levels on growth, yield and quality of Indian mustard cultivars. Indian Journal Agronomy. 2007;52:154-157.
7. Jarupula Suman BS, Dwivedi AK, Dwivedi Pandey SK. Interaction Effect of Phosphorus and Sulphur on Yield and Quality of Soybean in a Vertisol. Int. J. Curr. Microbiol. App. Sci. 2018;7(03):152-158.
8. Teotia US, Mehta VS, Ghosh D, Srivastava PC. Phosphorus-Sulphur interaction in moongbean (*Vigna radiata* L. Wilczek): I. Yield, phosphorus and sulphur contents. Legume Res. 2000;23:106-109.
9. Reddy KS, Ganeshamurthy AN. Effect of Integrated use of farmyard manure and sulphur in soybean and wheat cropping system on nodulation, dry matter production and chlorophyll content of soybean on swell-shrink soils in central India. Journal of Agronomy and Crop Science. 2000;185:91-97.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124390>