



Effect of Sulphur and Zinc on Growth and Yield of Indian Mustard [Brassica juncea (L.)]

Anil Kumar ^{a*}, Bhagwan Suman ^a, Om Prakash Gurjar ^a, Chandrakanta Jakhar ^a and Indra Raj Yadav ^b

^a Department of Agriculture, Faculty of Agriculture and Veterinary Science, Mewar University Gangrar, Chittorgarh Rajasthan, 312901, India.

^b Department of Soil Science and Agricultural Chemistry, Rajmata Vijayaraje Scindia Krishi Vishwavidyalaya, Gwalior, 474002, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/arja/2024/v17i4601>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126732>

Short Research Article

Received: 16/09/2024
Accepted: 21/11/2024
Published: 26/11/2024

ABSTRACT

A field experiment was conducted at experimental farm, Department of Agronomy, Faculty of Agriculture and Veterinary Sciences, Mewar University Gangrar, Chittorgarh (Rajasthan) during Rabi season of 2023-24 to effect of Sulphur and Zinc on growth and yield of mustard (*Brassica juncea* L.), variety "Bio-902" was used in this study. The experiment was laid out in factorial randomized block design with three replications. Maximum growth attributes such as, plant height (146.30 and 143.90 cm), number of branches per plant (8.28 and 8.05) and dry matter accumulation (326.17 and 324.32 g/plant) and yield attributes such as number of siliqua per plant (285.20 and 282.49), number of seed per siliqua (13.20 and 13.10), seed yield (20.40 and 20.18 q/ha), stover

*Corresponding author: E-mail: anilmehara07@gmail.com;

Cite as: Kumar, Anil, Bhagwan Suman, Om Prakash Gurjar, Chandrakanta Jakhar, and Indra Raj Yadav. 2024. "Effect of Sulphur and Zinc on Growth and Yield of Indian Mustard [*Brassica Juncea* (L.)]". Asian Research Journal of Agriculture 17 (4):908-13. <https://doi.org/10.9734/arja/2024/v17i4601>.

yield (69.84 and 69.58 q/ha) and maximum net return (66449.75 and 66557.00 Rs/ha) was recorded with combine application of 60 kg/ha Sulphur and 7.5 kg/ha Zn. Therefore, it was concluded that application of 40 kg Sulphur and 7.5kg zinc ha⁻¹ found suitable to produce good yield of Indian mustard. Maximum values of growth parameters and yield also recorded with application 60 Sulphur kg ha⁻¹ and 7.5 zinc kg ha⁻¹.

Keywords: Sulphur; zinc; indian mustard; yield; profitability.

1. INTRODUCTION

“Indian mustard belongs to family Cruciferae family, grown during rabi season in Northern parts of India. In India the area under mustard crop is 7.32 million ha producing about 10.02 million tonnes of seeds with an average productivity of 1348 kg ha⁻¹ and Area under cultivation in Rajasthan was 3.85 million ha, having production of 4.63 million tones and productivity of 1412 kg ha⁻¹ [1]. “The oil content in mustard seeds varies from 37-49 per cent, the seeds are highly nutritive containing 38-57 per cent eruric acid, and 27 per cent oleic acid” (Bhowmik et al., 2014).

“Micronutrients hold a dominant and significant role in the growth and metabolic operations of oilseed crops (Indian Mustard) is highly sensitive to sulphur (S), zinc (Zn), iron (Fe) and nitrogen (N) deficiency, thereby resulting in small leaves, chlorosis and dwarfness, leading to a decline in its productivity. Zinc and Fe are important micronutrients required in trace amounts by humans, animals, and plants” [2]. “Zinc being one of the essential micro-nutrients, plays significant role in various enzymatic and physiological activities of the plant system. It is also essential for photosynthesis and N-metabolism. It is important for the stability of cytoplasmic ribosomes, cell division, dehydrogenase, proteinase and peptidase enzymes and also helps in the synthesis of protein and carotene” [3].

“Sulphur is the most vital nutrients for growth and development oil seeds. It is considered to be the fourth important essential nutrient after nitrogen, phosphorus and potassium for the plant growth. In order to develop a sound fertilizer recommendation for improvement in the mustard yield, it is necessary to find out Sulphur and zinc requirements in mustard crop” (Mohiuddin et al. 2011).

2. MATERIALS AND METHODS

A field experiment was conducted during Rabi season of 2023-24 at experimental farm,

Department of Agronomy, Faculty of Agriculture and Veterinary Sciences, Mewar University Gangrar, Chittorgarh (Rajasthan). Soil of the experimental field was sandy loam in texture, saline in reaction with a pH value of 7.6, poor in organic carbon (0.16%), deficient in available zinc (0.48 ppm) and iron (1.2 ppm) low in available nitrogen (176 kg/ha) and phosphorus (20.2 kg/ha) but medium in available potassium (320 kg/ha). The experiment was laid out in factorial randomized block design with three replications where treatment level-I, S₁- 0 kg/ha, S₂- 20, S₃- 40 and S₄- 60 kg/ha sulphur and level second Zn₁-0, Zn₂- 0.25, Zn₃- 5.0 and Zn₄- 7.5 kg/ha Zn. The required quantities of fertilizers as per treatments were applied. The doses of NPK were applied in the form of urea, diammonium phosphate, murate of potash respectively. The growth attributes and yield parameters were calculated from output from the field. The profitability and productivity was calculated from field preparation to threshing operation and the cost of the seed yield was taken as per market rate.

3. RESULTS AND DISCUSSION

3.1 Growth Attributes

Data pertaining to plant height of mustard as affected by different levels of sulphur and zinc presented in Table 1. The result clearly stated from the data that plant height increased as the dose of sulphur increased up to 60 kg ha⁻¹ and Zn 7.5 kg/ha at all growth stages. Plant height of mustard was maximum with application of 60 kg sulphur ha⁻¹ being at par with 40 kg sulphur ha⁻¹ at 60 DAS and with 40 and 20 kg sulphur ha⁻¹ at 90 DAS. The minimum value of plant height was observed with no sulphur application. Response regarding plant height was higher with 7.5 kg zinc ha⁻¹ being at par with 5 kg and 2.5 kg zinc ha⁻¹ and found significantly superior over control at 60 and 90 DAS. The results of present investigation are in agreement with the finding of Singh et al. [4] Rajput et al. [5]. Number of branches plant⁻¹ of mustard was maximum with application of 60 kg sulphur ha⁻¹. The magnitude

changes the number of branches plant⁻¹ in terms of percentage of 44.89 and 47.56 % at 60 DAS and 14.02 and 30.90 % at harvest, to 20 kg sulphur ha⁻¹ and control respectively. Number of branches plant⁻¹ due to application of 7.5 kg zinc ha⁻¹ in the tuning of percentage viz. 25.71 and 10.00 % at 60 DAS and 34.07 and 36.71 % at 90 DAS over control. Dry matter accumulation was maximum with application of 60 kg sulphur ha⁻¹ in mustard being at par with 40 kg sulphur ha⁻¹ and found significantly superior 20 kg sulphur ha⁻¹ and control at 60 and 90 DAS. The percentage increment in dry matter accumulation due to application of 60 kg sulphur ha⁻¹ was 20.61 at 60 DAS, 10.89% at DAS by control. Response regarding of dry matter accumulation was higher with 7.5 kg zinc ha⁻¹ being at par with 5 kg zinc ha⁻¹ and found significantly superior 2.5 kg ha⁻¹ and over control at 60 and 90 DAS. In respect to percentage dry matter accumulation changes with the application of 7.5 kg zinc ha⁻¹ as 19.39% at 60 DAS and 19.34% at harvest over control. Dubey et al. [6], Kobrai et al. [2], Rabari et al. [7], Rajput et al. [5], Mathpal et al. [8] also reported the similar results.

3.2 Yield Attributes and Yield

The by different levels of sulphur and zinc presented in Table 2 and graphically illustration in Fig. 1. The highest number of siliquee plant⁻¹ (285.2) was recorded with application of 60 kg sulphur ha⁻¹ being at par with 40 kg sulphur ha⁻¹ and 20 kg sulphur ha⁻¹ and minimum value of number of siliquee plant⁻¹ (246.0) was observed with no sulphur

application. The maximum number of siliquee plant⁻¹ (282.49) was observed with 7.5 kg zinc ha⁻¹ being at par with 5 kg and 2.5 kg zinc ha⁻¹ and found significantly superior over control. The number of seeds siliquee⁻¹ of mustard was maximum with application of 60 kg sulphur ha⁻¹ being at par with 40 kg sulphur ha⁻¹ and found superior 20 kg sulphur ha⁻¹ and over control. The maximum number of seeds siliquee⁻¹ was observed with 7.5 kg zinc ha⁻¹ being at par with 5 kg and 2.5 kg zinc ha⁻¹ and found significantly superior over control. The result revealed that the seed yield increased as the doses of sulphur increased up to 60 kg sulphur ha⁻¹. The seed yield of mustard was maximum (20.40 q ha⁻¹) with application of 60 kg sulphur ha⁻¹. The minimum seed yield (16.50 q ha⁻¹) was noticed with control. The maximum yield (20.18 q ha⁻¹) obtained with the application of 7.5 kg zinc ha⁻¹. The data that stover yield increased as the dose of sulphur increased up to 60 kg sulphur ha⁻¹. The stover yield of mustard was maximum (69.84 q ha⁻¹) with application of 60 kg sulphur ha⁻¹ being at par with 40 kg sulphur ha⁻¹ and found significantly superior over 20 kg sulphur ha⁻¹ and control. The Stover yield higher with 7.5 kg zinc ha⁻¹ being at par with 5 kg zinc ha⁻¹ and found significantly superior over 2.5 kg ha⁻¹ and control. The number of siliquee plant⁻¹, number of seed siliquee⁻¹, test weight, seed yield, and stover yield and harvest index were increased with increasing rate of sulphur from control to 45 kg sulphur ha⁻¹ in mustard Debnath et al. [9], Rabari et al. [7], Jat et al. [10], Kour et al. [11], Ray et al. [12], Bhagwat et al. [13], Kumar et al. [14] also reported the similar results.

Table 1. Effect of sulphur and zinc levels on growth attributes of mustard

Treatments	Plant height (cm)			Number of branches per plant			Dry matter accumulation/plant (g)		
	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS
Levels of sulphur (kg ha⁻¹)									
0	15.2	84.3	127.7	2.04	5.00	5.75	14.56	121.40	269.79
20	19.1	90.8	139.0	2.06	6.20	7.13	15.00	133.31	296.48
40	20.5	94.2	145.9	2.08	7.00	8.05	15.45	144.46	321.28
60	20.8	96.5	146.3	2.09	7.20	8.28	15.73	146.78	326.17
S. Em. ±	0.53	1.85	2.89	0.06	0.12	0.14	0.37	2.77	7.76
CD (P=0.5) %	NS	5.35	8.34	NS	0.36	0.41	NS	8.01	22.41
Levels of zinc (kg ha⁻¹)									
0	16.8	84.9	122.2	2.04	5.20	5.98	14.73	122.29	271.75
2.5	19.6	88.1	136.3	2.06	6.30	7.25	14.91	133.70	297.62
5.0	20.7	90.9	140.3	2.08	6.90	7.94	15.27	144.01	320.03
7.5	20.8	91.0	143.9	2.09	7.00	8.05	15.83	145.94	324.32
S. Em. ±	0.53	1.85	2.89	0.06	0.12	0.14	0.37	2.77	7.76
CD (P=0.5) %	NS	5.35	8.34	NS	0.36	0.41	NS	8.01	22.41

Table 2. Effect of sulphur and zinc levels on yield attributes and yield of mustard

Treatments	Number of siliquae per plant	Number of seed per siliquae	Seed yield 9q/ha)	Stover yield (q/ha)
Levels of sulphur (kg ha⁻¹)				
0	246.0	11.20	16.5	58.19
20	268.0	12.40	18.2	63.76
40	281.2	13.00	19.9	68.91
60	285.2	13.20	20.4	69.84
S. Em. ±	6.17	0.25	0.36	1.53
CD (P=0.5) %	17.82	0.71	1.06	4.45
Levels of zinc (kg ha⁻¹)				
0	251.00	11.25	16.7	58.48
2.5	269.00	12.60	18.3	63.9
5.0	278.00	12.85	19.8	68.74
7.5	282.49	13.10	20.18	69.58
S. Em. ±	6.17	0.24	0.37	1.54
CD (P=0.5) %	17.82	0.71	1.06	4.45




Fig. 1. Effect of sulphur and zinc levels on yield attributes and yield of mustard

3.3 Economics

The economics of mustard as influenced by different levels of sulphur and zinc presented in Table 3 and graphically illustration. The cost of cultivation was calculated for all the treatment combinations. The maximum cost of cultivation of 31063 ha⁻¹ was noted with 60 kg sulphur ha⁻¹ and 7.5 kg zinc ha⁻¹. The maximum gross return of 103492 ha⁻¹ was also noted with 60 kg sulphur

ha⁻¹ and 7.5 kg zinc ha⁻¹. However, the maximum benefit cost ratio of 2.42 was recorded with treatment combination of 40 kg sulphur ha⁻¹ and 7.5 kg zinc ha⁻¹. This is due to the increased net return in corresponding to the cost of cultivation under the treatment combination. The finding are in close conformity with findings of Verma et al. [15], Dubey et al. [6], Chaurasiya et al. [16] and Komatineni, [17].

Table 3. Effect of sulphur and zinc levels on Economics of mustard

Treatments	Economics	
	Net returns (₹ ha ⁻¹)	B:C
Levels of sulphur (kg ha⁻¹)		
0	53000.00	2.10
20	59409.00	2.21
40	65183.50	2.31
60	66449.75	2.32
Levels of zinc (kg ha⁻¹)		
0	52908.5	2.02
2.5	59500.5	2.20
5.0	65076.25	2.31
7.5	66557.00	2.32

4. CONCLUSION

The findings of present investigation revealed that application of 40 kg sulphur and 7.5kg zinc ha⁻¹ found suitable to produce good yield of Indian mustard. Maximum values of growth parameters, yield also recorded with application 60 sulphur kg ha⁻¹ and 7.5 zinc kg ha⁻¹. On the basis of economics, it may be concluded that application of 40 kg sulphur and 7.5 kg zinc ha⁻¹ in mustard was found more remunerative which recorded the Benefit cost ratio of 2.31 and 2.32, respectively.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Anonymous. Directorate of Economics and Statistics, Department of Agriculture and Cooperation (DAC). Annexure-3.1, p. 163 and Annexure-6.1 p. 169 on Mustard Crop Survey Report; 2021.
- Kobrai S, Shamsi K, Rasekhi B. Micronutrient fertilizer and soybean nutrition. *Annals of Biological Research*. 2011;2:468-475.
- Kumar M, Singh PK, Yadav KG, Chaurasiya A, Yadav A. Effect of nitrogen and sulphur nutrition on growth and yield of Indian mustard (*Brassica juncea* L.) in western UP. *Journal of Pharmacognosy and Phytochemistry*. 2017;25(1):445-448.
- Singh R, Singh SB, Manhas SS, Kumar A. Effect of different levels of sulphur and varieties on growth, yield and quality of Indian mustard. *International Journal of Plant Science*. 2012;7(2):290-294.
- Rajput RK, Singh S, Varma J, Rajput P, Singh M, Nath S. Effect of different levels of nitrogen and sulphur on growth and yield of Indian mustard (*Brassica juncea* (L.) Czern and Coss.) in salt affected soil. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(1):1053-1055.
- Dubey SK, Trapathi SK, Singh Bhagwan. Effect of sulphur and zinc levels on growth, yield and quality of mustard (*Brassica juncea* L. Czern & Coss.). *Journal of Crop Science and Technology*. 2013;2:159-166.
- Rabari KV, Patel KM, Patel B, Desai NH. Influence of ferrous sulphate and zinc sulphate on pod yield of groundnut. *International Journal of Agriculture Sciences*. 2018;10:5725-5726.
- Mathpal B, Dharvankar M, Singh G. Role of sulphur in improving growth and yield of Indian mustard (*Brassica juncea* L.). *The Pharma Innovation Journal*. 2023;12(6):3431-3436.
- Debnath A, Kole SC, Sukhim JM. Evaluation of the efficacy of different sulphur amendments and sulphur oxidizing bacteria in relation to its transformation in soil and yield of mustard (*Brassica juncea*). *Research on Crops*. 2014;15(3):578-584.
- Jat G, Sharma KK, Choudhary R. Effect of FYM and mineral nutrients on yield, content and uptake of nutrients in mustard. *Annals of Agricultural Research*. 2015; 34(3).
- Kour S, Gupta M, Bharat R, Sharma V. Effect of zinc and boron on yield, nutrient

- uptake and economics of mustard (*Brassica juncea* L.) cropping sequence. Bangladesh Journal of Botany. 2017; 46(2):817-821.
12. Ray K, Sengupta K, Pal AK, Banerjee H. Effects of sulphur fertilization on yield, S uptake and quality of Indian mustard under varied irrigation regimes. Plant, Soil and Environment. 2015;61(1):6-10.
 13. Bhagwat GJ, Gokhale DN, Waghmare PK, Bhalerao. Effect of micronutrients application and economics of soybean (*Glycine max* L.) crop. International Journal of Current Microbiology Applied Sciences. 2018;6:1860-1865.
 14. Kumar V, Singh RK, Kumar DM. Effect of farm yard manure and Sulphur on production of Indian mustard: A review. Journal of Pharmacognosy and Phytochemistry. 2019;8(3):2290-2294.
 15. Verma CK, Prasad K, Yadav DD. Studies on response of sulphur, zinc and boron levels on yield, economics and nutrients uptake of mustard (*Brassica juncea* L.). Crop Research. 2012;44(1/2):75-78.
 16. Chaurasiya A, Singh S, Singh V, Singh N, Singh A. Effect of nitrogen and sulphur nutrition on interaction effect, quality parameters, nutrient content or uptake & economics of Indian mustard (*Brassica juncea* L.) in western UP. IJCS. 2019;7(1):787-791.
 17. Komatineni A. Effect of potassium & zinc on growth & yield of mustard. Thesis for: M.Sc (Ag.) AGRONOMY, SHUSTS, Prayagraj (U.P); 2023.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/126732>