



# Impact of New Herbicides on Weed Population and Weed Control Efficiency in Scented and Hybrid Rice (*Oryza sativa* L.)

Jaykar Singh <sup>a\*</sup>, V.K. Verma <sup>a++</sup>, D.D. Yadav <sup>a++</sup>,  
Sanjiv Kumar <sup>a++</sup>, M.Z. Siddqui <sup>a++</sup>, Sarvesh Kumar <sup>b++</sup>,  
Anil Kumar <sup>c++</sup>, Vineet Dheer <sup>a</sup>, Krishna Kumar Singh <sup>c</sup>  
and Suryabhan <sup>a</sup>

<sup>a</sup> Department of Agronomy, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur (U.P.) 208002, India.

<sup>b</sup> Department of Soil Conservation and Water Management, Chandra Shekhar Azad university of Agriculture and Technology, Kanpur (U.P.) 208002, India.

<sup>c</sup> Department of Soil Science and Agricultural Chemistry, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur (U.P.) 208002, India.

## Authors' contributions

This work was carried out in collaboration among all authors. Author JS has designed, conceived and performed the field experiment and wrote the original draft, reviewed and edited the manuscript. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jsrr/2024/v30i112644>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125907>

**Original Research Article**

**Received: 05/09/2024**

**Accepted: 09/11/2024**

**Published: 29/11/2024**

<sup>++</sup>Professor;

\*Corresponding author: E-mail: [jaykar6697@gmail.com](mailto:jaykar6697@gmail.com);

**Cite as:** Singh, Jaykar, V.K. Verma, D.D. Yadav, Sanjiv Kumar, M.Z. Siddqui, Sarvesh Kumar, Anil Kumar, Vineet Dheer, Krishna Kumar Singh, and Suryabhan. 2024. "Impact of New Herbicides on Weed Population and Weed Control Efficiency in Scented and Hybrid Rice (*Oryza Sativa* L.)". *Journal of Scientific Research and Reports* 30 (11):1182-90. <https://doi.org/10.9734/jsrr/2024/v30i112644>.

## ABSTRACT

Field experiments were carried out to study the effect of herbicide and micronutrient spray on different varieties of rice (*Oryza sativa* L.) at Students' Instructional farm, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur (U.P.), during *Kharif* season 2022 and 2023. The experiment was laid out in split-split plot design with three replications and 24 treatment combinations comprising of two varieties viz. Arize 6444 Gold ( $V_1$ ) and PB 1509 ( $V_2$ ) were kept in main plots and three herbicidal treatments viz. Control ( $W_1$ ), Bispyribac-sodium 10% SC @ 20 g a.i.  $ha^{-1}$  at 25 DAT ( $W_2$ ), Triafamone 20% + Ethoxysulfuron 10 WG @ 225g  $ha^{-1}$  at 25 DAT ( $W_3$ ) were kept in Sub plots and four micronutrients application viz. Zinc Sulphate 0.5% + Ferrous Sulphate 1% foliar spray at 45 and 60 DAT ( $M_1$ ), Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% foliar spray at 45 and 60 DAT ( $M_2$ ), Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% + Borax 0.2% foliar spray at 45 and 60 DAT ( $M_3$ ), Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% + Borax 0.2% + Silicon 0.1% foliar spray at 45 and 60 DAT ( $M_4$ ) were kept in Sub-sub plots. Result revealed that Arize 6444 Gold variety had better weed suppression compare to PB 1509. On other side, herbicidal application with Triafamone 20% + Ethoxysulfuron 10 WG @ 225g  $ha^{-1}$  spray at 25 DAT, provided the most effective control against weed flora, weed fresh weight, dry weight and weed control efficiency, among all the growth stages. However, foliar spray of Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% + Borax 0.2% + Silicon 0.1% foliar spray at 45 and 60 DAT had a notable impact on reducing dry weed weight and significant influenced on weed control efficiency.

**Keywords:** Foliar spray; herbicide; micronutrient; rice; weed control efficiency; weeds.

## 1. INTRODUCTION

Rice (*Oryza sativa* L.) is a globally important cereal crop, particularly known for its wider adaptability in wet and submerged growing conditions. As a member of the Poaceae family, rice serves as a staple food for over 60% of the global population, making it a cornerstone of agricultural economies worldwide. The global cultivation area of rice approximately 165.21 million hectares, with an annual production of 509.26 million tonnes and an average productivity of 4600 kg per hectare during the 2020-21 period (USDA 2021). In India, it covering 46.38 million hectares and producing 130.29 million tonnes of grain, with an average productivity of 2809 kg  $ha^{-1}$  (DAC & FW 2022). Rice primarily consumed by humans, playing a critical role in global food security. It serves as the main source of energy for a significant portion of the world's population, particularly in Asia. The grain is highly caloric, consisting predominantly of 75% starch, which makes it an efficient energy source. It also contains 6-7% protein, 2-2.5% fat, 0.8% cellulose and 5-9% ash, making it a valuable component of a balanced diet (Resurreccion and Juliano 1981). Aromatic rice varieties, known for their distinct flavours and consumer appeal, play an important role in global agriculture. The key compound responsible for their unique 'popcorn-like' aroma is 2-acetyl-1-pyrroline (2-AP), identified as the

main flavour component (Yoshihashi et al. 2002). Despite its significance, rice cultivation faces challenges, particularly from weeds, which compete for essential resources and can host pests and diseases. Herbicides like Bispyribac-sodium and Council (Triafamone 20% + Ethoxysulfuron 10 WG) have proven effective in controlling weeds, enhancing rice crop yield and quality. In addition, micronutrients such as iron, zinc, manganese, silicon and boron are vital for the healthy growth and development of rice plants, contributing to chlorophyll synthesis, protein production, disease resistance and pollen viability. This study focuses on understanding the combined effects of herbicides and micronutrient application on transplanted rice. By investigating their impact on growth, yield and quality, the research aims to provide valuable insights for optimizing agronomic practices, ultimately improving rice productivity and contributing to global food security.

## 2. MATERIAL AND METHODS

A field experiment was conducted at Students' Instructional farm, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur (U.P.). It is situated at an elevation of 125.9 m above the mean sea level 26°41'48" North latitude and 80°23'21" East longitude. It is situated in the alluvial belt of Indo-Gangetic Plain

in the Central part of Uttar Pradesh. Geographically, this experimental site falls under semi-arid with hot summer, moderate rainfall and cold winter. The average annual precipitation of the area varies from 800 to 900 mm with a mean annual precipitation of about 868 mm, mainly through south-west monsoon rains confined within June to last week of September with occasional frost and shower in winter season from North-East monsoon during December and January. The weekly mean minimum and maximum temperature during the entire crop season ranged from 9.90 to 11.40 °C and 43.90 to 42.20 °C, total rainfall received was 836.20 mm and 747.80 mm. The experimental soil was sandy loam in texture with sand 63.24%, clay 13.08% and 23.68% silt, a bulk density of 1.34 g cm<sup>-3</sup> (0-15 cm), pH of 7.64, electrical conductivity of 0.45 dSm<sup>-1</sup>, organic carbon at 0.49%, low in available nitrogen (221.41 kg ha<sup>-1</sup>), available Zinc (0.70 ppm) and medium in Phosphorus (13.04 kg ha<sup>-1</sup>), potassium (214.84 kg ha<sup>-1</sup>) and available Fe (6.74 ppm). The Experiment consisted of 24 treatment combinations comprising of two varieties viz. Aromatic rice Arize 6444 Gold (V<sub>1</sub>) and PB 1509 (V<sub>2</sub>) were kept in main plots and three herbicidal treatments viz. Control (W<sub>1</sub>), Bispyribac-sodium 10% SC @ 20 g a.i. ha<sup>-1</sup> at 25 DAT (W<sub>2</sub>), Triafamone 20% + Ethoxysulfuron 10 WG @ 225g ha<sup>-1</sup> at 25 DAT (W<sub>3</sub>) were kept in Sub plots and four micronutrients applications viz, Zinc Sulphate 0.5% + Ferrous Sulphate 1% foliar spray at 45 and 60 DAT (M<sub>1</sub>), Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% foliar spray at 45 and 60 DAT (M<sub>2</sub>), Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% + Borax 0.2% foliar spray at 45 and 60 DAT (M<sub>3</sub>), Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% + Borax 0.2% + Silicon 0.1% foliar spray at 45 and 60 DAT (M<sub>4</sub>). The treatments were arranged in a split-split plot design with three replications and plot size 4m width and 5m length. A common procedure was followed in raising seedlings (seed rate- 15 and 45 kg ha<sup>-1</sup> for hybrid and Basmati, respectively) in the seedbed then uprooting of rice Healthy seedlings, around 25 days old and carefully transplant seedlings into the main field at a spacing of 20x10 cm as row to row and plant to plant using 2-3 seedlings per hill in well prepared puddle field. Weed management by herbicide was spray as post-emergence at 25 DAT with the help of a knapsack sprayer attached with flat fan nozzle using 600 Liters of water per ha. The plot wise crop was harvested by hand with sickles at the time when upper portion of spikelet's looked

straw colored and grain has hard, containing 15-18% moisture and left for three days for sun drying then threshing was done manually with the assistance of sticks. All the application fertilizers were done separately in each plot as per treatments. Application of recommended dose of fertilizer, half dose of nitrogen and a full dose of phosphorus and potash were given as basal dose and remaining nitrogen applied as a top dressing in two equal splits and micronutrient are applied by foliar spray as per treatments. Observed parameters like total number of weed flora (m<sup>-2</sup>) at 45, 60 and 90 DAT. From each plot at three places randomly using a quadrat of 50cm x 50cm (0.25 m<sup>2</sup>) and multiplied by four to convert into m<sup>-2</sup>. Weed fresh weight and weed dry weight (g m<sup>-2</sup>). Weed sample taken from each plot, for the observation of fresh and dry weight of weed was recorded at 30, 60 days after transplanting and at harvest stage. Three plants were uprooted from boarder effect row of each plot and weight of them just after cleaning soil from roots by the help of electric balance for fresh weight and Weed dry weight of weed was recorded by weight was divided by 3 to find the weight of per plant in grams and after sun drying, samples were dried in an oven at 70±1°C for 48 hrs or till constant and weighed. it was expressed in g m<sup>-2</sup>.

Weed Control Efficiency is a percentage that measures how much a weed control treatment reduces the number of weeds. It's calculated by comparing the number of weeds in a treated plot to the amount in an untreated plot.

WCE =

$$\frac{\text{Weed dry weight of control plot} - \text{Weed dry weight of treated plot}}{\text{Weed dry weight of control plot}} \times 100$$

The collected data on various aspects of weeds during two years of experimentation Data obtained for statistical analysis of variance difference among mean of different treatments. The treatments means were compared using the Least Significant Differences test at 5% level of probability by using the Split-split Plot Design procedure as given by Fisher and Yates (1949). The data on population of individual weed species and their dry matter and spike deformities were analyzed after square root transformation ( $X = \sqrt{x+0.5}$ ). The treatment comparisons were made at 5% level of significance.

### 3. RESULTS AND DISCUSSION

#### 3.1 Effect on Weed Population (m<sup>-2</sup>)

The effect of micronutrients and herbicide application on weed density, weed fresh and dry weight (species wise and total) and weed control efficiency in different rice variety. The major weeds *Echinochloa colona* L., *Echinochloa crusgally* L., *Cyperus rotundus* L., *Commenlina bengalensis* L., *Leptochloa chinensis* L., (m<sup>-2</sup>) and other weeds (*Phyllanthus niruri* L., *Eleusin indica* L., *Digitairia ciliaris* L., *Cynodon dactylon* and *Sorghum helipense* L., etc) were found in rice crop.

##### 3.1.1 Effect of rice variety on weed population (m<sup>-2</sup>)

The population of total weeds (m<sup>-2</sup>) depicted in (Figs. 1, 2 and 3) and texted on pooled basis from both the year 2022 and 2023. The minimum population of total weeds (7.52 (57.02), 7.15 (51.64) and 6.07 (37.42) m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) were recorded in Arize 6444 gold variety and maximum population of total weeds (7.87 (62.50), 7.49 (56.59) and 6.36 (40.96) m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) were recorded in PB 1509 variety of rice. This reduction may be attributed to the competitive growth characteristics of Arize 6444 Gold, which effectively suppresses weed establishment and growth (Joshi et al. 2019).

##### 3.1.2 Effect of herbicidal treatments on weed population (m<sup>-2</sup>)

Among both the herbicidal treatments the application of Triafamone 20% + Ethoxysulfuron 10 WG @ 225g ha<sup>-1</sup> at 25 DAT were recorded minimum weed population (6.08 (37.49), 5.50 (30.75) and 4.39 (19.76) m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) which was followed by Bispyribac-sodium10% SC@ 20 g a.i. ha<sup>-1</sup>. The maximum weed population (10.17 (103.91), 10.37 (107.93) and 9.41 (88.95) m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) were recorded with control plot. T Triafamone 20% + Ethoxysulfuron 10 WG @ 225g ha<sup>-1</sup> is an effective herbicide combination that significantly reduced the fresh weight of broadleaf weeds and sedges in rice crops (Basu et al. 2023).

##### 3.1.3 Effect of micronutrients on weed population (m<sup>-2</sup>)

Significantly lowest weed population (7.58 (58.02), 7.21 (52.52) and 6.13 (38.05) m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) were recorded with the foliar spray of Zinc Sulphate 0.5% + Ferrous Sulphate 1% at 45 and 60 DAT which was

followed by Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% at 45 and 60 DAT. However, the maximum weed population (7.91 (63.10), 7.53 (57.13) and 6.39 (41.36) m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) were recorded with foliar spray of Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% + Borax 0.2% at 45 and 60 DAT. Lower weed density in hybrid rice with micronutrient application occurs because micronutrients enhanced the crop's growth, vigour, and competitiveness, allowing rice to better outcompete weeds for resources like light, water and nutrients (Ghosh et al. 2017).

#### 3.2 Effect on Fresh Weight Weeds (g m<sup>-2</sup>)

##### 3.2.1 Effect of rice variety on fresh weight of weeds (g m<sup>-2</sup>)


The fresh weight of weeds (g m<sup>-2</sup>) represented through graph in Figs. (1, 2 and 3) and texted on pooled basis from both the year 2022 and 2023. The minimum fresh weight of weeds (8.33 (69.96), 8.29 (69.26) and 6.66 (44.90) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) was recorded in Arize 6444 gold variety. While, maximum fresh weight of total weeds (8.69 (75.95), 8.68 (75.94) and 6.98 (49.18) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) was recorded in PB 1509 variety of rice (Arthanari 2023).

##### 3.2.2 Effect of herbicidal application on fresh weight of weeds (g m<sup>-2</sup>)


Among both the herbicidal treatments the application of Triafamone 20% + Ethoxysulfuron 10 WG @ 225 g ha<sup>-1</sup> at 25 DAT recorded minimum fresh weight of weeds (5.93 (35.71), 5.91 (35.43) and 4.51 (20.81) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) which was followed by Bispyribac-sodium10% SC@ 20 g a.i. ha<sup>-1</sup> (6.48 (42.44), 6.44 (41.91) and 5.27 (28.24) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively). On the contrary, the maximum fresh weight of weeds (13.12 (172.63), 13.12 (172.68) and 10.69 (114.69) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) was recorded with control plot (Pouramir et al. 2020).

##### 3.2.3 Effect of micronutrients on fresh weight of weeds (g m<sup>-2</sup>)


Significantly lowest fresh weight of weeds (8.40 (71.04), 8.37 (70.62) and 6.73 (45.77) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) was recorded with the foliar spray of Zinc Sulphate 0.5% + Ferrous Sulphate 1% at 45 and 60 DAT. While,



**Fig. 1. Effect of herbicidal treatments on varieties**



**Fig. 2. Effect of herbicidal treatments on weed management**



**Fig. 3. Effect of micronutrients on weed management**

the maximum fresh weight of total weeds (8.71 (76.35), 8.70 (76.15) and 6.99 (49.33) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) was recorded with foliar spray of Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% + Borax 0.2% at 45 and 60 DAT (Mohapatra et al. 2021).

### 3.3 Effect on Dry Weight of Weeds (g m<sup>-2</sup>)

#### 3.3.1 Effect of rice variety on dry weight of weeds (g m<sup>-2</sup>)

The dry weight of total weeds (g m<sup>-2</sup>) demonstrated in Figs. (1, 2 and 3) and texted on pooled basis from both the year 2022 and 2023. The minimum dry weight of weeds (4.60 (21.63), 4.63 (21.97) and 4.18 (17.99) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) was recorded in Arize 6444 gold variety. However, the maximum dry weight of weeds (4.79 (23.40), 4.85 (23.98) and 4.37 (19.61) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) was recorded in PB 1509 variety of rice (Dehghannezhad et al. 2023).

#### 3.3.2 Effect of herbicidal application on dry weight of weeds (g m<sup>-2</sup>)

Among both the herbicidal treatments the application of Triafamone 20% + Ethoxysulfuron 10 WG @ 225g ha<sup>-1</sup> at 25 DAT was recorded minimum dry weight of weeds (3.32 (11.52), 3.35 (11.70) and 2.88 (8.80) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) which was followed by Bispyribac-sodium 10% SC @ 20 g a.i. ha<sup>-1</sup> (3.60 (13.49), 3.63 (13.66) and 3.33 (11.62) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively). Contrary, the maximum dry weight of weeds (7.15 (51.61), 7.24 (52.98) and 6.62 (44.30) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) was recorded with control plot (Joshi et al. 2019).

#### 3.3.3 Effect of micronutrients on dry weight weeds (g m<sup>-2</sup>)

The effect of adding more micronutrients, such as borax and silicon, showed a reduction in dry weed weights across the intervals. Significantly lowest dry weight of weeds (4.63 (21.96), 4.68 (22.37) and 4.22 (18.33) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) was recorded with the foliar spray of Zinc Sulphate 0.5% + Ferrous Sulphate 1% at 45 and 60 DAT which was followed by Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% at 45 and 60 DAT (4.63 (22.01), 4.69 (22.45) and 4.23 (18.38) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively). While, the maximum dry weight of weeds (4.80 (23.51),

4.85 (24.05) and 4.38 (19.68) g m<sup>-2</sup> at 45, 60 and 90 DAT, respectively) was recorded with foliar spray of Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% + Borax 0.2% at 45 and 60 DAT (Pouramir et al. 2020).

### 3.4 Effect on Weed Control Efficiency (WCE)

#### 3.4.1 Effect of rice variety on weed control efficiency (WCE)

The data on weed control efficiency are represented through graph in Figs. (1, 2 and 3) and texted on pooled basis from both the year 2022 and 2023 experimentation. The maximum weed control efficiency (54.98, 55.20 and 54.54% at 45, 60 and 90 DAT, respectively) was recorded in Arize 6444 gold variety however, the minimum weed control efficiency (53.62, 53.89 and 53.15% at 45, 60 and 90 DAT, respectively) was recorded in PB 1509 variety of rice. Weed Control Efficiency (WCE) was higher in the variety Arize 6444 gold, across all the growth stage compared to PB 1509 (Ghosh et al. 2017).

#### 3.4.2 Effect of herbicides on weed control efficiency (WCE)

Significantly better weed control efficiency were recorded with Triafamone + Ethoxysulfuron 10 WG @ 225g ha<sup>-1</sup> at 25 DAT (83.86, 84.71 and 83.86% at 45, 60 and 90 DAT, respectively). While, the minimum weed control efficiency (79.08, 78.93 and 77.67% at 45, 60 and 90 DAT, respectively) was recorded with the application of Bispyribac-sodium 10% SC @ 20 g a.i. ha<sup>-1</sup> at 25 DAT (Basu et al. 2023).

#### 3.4.3 Effect of micronutrients on weed control efficiency (WCE)

The highest WCE was observed with the spray of Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% + Borax 0.2% at 45 and 60 DAT at different growth (54.70% at 45 DAT, 54.91% at 60 DAT, and 54.23% at 90 DAT). While, the lowest WCE was noted with the foliar spray of Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% + Borax 0.2% + Silicon 0.1% at 45 and 60 DAT, but differences among treatments were not statistically significant as the value shows no significance (NS) at different growth stage.

## 4. CONCLUSION

Therefore, assess the "Effect of herbicides and micronutrients spray on different varieties of rice (*Oryza sativa* L.)" on weeds flora, weed fresh



weight, dry weight and weed control efficiency. There are three major conclusions can be drawn from research. Firstly, the effect of variety Arize 6444 Gold had better weed suppression ability compared to PB 1509. Secondly, weed management treatment with Triafamone 20% + Ethoxysulfuron 10 WG @ 225g ha<sup>-1</sup> spray at 25 DAT, provided the most effective control of weed flora, weed fresh weight, dry weight and weed control efficiency, across all the stages of growth. Thirdly, among micronutrient treatments, the application of Zinc Sulphate 0.5% + Ferrous Sulphate 1% + Manganese Sulphate 0.1% + Borax 0.2% + Silicon 0.1% foliar spray at 45 and 60 DAT had a notable impact on reducing dry weed weight and significantly influenced on weed control efficiency.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- DAC & FW, (2022). Directorate of Economics and Statistics, Department of Agriculture and Farmers Welfare, Ministry of Agriculture and Farmers Welfare, Government of India.
- Arthanari, P. M. (2023). Weed management with Triafamone herbicide in transplanted rice ecosystem. *Emirates Journal of Food and Agriculture*, 4, 351-356.
- Basu, B. J., Swathi, P., Rao, N. S., and Naik, V. S. (2023). Efficacy of Triafamone 18. 52% SC on weed control and yield in direct sown rice. *International Journal of Environment and Climate Change*, 13(10), 4414-4422.

- Dehghannezhad, H., Zaefarian, F., Abbasi, R., and Otaghsara, N.M. (2023). weed management methods in direct-seeded rice (*Oryza sativa* L.). *Journal of Crop Production*, 16(4), 21-40.
- Ghosh, D., Singh, U. P., Brahmachari, K., Singh, N. K., and Das, A. (2017). An integrated approach to weed management practices in direct-seeded rice under zero-tilled rice-wheat cropping system. *International Journal of Pest Management*, 63(1), 37-46.
- Joshi, N., Pandey, S. T., Singh, V. P., Kumar, A., and Gautam, P. (2019). Weed management in direct-seeded rice (*Oryza sativa*) + brahmi (*Bacopa monnieri*) intercropping system. *Indian Journal of Agronomy*, 64(1), 23-27.
- Mohapatra, S., Tripathy, S. K., Tripathy, S. and Mohanty, A. K. (2021). Effect of sequential application of herbicides on productivity and profitability of transplanted rice. 129-134.
- Pouramir, F., Yaghoubi, B., and Aminpanah, H. (2020). Efficacy of new herbicides triafamone+ ethoxysulfuron, flucetosulfuron and pyrazosulfuron-ethyl on paddy fields weed control. *Iranian Journal of Field Crop Science*, 50(4), 127-136.
- Resurreccion, A. P. and Juliano, B. O. (1981). Properties of poorly digestible fraction of protein bodies of cooked milled rice. *Plant Foods for Human Nutrition*, 31, 119-128.
- USDA (2021). United States Department of Agriculture, National Agricultural Statistics Service. Farm Labor. Washington, D.C.
- Yoshihashi, T., Huong, N. T. T., and Inatomi, H. (2002). Precursors of 2-acetyl-1-pyrroline, a potent flavor compound of an aromatic rice variety. *Journal of Agricultural and Food Chemistry*, 50(7), 2001-2004.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/125907>