



## **Effects of Different Rates of Plant Nutrients on Yield Attributes and Yield of Maize (*Zea mays* L.)**

**Harender<sup>1\*</sup>, Samar Singh<sup>1</sup>, Narender Singh<sup>1</sup>, Kavinder<sup>1</sup>, Manjeet<sup>1</sup> and Naveen Rathi<sup>1</sup>**

<sup>1</sup>*Department of Agronomy, CCS Haryana Agricultural University, Hisar-125004, India.*

### **Authors' contributions**

*This work was carried out in collaboration between all authors. Author Harender designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors SS, NS, Kavinder, Manjeet and NR managed the analyses of the study. Author Kavinder managed the literature searches. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/CJAST/2017/37315

*Editor(s):*

(1) Ming-Chih Shih, Professor, Department of Health and Nutrition Science, Chinese Culture University, Taiwan.

*Reviewers:*

(1) Khalid A. Khalid, National Research Centre, Egypt.

(2) Temegne Nono Carine, University of Yaounde I, Cameroon and The University of Bamenda, Cameroon.

Complete Peer review History: <http://www.sciencedomain.org/review-history/21936>

**Original Research Article**

**Received 10<sup>th</sup> October 2017**  
**Accepted 14<sup>th</sup> November 2017**  
**Published 16<sup>th</sup> November 2017**

### **ABSTRACT**

**Aim:** Experiment was conducted to study the effect of different rates of plant nutrients on yield attributes and yield of Maize (*Zea mays* L.).

**Study Design:** A field experiment in randomized block design consists of 12 treatments combination with three replications.

**Place and Duration of Study:** Regional Research Station, Karnal of CCS Haryana Agricultural University during *kharif* seasons of year 2015.

**Methodology:** 12 treatments combination viz., T<sub>1</sub> - Control (no fertilizer), T<sub>2</sub> - N (150 kg/ha), T<sub>3</sub> - NP (150, 60 kg/ha), T<sub>4</sub> - NPK (150, 60, 60 kg/ha), T<sub>5</sub> - NPK + S (160, 60, 60, 40 kg/ha), T<sub>6</sub> - NPK + Zn (150, 60, 60, 25 kg/ha), T<sub>7</sub> - NPK (150, 60, 60 kg/ha) + Fe (foliar application of FeSO<sub>4</sub> @ 1% twice i.e. 30 and 45 DAS), T<sub>8</sub> - NPK (150, 60, 60 kg/ha) + Mn (foliar application of MnSO<sub>4</sub> @ 0.5% twice i.e. 30 and 45 days after sowing (DAS)), T<sub>9</sub> - NPK + S + Zn (150, 60, 60, 40, 25 kg/ha), T<sub>10</sub> - NPK + S (150, 60, 60, 40, 25 kg/ha) + Zn + Fe (foliar application of MnSO<sub>4</sub> @ 0.5% twice i.e. 30 and 45 DAS), T<sub>11</sub> - NPK + S + Zn (150, 60, 60, 40, 25 kg/ha) + Fe + Mn (foliar application of FeSO<sub>4</sub> @ 1% and MnSO<sub>4</sub> @ 0.5% twice i.e. 30 and 45 DAS) and T<sub>12</sub> - soil test based fertilizer application (150, 60, 40 kg/ha) laid out in randomized block design.

\*Corresponding author: E-mail: rickydagar38@gmail.com;

**Results:** The results revealed that application of recommended NPK with micronutrients (Fe, Zn, Mn) is statically at par to alone NPK application in terms of yield and yield attributes. Maximum grain yield (73400 kg/ha), cob length (15.7cm), cob girth (3.6 cm) and test weight (21.0g) was recorded in treatment T<sub>5</sub> which is significantly superior over T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub> where at least single primary macro nutrient lacking.

**Conclusion:** Finding suggests that use of recommended NPK in combination with Sulphur increase yield attributes and productivity.

*Keywords: Micronutrients; foliar spray; productivity; maize; NPK.*

## 1. INTRODUCTION

Globally, maize is referred as 'Miracle crop' or 'Queen of the Cereals' due to its high productivity potential compared to other family members of *Poaceae* [1]. Maize is a dual-purpose crop used as grain for human consumption and stover solely fed to the livestock. It also serves as a basic raw material to thousands of industries viz., starch, oil, protein, alcoholic beverages, food sweeteners, pharmaceutical, cosmetic, film, textile, gum, package, paper industries etc. [2]. Maize was grown in an area of 12 thousand ha in Haryana, with production of 27 thousand tonnes and productivity of 2.25 tonnes/ha during the year 2016 [3]. Haryana state has an ample scope to increase its acreage and productivity. Strong market demand and resilience of maize to abiotic and biotic stresses have increased the area and production of maize in the country over the past decade. Productivity of maize, however, has not increased proportionately and significant yield gaps are evident across maize growing areas in the country. Adaptation of 4R principle-based site-specific nutrient management decision support tools provides the opportunity for large-scale adoption of improved nutrient management across maize ecologies [4].

Nutrient removal is far excess of their replenishment under intensively cropped cereal systems in India, which has led to wide spread multi-nutrient deficiencies in soils. As a result of improved agronomic, breeding, and biotechnological advancements in maize systems, yields have reached at far higher levels than achieved ever before. However, greater yields of maize have always been accompanied by a significant removal of macro and micro nutrient from the soil. While managing plant nutrients in maize systems, nitrogen (N), phosphorus (P), and potassium (K) remain the major ones for increased productivity. However, cultivation of high yielding maize systems will likely exacerbate the problem of secondary and micronutrient deficiencies, not only because

larger amounts are removed, but also because the application of large amounts of N, P, and K to achieve higher yield targets often stimulates the deficiency of secondary and micronutrients. Information on crop yield response to fertilizer application, agronomic efficiency and return on investment (ROI) to fertilizer application is also essential for determining optimum dose of nutrients. Soils of the major maize growing areas in India are inherently low in soil organic matter and nitrogen. Nitrogen is the major limiting plant nutrient routinely supplemented through application of fertilizers. Through the yield increase in maize due to N fertilization was substantial (92%), the average agronomic efficiency of N in maize, indicated low N use efficiency [4]. [5] reported variable maize yield response to N fertilizer application, ranging from 4000-5160 kg per ha with an average response of 2154 kg per ha.

Phosphorous response is highly variable and is influenced by soil characteristics and growing environment of the crop. Phosphorus application rate, therefore, must be based on expected response of a particular location. Phosphorus application based on yield response alone does not take into account the nutrient removal by crops where response is low or negligible. Finally, management of phosphorus fertilizer for maize systems must take account of residue and organic amendments applied to the soil [6]. The aim of the study was to evaluate the effect of different nutrients on yield attributes and productivity of maize in Haryana Locality.

## 2. MATERIALS AND METHODS

A field experiment in randomized block design consists of 12 treatments combination with three replications was conducted at the Regional Research Station, Karnal of CCS Haryana Agricultural University during *kharif* seasons of year 2015. The treatments were T<sub>1</sub> - T<sub>12</sub> i.e. T<sub>1</sub> - Control (no fertilizer), T<sub>2</sub> - N (150 kg/ha), T<sub>3</sub> - NP (150, 60 kg/ha), T<sub>4</sub> - NPK (150, 60, 60 kg/ha),

T<sub>5</sub> - NPK + S (160, 60, 60, 40 kg/ha), T<sub>6</sub> - NPK + Zn (150, 60, 60, 25 kg/ha), T<sub>7</sub> - NPK (150, 60, 60 kg/ha) + Fe (foliar application of FeSO<sub>4</sub> @ 1% twice i.e. 30 and 45 DAS), T<sub>8</sub> - NPK (150, 60, 60 kg/ha) + Mn (foliar application of MnSO<sub>4</sub> @ 0.5% twice i.e. 30 and 45 DAS), T<sub>9</sub> - NPK + S + Zn (150, 60, 60, 40, 25 kg/ha), T<sub>10</sub> - NPK + S (150, 60, 60, 40, 25 kg/ha) + Zn + Fe (foliar application of MnSO<sub>4</sub> @ 0.5% twice i.e. 30 and 45 DAS), T<sub>11</sub> - NPK + S + Zn (150, 60, 60, 40, 25 kg/ha) + Fe + Mn (foliar application of FeSO<sub>4</sub> @ 1% and MnSO<sub>4</sub> @ 0.5% twice i.e. 30 and 45 DAS) and T<sub>12</sub> - soil test based fertilizer application (150, 60, 40 kg/ha). The experimental site was located at latitude of 29° 43' 42.19" N longitude of 76° 58' 49.88" E and at an altitude of 253 m above mean sea level. The soil of experimental field was deep with silty clay loam in texture, slightly alkaline pH (8.2), medium in organic carbon (0.46%), available P<sub>2</sub>O<sub>5</sub> (15 kg/ha), K<sub>2</sub>O (127 kg/ha) and low in available N (120 kg/ha). The experimental site had been used over the years for continuous maize cropping. Maize crop was in alternation with wheat crop grown in *spring* season.

In experiment gross plot size was 4.2 m x 5.0 m with net plot size 2.8 m x 5.0 m. Maize variety HPQM 1 available from Regional Research Station, Karnal was sown on flat bed at the spacing of 70 cm x 20 cm with seed rate of 20 kg/ha. HQPM 1 is hybrid variety of maize which is cross of HKI 193-1x HKI 163. Pre-sowing irrigation was applied to the field to facilitate preparatory tillage and seed germination. The seed bed was prepared by four harrowing followed by cultivator twice and planking. Furrows were opened in dry condition to facilitate the dibbling of maize. 1/4<sup>th</sup> dose of nitrogen (37.5 kg/ha), full dose of phosphorus (60 P<sub>2</sub>O<sub>5</sub> kg/ha) and full dose of potash (60 K<sub>2</sub>O kg/ha) through urea, DAP and MOP respectively, were applied as a basal dose at the time of sowing and remaining 3/4<sup>th</sup> dose of N (112.5 kg/ha) was top dressed through urea in 3 equal splits i.e. knee-high stage, tasseling stage and dough stage. Maize hybrid as per treatment was sown by dibbling method on dry ridges opened at 70 cm with plant to plant spacing of 20 cm immediately followed by irrigation up to half of the ridge to ensure proper soil moisture for better germination of seed. Crop received very good rainfall during the crop growth period. Recommended package of practices was followed for all other operations. The length of ten randomly selected cobs from each net plot was measured from base to tip and average was

recorded as cob length. Cob girth was measured for 5 cobs from each net plot and average of 5 cobs was recorded as cob girth. Numbers of rows/cob were counted for 5 cobs from each net plot and average of 5 cobs was calculated. Number of kernels per cob was calculated by multiplying the number of grains/row by the number of rows/cob. All the cobs were harvested from net plot area and converted on hectare basis. The weight of 100 seeds drawn at random from the grains of 10 cobs from each plot was recorded. Five cobs were selected from each plot and after sun drying to 15% moisture, the grains were separated from cobs and weight of grains was measured and converted on hectare basis. Straw yield was recorded after remaining the cobs at harvest from net plots after sun drying to 15% moisture and expressed in kg/ha. The shelling was calculated by the given formula on sun dry weight basis:

$$\text{Shelling (\%)} = \frac{\text{Grain yield}}{\text{Cob yield without husk}} \times 100$$

The data was analyzed using analysis of variance (ANOVA) as applicable to randomized complete block design. The significance of the treatment effects was determined using F-test at 5% significance level.

### 3. RESULTS AND DISCUSSION

#### 3.1 Yield and Yield Attributes

##### 3.1.1 Cob length and Cob girth

The perusal of data (Table 1) indicates that cob length was significantly higher in all nutrient treatments compared to control. Highest cob length (15.7 cm) was recorded in treatment T<sub>5</sub> followed by T<sub>6</sub> (15.5 cm), T<sub>9</sub> (15.5 cm) and lowest in T<sub>1</sub> (10.5 cm). [7] and [8] also reported that highest cob length (19.0 cm) was obtained from the treatment with 250-76-88-7.4 kg N-P-K-Zn/ha. Results also indicate that cob girth differed significantly in different nutrient treatments. Treatment T<sub>5</sub> recorded highest cob girth (3.6 cm) followed by T<sub>6</sub> (3.5 cm) and T<sub>9</sub> (3.5 cm). Significantly lower cob girth compared to other nutrient treatments were recorded in control (2.6 cm) followed by N (2.8 cm) and NP (3.0 cm) treatments. The result obtained during the investigation were similar with the finding of [7] in reference of cob girth who obtained maximum cob girth (3.2 cm) from the treatment with 250-76-88-7.4 kg N-P-K-Zn/ha.

**Table 1. Effect of different rate of nutrient on yield attributing characteristics of maize**

Treatments	Cob length (cm)	Cob girth (cm)	Number of rows/cob	No. of kernels /cob	No. of cobs /ha(000/ha)
T <sub>1</sub>	10.5	2.6	12.7	373.3	67.9
T <sub>2</sub>	12.4	2.8	12.7	422.7	67.6
T <sub>3</sub>	13.5	3.0	12.8	441.3	68.7
T <sub>4</sub>	14.5	3.4	14.8	448.0	68.1
T <sub>5</sub>	15.7	3.6	15.7	527.3	69.8
T <sub>6</sub>	15.5	3.5	15.8	474.0	69.4
T <sub>7</sub>	14.9	3.4	15.3	504.7	68.9
T <sub>8</sub>	14.6	3.4	14.5	504.7	68.7
T <sub>9</sub>	15.5	3.5	15.5	497.3	69.0
T <sub>10</sub>	15.1	3.4	15.4	491.3	68.6
T <sub>11</sub>	15.0	3.4	14.7	505.3	68.7
T <sub>12</sub>	15.1	3.4	15.5	504.7	68.6
SE(m)±	0.58	0.65	2.0	40.6	0.67
CD (P=0.05)	1.71	0.25	NS	NS	NS

### **3.1.2 Number of rows/cob, kernels/cob, cobs/ha**

The no. of rows/cob, no. of kernel/cob and cobs/ha did not affect by different rates of nutrients (Table 1). [9] and [10] also reported highest number of rows/cob in treatment of recommended dose of fertilizer + 1 spray of multi-nutrients whereas lowest no. of rows/cob observed in control. Application of different nutrients in maize produced significantly higher no. of kernels/cob compared to control treatment. Treatment T<sub>5</sub> recorded highest no. of kernels/cob (527.3) followed by T<sub>11</sub> (505.3), T<sub>8</sub> (504.7) and T<sub>12</sub> (504.7). Similar results were also reported by [11] and [12].

Treatment, T<sub>5</sub> recorded highest no. of cobs/ha (69800) followed by T<sub>6</sub> (69400) and T<sub>9</sub> (69000). [13] and [14] also reported similar results, with highest number of cobs/plant from the treatment of 25% RDF+ biofertilizers (*Azotobacter*+PSB) + green manuring with sunhemp+compost.

### **3.1.3 Test weight**

The data (Table 2) indicated that all the treatments except treatment T<sub>2</sub> produced significantly bolder kernels as compared to control. Treatment T<sub>5</sub> recorded highest test weight (21.0) which was statistically at par with all treatment except T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub> where at least one primary macro nutrient lacking. [9] and [15] also found maximum test weight when maize plants received basal dose of conventional fertilizer with single spray of multi-nutrients solution.

### **3.1.4 Grain yield and straw yield (q/ha)**

Among all treatments, highest grain yield (73.4 q/ha) was recorded in treatment T<sub>5</sub> followed by T<sub>9</sub> (73.2 q/ha), T<sub>6</sub> (72.8 q/ha), T<sub>11</sub> (72.2 q/ha) and T<sub>7</sub> (72.0 q/ha) as shown in (Table 2). Treatments where at least one primary macro nutrient lacking (T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub>) produce significantly lower grain and straw yield compare to rest of the other treatments. The results were confirmed with the finding of [4] and [8] as they found maximum grain yield by the use of farmyard manure and in combination with inorganic fertilizers.

The application of NPK and NPK with micronutrients produced similar straw yield and produced significantly superior to the treatment NP, N and control. Treatment of NPK over NP (107.8 q/ha), NP over N (84.7 q/ha) and N over control (76.8 q/ha) produced significantly higher straw yield. Among the treatments significantly higher straw yield/ha was recorded in T<sub>5</sub> (110.0 q/ha) followed by T<sub>9</sub> (109.8 q/ha) and T<sub>6</sub> (109.1 q/ha). The highest biological yield was also influenced which might be attributed to the additional availability of nutrients [16,8] and [17]. Treatment T<sub>5</sub> produce significantly higher grain and straw yield over T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub>. Treatment T<sub>5</sub> produce 80%, 40% and 30% higher grain and 79%, 43% and 30% higher straw yield over T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub>.

### **3.1.5 Shelling (%)**

Recovery of grains from cob was not affected by different rates of plant nutrients (Table 2). Highest recovery of grains from cob (80.2) was recorded in treatment T<sub>5</sub>. Lowest were recovered in treatment

**Table 2. Effect of different nutrient treatments on test weight, grain yield, straw yield/ha and shelling percentage of maize**

Treatments	Test weight (g) (100 grains)	Grain yield (q/ha)	Straw yield (q/ha)	Shelling (%)
T <sub>1</sub>	12.8	40.9	61.3	73.0
T <sub>2</sub>	14.2	51.2	76.8	75.6
T <sub>3</sub>	18.4	56.5	84.7	76.0
T <sub>4</sub>	20.4	71.8	107.8	79.0
T <sub>5</sub>	21.0	73.4	109.8	80.2
T <sub>6</sub>	19.8	72.8	109.1	79.2
T <sub>7</sub>	19.4	72.0	108.0	79.0
T <sub>8</sub>	19.8	71.3	107.0	78.9
T <sub>9</sub>	19.3	73.2	110.0	79.2
T <sub>10</sub>	19.3	71.0	106.6	79.1
T <sub>11</sub>	19.8	72.2	108.3	79.1
T <sub>12</sub>	20.0	71.4	107.1	79.0
SE(m)±	0.59	1.78	2.24	1.85
CD (P=0.05)	1.75	5.25	6.63	NS

T<sub>1</sub> (73.0). Similar, results were reported by [10] and [18] in respect to shelling per cent, as they also found significantly higher shelling percentage with P<sub>50</sub>-PEC+P<sub>50</sub>-F.

#### 4. CONCLUSIONS

Application of NPK with S resulted in highest yield attributes and yield of maize crop. No. of rows/cob, no. of kernel/cob, no. of cobs/ha and shelling percentage of maize were not affected by different rate of nutrient applied. So, it is concluded from the study that use of recommended NPK in combination with S improves/increases yield attributes and productivity of maize.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Choudhary AK, Choudhary LB. Genetic studies in some crosses of maize (*Zea mays* L.). Journal Research (BAU). 2002; 14(1):87-90.
- Kumar B, Chikkappa GK, Jat SL, Parihar CM, Yathish KR, Singh V, Hooda KS, Dass AK, Mukri G, Sekhar JC, Kumar R, Kumar RS. Maize biology: An introduction, directorate of maize research, Indian Council of Agricultural Research (ICAR). Technical Bulletin. 2012/2:32.
- Anonymous. Annual Kharif-maize progress report. Directorate of Maize Research, IARI, New Delhi; 2016.
- Jat SL, Parihar CM, Singh AK, Jat ML, Sinha AK, Mishra BN, Meena H, Paradkar VK, Singh CS, Singh D, Singh RN. Integrated nutrient management in quality protein maize planted in rotation with wheat: Effect on productivity and nutrient use efficiency under different agro-ecological conditions. Indian Journal of Agricultural Sciences. 2013;83(4):391-396.
- Satyanarayana T, Jat ML, Kaushik M, Parihar CM, Jat SL, Tetarwal JP, Jat RK, Saharawat YS. Fertiliser best management practices for maize systems. Indian Journal of Fertilizers. 2013;9:80-94.
- Walia SS, Kler DS. Effect of organic and inorganic sources of nutrition on growth, macro and micro nutrient uptake in maize under maize wheat sequence. Indian Journal of Ecology. 2010;37:27-29.
- Paramsivan M, Malarvizhi P, Thiyageswari S. Balanced use of inorganic fertilizers on maize yield, nutrient uptake and soil fertility in alfisols. Karnataka Journal of Agricultural Sciences. 2012;25(4):423-426.
- Balai ML, Verma A, Nepalia V, Kanthaliya PC. Productivity and quality of maize as influenced by integrated nutrient management under continuous cropping and fertilization. Indian Journal of Agricultural Sciences. 2011;81(4):374-376.
- Ghaffari A, Ali A, Tahir M, Waseem M, Ayuv M, Iqbal A, Mohsin AU. Influence of integrated nutrients on growth, yield and

- quality of maize. American Journal of Plant Sciences. 2011;2:63-69.
10. Paramesh V, Dhar S, Vyas AS, Dass A. Studies on impact phosphoenriched compost, chemical fertilizer and method of zinc application on yield, uptake and quality of maize. Indian Journal Agricultural Sciences.2014;59(4):613-618.
  11. Iqbal S, Khan HZ, Ehsanullah AN, Zamir MSI, Javeed HMR. Nitrogen management studies in maize hybrids. Cercetari Agronomice in Moldoba. 2013;46(3):39-48.
  12. Maqsood M, Abid AA, Iqbal A, Hussain MI. Effect of variable rates of nitrogen and phosphorus on growth and yield of maize. Journal of Biological Sciences. 2001;1(1): 19-20.
  13. Kalhapure AH, Shete BT, Dhonde MB. Integrated nutrient management in maize for increasing production with sustainability. International Journal of Agriculture, Food Science and Technology. 2013;4(3):195-206.
  14. Sujatha MG, Lingaraju BS, Palled YB, Ashalatha KV. Importance of integrated nutrient management practices in maize under rainfed condition. Karnataka Journal of Agricultural Sciences. 2008;21(3):334-338.
  15. Seepat S, Rai RK. Effect of phosphorus levels and sources on productivity, nutrient uptake and soil fertility of maize-wheat cropping system. Indian Journal of Agronomy. 2013;58(3):292-297.
  16. Ali K, Khalil KS, Munsif F, Rab A, Nawab K, Khan AZ, Kamal A, Khan ZH. Response of maize to various nitrogen sources and tillage practices. Sarhad Journal of Agriculture. 2012;28(1):9-14.
  17. Dibaba DH, Hunshal CS, Hiremath SM, Awaknavar JS, Wali MC, Nadagouda BT, Chandrashekar CP. Growth and yield of maize hybrids has influenced by application of NPK and S levels. Karnataka Journal of Agricultural Sciences. 2014; 27(4):454-459.
  18. Banerjee M, Bhuiya GS, Malik GC, Maiti D, Dutta S. Precision nutrient management through use of LCC and nutrient expert in hybrid maize. Universal Journal of Food and Nutrition science. 2014;2(2): 33-36.

© 2017 Harender et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://sciencedomain.org/review-history/21936>