



Serum Ferritin as an Indicator of Neonatal Anaemia

Asinobi Isaac Nwabueze ^{a,b*}

^a *Department of Paediatrics, College of Medicine, Enugu State University of Science and Technology, Enugu, Nigeria.*

^b *Department of Paediatrics, Enugu State University of Science and Technology Teaching Hospital, Enugu, Nigeria.*

Author's contribution

The sole author designed, analyzed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/IBRR/2023/v14i4312

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100035>

Original Research Article

Received: 02/03/2023

Accepted: 14/05/2023

Published: 15/05/2023

ABSTRACT

Introduction: Haemoglobin levels are usually used to diagnose neonatal anaemia. By the time haemoglobin levels drop, body iron is depleted. Serum ferritin is the standard measure for assessment of iron levels in neonates and detects iron deficiency earlier than haemoglobin levels.

Aims: To determine the prevalence of low haemoglobin and ferritin levels amongst term and preterm newborn babies. To determine the value of serum ferritin in the diagnosis of low iron stores amongst term and preterm newborns.

Methodology: This was a cross sectional descriptive study carried out at the Neonatal Intensive Care Unit of the University of Nigeria Teaching Hospital (UNTH), Enugu, Nigeria between June and December 2014. The study included 140 newborns of all birth weights delivered at the UNTH. These were categorized into preterm (gestational age <37 completed weeks) and term (gestational age ≥37 completed weeks). Babies with C-reactive protein levels > 10mg/dl, who were intra-uterine growth restricted, and whose mothers had conditions associated with low iron stores were excluded from the study. Anthropometric measurements were done for all subjects. Haemoglobin estimation and ferritin assay were carried out and the prevalence of neonatal anaemia was determined using each of these.

*Corresponding author: E-mail: isaacasinobi@esu.edu.ng, isaacasinobi@gmail.com;

Results: The range of haemoglobin concentration in the study population was 12.22g/gl – 22.80g/dl. The mean serum haemoglobin concentrations were 15.69mg/dl \pm 1.58 and 16.45 \pm 1.92 in the preterm and term babies respectively ($t = 2.557$, $P = .0116$). The prevalence of low haemoglobin concentrations amongst both preterm and term babies was zero= .024). The range of serum ferritin level in the study population was 20.6 μ g/l - 296 μ g/l. The mean serum ferritin levels were 63.13 μ g/l \pm 23.93 and 133.67 μ g/l \pm 50.14 in the preterm and term babies respectively ($t = 10.623$, $P < .001$). The prevalence of low serum ferritin in the study population was 22.14%, but was higher in preterm than term babies 35.7% vs 8.6%: (OR – 5.926, 95% C.I OR = 2.248 – 15.619)($P < .001$).

Conclusion: Serum ferritin assay is more useful than haemoglobin as an indicator of anaemia during the neonatal period.

Keywords: Anaemia; ferritin; iron; prevalence; indicator.

1. INTRODUCTION

Anaemia is characterized by a reduction in red cells mass or haemoglobin concentration to levels that are insufficient to meet the metabolic needs of the body [1,2]. It is usually defined as a haemoglobin concentration or haematocrit > 2 standard deviations below the mean for age [2]. Globally, the commonest cause of this reduction is iron deficiency [1] Anaemia has been documented as the commonest haematological disorder diagnosed at birth [2]. Despite this, there is a dearth of literature on this subject [3]. Studies have shown a prevalence of neonatal anaemia of 17.5 – 21% [3]. Risk factors include premature birth, low birth weight, low maternal blood levels, bleeding in pregnancy (placenta previa and abruption placenta and unsafe delivery practices [2]. Anaemia during the neonatal period has been linked to late neurological deficits, and is a leading risk of perinatal mortality [3]. Healthy newborn babies are however often not routinely screened for anaemia, but are usually only screened following hospitalization.

“Accurate determination of iron status is crucial for diagnostic and screening purposes” [4] “Several markers are available to assess total body iron and iron stores” [5]. “Some of these include serum iron, haemoglobin (Hb) concentrations, ferritin concentrations, mean corpuscular volume (MCV), total iron binding capacity (TIBC), transferrin saturation, red cell distribution width, zinc protoporphyrin (ZnPP), and serum transferrin receptor (sTfR)” [5]. “However, each of these has major limitations, some of which include a lack of association with gestational age, and a lack of specificity for iron deficiency” [5]. “There is thus no agreement on the specific laboratory criteria for iron deficiency” [6]. “There is also no consensus concerning

whether to use single or multiple criteria, and on which iron status variables to use” [6]. “The usual requirement is either a low serum ferritin or a combination of multiple criteria, such as, abnormal values for any two of three iron status variables” [7].

In clinical settings however, haemoglobin estimation alone is usually used for decision making [3]. This is as a result of its ease of estimation and interpretation of results [8]. For decades, defining cut-off values for low haemoglobin in infants and children has been a source of debate [9]. In addition, cut-offs for neonates are not included in most guidelines and recommendations [9]. Among both term and preterm babies, haemoglobin concentrations of 15mg/dl have been reported as optimal [10]. However, minimal acceptable levels are much lower than this (12 g/dl for preterm infants and 11 g/dl for full-term neonates) [10]. “That notwithstanding, haemoglobin concentration alone cannot be used to diagnose iron deficiency” [1]. “This is due to the hierarchical loss of tissue iron” [11-13]. “This means that, in the presence of iron deficiency, there is a preference to maintain red blood cell (RBC) iron for the synthesis of haemoglobin, at the expense of brain, heart and skeletal muscle iron stores” [11-13]. Thus, by the time measured serum iron deficiency is obtained, iron stores have already been depleted” [11].

“Ferritin is the major storage form of iron” [4,11]. “In the body, small amounts of ferritin are secreted into the plasma” [14]. “The concentration of this is positively correlated with the size of the total body iron stores in the absence of inflammation” [14]. “Serum ferritin levels are reduced only in iron deficiency, and are the most sensitive biomarker to test for early stages of iron deficiency as well as iron

deficiency anemia". [4,11,15]. "In addition, the body ferritin levels, in contrast to haemoglobin, are not affected by residential elevation above sea level or smoking behavior" [14]. Different cut-offs have been described for defining low ferritin [16-18]. However, levels below 35µg/l have been found to be associated with depletion of brain stores and with consequent effects on brain function [19]. Serum ferritin estimation has been recommended as the standard for the assessment of body iron stores in both term and preterm babies [4,11].

Serum ferritin being an acute phase reactant, can however be increased in the presence of inflammation [11,14]. Thus, its interpretation requires exclusion of other causes of inflammation [20]. The author thus hypothesizes that, amongst preterm and term babies, serum ferritin may be a better indicator of anaemia than haemoglobin. This study thus aims to determine the prevalence of low haemoglobin and ferritin levels amongst term and preterm newborn babies. It also aims to determine the value of serum ferritin in the diagnosis of low iron levels amongst term and preterm newborns.

2. MATERIALS AND METHODS

This cross sectional descriptive study was carried out at the Neonatal Intensive Care Unit of the University of Nigeria Teaching Hospital (UNTH), Enugu, Nigeria between June and December, 2014. The study included 140 neonates. These were categorized as follows: 70 preterm babies (gestational age < 37 completed weeks) and 70 term babies (gestational age ≥ 37 completed weeks). Babies with C-reactive protein levels > 10mg/dl (to exclude other causes of inflammation), who were intra-uterine growth restricted, and whose mothers: had ante partum haemorrhage or other bleeding episodes during pregnancy; had severe anaemia (haemoglobin cut-off point of less than 11g/dl defines maternal anaemia in the later stages of pregnancy [21]); diabetes mellitus or hypertension; and who smoked were excluded from the study. Subjects were enrolled consecutively until the calculated sample size was reached.

Data was collated and analysed using Statistical Package for Social Sciences (SPSS) Version 20. Relationships between continuous variables were determined using correlation and linear regression analysis. Means of continuous variables were compared using Student's t-test, while associations between categorical variables

were determined using chi-square and logistic regression analysis as applicable. All tests were considered significant at $P < .05$.

2.1 Blood Sample Collection

Umbilical venous blood was collected from a double clamped segment of the umbilical cord during delivery. "This was then placed into a small study designated storage box at room temperature designated. Subsequently, the Howard Kelly forceps on one end of the section of the cord was removed. The umbilical vein was identified and depending on its size, a 5,6 or 8 Fr gauge nasogastric tube attached to a 10ml syringe was inserted and at least 6ml of blood was withdrawn. Where this did not work, the blood was obtained by venopuncture of the side of the cord corresponding to the identified umbilical vein. A drop (approximately 0.2 ml) of the blood obtained was first immediately dropped onto a microcuvette which was inserted into the Hemocue® Hb 201⁺ for estimation of haemoglobin concentration. Serum was then obtained from the remaining blood for both CRP and ferritin estimation at the Haematology laboratory of UNTH using the Diagnostic Automation 800 ELISA machine®. Low ferritin was regarded as a measured serum level of less than 35µg/l" [11].

2.2 Sample Size Determination

"The sample size (n) for an infinite population of more than 10,000 was first obtained using the formula for the comparison of proportions" [22]:

$$n = \frac{[P1(1 - P1) + P2(1 - P2)]}{(P1 - P2)^2} \times Cp \text{ power}$$

Where:

P1 = Proportion of preterm babies from a previous study(10%) [23] P2 = Proportion of term babies from a previous study (18%) [23] Cp power = 13 when p value is 0.05 and power is 95%, Therefore:

$$n = \frac{[(0.1)(0.9) + (0.18)(0.82)] \times 13}{(-0.08)^2} = 483$$

"Since this study was done on a finite population (less than 10,000), the sample size for a finite population was then derived using the formula below" [24].

$$nf = \frac{no}{(1+ no/N)}$$

Where:

nf = final (or minimum) sample size
 n_0 = initial sample size (derived above)
 N = population of preterm births over a 12 month period in UNTH i.e. 70.

$$nf = \frac{483}{(1 + 483/70)} = 61$$

An attrition rate of 10% was used in the study to account for possible sample loss. Thus, the total minimum sample size was calculated to be 67 preterm babies, which was rounded off to 70 each.

3. RESULTS

3.1 Study Characteristics

The baseline characteristics of the study population is shown in Table 1. There were 68 males and 72 females, giving a male to female ratio of 0.9:1. Mothers of 84 (60%) babies reside in urban areas while mothers of 56 (40%) babies reside in rural areas. Majority of the subjects (36.4%) were of the upper socioeconomic class.

The gestational age of the study population ranged from 25 weeks to 39 weeks, with birth weight ranging from 0.55kg to 5.2kg. The distribution of other anthropometric parameters

amongst the study population is shown in Tables 2 and 3.

3.2 Haemoglobin Concentrations in the Study Population

The range of haemoglobin concentration in the study population was 12.22g/gl – 22.80g/dl. The mean serum haemoglobin concentrations were 15.69mg/dl \pm 1.58 and 16.45 \pm 1.92 in the preterm and term babies respectively ($t = 2.557$, $P = .0116$). The prevalence of low haemoglobin concentrations amongst both preterm and term babies was zero.

3.3 Ferritin Levels in the Study Population

The range of serum ferritin level in the study population was 20.6 μ g/l - 296 μ g/l. The mean serum ferritin levels were 63.13 μ g/l \pm 23.93 and 133.67 μ g/l \pm 50.14 in the preterm and term babies respectively ($t = 10.623$, $P < .001$). The prevalence of low serum ferritin in the study population was 22.14%, but was higher in preterm than term babies 35.7% vs six 8.6%: (OR – 5.926, 95% C.I OR = 2.248 – 15.619) ($P < .001$). In addition, preterms were found to be six times more likely than term babies to have low serum ferritin levels (OR – 5.926, 95% C.I OR = 2.248 – 15.619) (Table 4).

Table 1. Demographic, maternal and neonatal variables

Characteristics	Preterm n = 70 (50%)	Term n = 70 (50%)	Total n = 140 (100%)
Gender			
Male	34 (24.3)	34 (24.3)	68
Female	36 (25.7)	36 (25.7)	72
	70	70	140
Tribe			
Ibo	68	64	132
Yoruba	1	3	4
Hausa/Fulani	1	3	4
	70	70	140
Socioeconomic Class			
Upper	27	24	51
Middle	15	26	41
Lower	28	20	48
	70	70	140

Table 2. Anthropometric indices of the preterm babies

Gestational age (weeks)	N (70)	Weight (g) Mean (SD)	Length (cm) Mean (SD)	OFC (cm) Mean (SD)	CC (cm) Mean (SD)
<28	7	680 (0.80)	24.29 (4.82)	23.71 (1.07)	21.14 (1.21)
28 - <32	27	1610 (0.46)	39.59 (4.41)	27.49 (9.73)	29.24 (3.29)
32 – 36	36	2288 (0.51)	45.39 (3.62)	33.64 (1.73)	31.46 (3.58)
Total	70	1861 (0.68)	41.06 (7.44)	30.27 (7.11)	29.55 (4.43)

Table 3. Anthropometric indices of the term babies

Gestational age (weeks)	N (70)	Weight (g) Mean (SD)	Length (cm) Mean (SD)	OFC (cm) Mean (SD)	CC (cm) Mean (SD)
37	32	2840 (270)	47.27 (2.26)	34.94 (1.38)	33.30 (1.63)
38	28	3750 (610)	49.84 (2.53)	36.25 (0.91)	35.21 (1.93)
39	10	4210 (700)	51.80 (3.49)	37.20 (0.95)	36.10 (1.35)
Total	70	3400 (0.73)	48.92 (3.03)	35.79 (1.42)	34.49 (2.04)

Table 4. Frequency of low serum ferritin levels in preterm and term subjects

Ferritin levels (µg/l)	Preterm	Term	Significance	OR	95 % C.I for OR
Low n (%)	25 (35.7)	6 (8.6)	$P < .001$	5.926	2.248 – 15.619
Normal n (%)	45 (64.3)	64 (91.4)			

(µg/l = microgram per liter, C.I = confidence interval, OR = odds ratio)

4. DISCUSSION

The mean haemoglobin concentration of newborn babies in this study was similar to the values of 12.54 ± 2.54 g/dl and 13.44 ± 2.23 g/dl ($P = 0.02$) obtained by Adediran et al. [25] in South-West Nigeria amongst term anaemic and non anaemic respectively. However, higher values were recorded by Tiruneh et al. [26] in Ethiopia and Esslami et al. [27] in Iran. This difference could be explained by the slightly higher and much higher sample sizes used in the former and latter studies respectively. It should however be noted that there is paucity of literature on mean haemoglobin levels in the entire population of newborns. Most literature concentrate on subsets of newborns such as low birth weight, anaemic, or categories of preterm neonates [28-30].

Haemoglobin levels were significantly lower in preterms when compared with term babies. This is similar to the findings by Eslami et al. [27] of a mean hemoglobin value of cord blood in preterm neonates of 14.77 g/dl \pm 1.69, and in term neonates of 15.4 g/dl \pm 5.07 SD and SD. ($P=0.036$). It has previously been documented that between 22 and 40 weeks gestation, haemoglobin concentrations increase linearly by approximately 0.21g/dl per week of gestation [N]. This reflects the increasing metabolic needs of the growing foetus with a concurrent increase in cell mass and body size [31].

Using the recommended cut-off of 11g/dl and 13g/dl in preterm and term babies yielded a zero prevalence for foetal anaemia. Several other cut-offs have been recommended for both healthy and sick neonates [1,6,31,32]. The World Health Organization (WHO) recommends 11g/dl as the cut-off to define anaemia in infants and children

[1]. However, these cut-offs were derived from a few western studies and have not been revised since. Also, this value was applied to infants from six months of age and did not include newborn babies [1]. Domellöf et al [6] conducted a study to re-evaluate the diagnostic criteria for iron deficiency amongst infants. A slightly lower cut-off of 10.5g/dl was obtained [6]. This however was also limited to infants 6-9 months of age [6]. Haemoglobin cut-offs to establish anemia amongst neonates remains a subject of debate. Using higher haemoglobin cut-offs, several other studies have documented a prevalence for foetal anaemia ranging from 5.7-65.6% [24,32,33] Adediran et al [25] in South West Nigeria reported a prevalence of 28.9%, while a prevalence of 65.6% was obtained in Abakaliki, South East Nigeria [34]. Values of 26.4 have been reported in other parts of Nigeria [35] An even lower prevalence has been observed in other parts of Africa [24, 34] and in Western countries [36-38]. In contrast to this however, the prevalence of iron deficiency without anaemia amongst toddlers in the US has been estimated at between 6.6% and 15.2% [39].

In our study, using a serum ferritin concentration of <35 µg/L to define low serum ferritin levels, the prevalence of low serum ferritin was 22.14%. Several studies have reported on the prevalence of low ferritin levels amongst newborn babies [9,11,16]. Ferritin measurements and corresponding cut-offs facilitate the monitoring of iron deficiency trends and the assessment of the impact of health and nutrition interventions [14]. However, the lack of normative values for serum ferritin concentration amongst neonates with gestational age between 23 and 41 weeks has led to difficulty in establishing the prevalence of low iron stores amongst this population [9]. "The WHO/CDC in 2005, stated that serum ferritin

values of less than 12 µg/l in children <5 years are indicative of a depletion of iron stores” [14]. “In neonates however, and more especially in preterms, rapid growth and development creates a greater demand for iron, with larger iron stores needed to meet this demand” [11,12,40,41]. “Thus, at serum ferritin levels of <15µg/l, the neonate already has deficient iron stores with a significant risk for neurodevelopmental problems” [19]. “Several authors have proposed different values for the definition of low ferritin in neonates” [4,11,19]. “Sidappa and co-workers [19] in USA, 2004, were able to estimate from existing nomograms that newborn ferritin concentrations less than 35µg/l represent a risk to the developing brain”. “This value was thus used as a cut-off in their neurodevelopmental studies” [42].

The prevalence of low serum ferritin obtained in this study is much lower than the 59.2% obtained by Adediran et al. [43] in an earlier study on term babies in South-West Nigeria. This is despite the use of a higher ferritin cut-off of 60µg/l. The use of term babies alone also dampened the effect as ferritin levels are known to increase with gestational age. A much lower prevalence of 19.8% was obtained by Zhang et al [16] in a cohort of full term normal birth weight infants in China, using similar cut-offs as ours. However, the authors excluded babies with a C-reactive protein (CRP) of > 5 mg/l, compared with 10mg/l used in this current study. It has been widely agreed that, in addition to ferritin, an independent indicator of the acute phase response, such as C-reactive protein (CRP), α-1 antichymotrypsin (ACT), α-1- acid-glycoprotein (AGP) and serum amyloid A, should be measured [14,44-46]. CRP is the most commonly used as it responds quickly to inflammation [19,20,47-49] with values of greater than 10mg/dl being the cut-off signifying infection [18,19,45-47]. Using lower cut-offs than this can result in the inclusion of subjects whose serum ferritin concentrations may be reflective on an inflammatory process, thereby reducing the number of newborns seen as having low serum ferritin, and thus reduce the prevalence of low serum ferritin.

According to the American Academy of Family Physicians, the sensitivity of serum ferritin for the diagnosis of iron depletion is 89% compared with 26% for hemoglobin [15]. In addition, a ferritin cut off of ≤ 30 ng/mL is 92% sensitivity and 98% specificity for the diagnosis of iron deficiency anemia, and is the best screening test for this disorder [15]. As iron deficiency without anaemia

is poorly recognised by clinicians despite its high prevalence, adequate evaluation using appropriate tests will identify early stages of iron deficiency and will potentially result in iron therapy, preventing iron deficiency anemia [15,39].

5. CONCLUSION

The assessment of low iron stores in newborn babies is crucial to their subsequent well-being. There is a wide variation of haemoglobin levels amongst newborn babies. The prevalence of anaemia was zero using haemoglobin levels to determine iron stores, and 22.14% using ferritin concentrations. Using serum ferritin in the diagnosis of low iron stores is thus more beneficial.

CONSENT AND ETHICAL APPROVAL

The study was approved by the University of Nigeria Teaching Hospital Health Research Ethics Committee. Written informed consent was obtained from the parents of the study participants.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. WHO. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Vitamin and Mineral Nutrition Information System. Geneva, World Health Organization (WHO/NMH/NHD/MNM/11.1); 2011. Available:<http://www.who.int/vmnis/indicators/haemoglobin.pdf>, accessed [Nov 30, 2013].
2. Walter AW(ed), Perinatal Anemia. MSDS Manual. Merck & Co, Inc., Rahway, NJ, USA Last review/revision Dec; 2022. Available:<https://www.msdsmanuals.com/professional/pediatrics/perinatal-hematologic-disorders/perinatal-anemia>. Accessed: April 30, 2023.
3. Alqahtany FS. Anaemia of the newborn: A review. *Eur J Mol Clin Med.* 2020;07(02): 3590.
4. WHO guideline on use of ferritin concentrations to assess iron status in individuals and populations. Geneva: World Health Organization. Licence: CC BY-NC-SA 3.0 IGO; 2020.

5. Beard J, deRegnier RA, Shaw M, Rao R, Georgieff M. Diagnosis of iron deficiency in infancy. *Lab Med* 2007;38:103–108.
6. Domellöf M, Hernell O, Dewey KG, Lönnerdal B, Cohen RJ. The Diagnostic Criteria for Iron Deficiency in Infants Should Be Reevaluated, *J Nutr.* 2002; 132(12):3680-6. Available:https://doi.org/10.1093/jn/132.12.3680.
7. Cook JD, Finch CA, Smith NJ, Evaluation of the iron status of a population. *Blood.* 1976;48:449-455.
8. Karakochuk CD, Hess SY, Moorthy D, Namaste S, Parker ME, Rappaport AI et al. HEmoglobin MEasurement (HEME) Working Group. Measurement and interpretation of hemoglobin concentration in clinical and field settings: a narrative review. *Ann N Y Acad Sci.* 2019;1450(1):126–46.
9. Addo OY, Yu EX, Williams AN, Young MF, Sharma AJ, Mei Z et al. Evaluation of hemoglobin cutoff levels to define anemia among healthy individuals. *MDAMA Network Open.* 2021;4(8):e2119123. DOI:10.1001/jamanetworkopen.2021.19123
10. Linderkamp O, Zilow EP, Zilow G. Kritische Hämoglobinwerte bei Neugeborenen, Säuglingen und Kindern [The critical hemoglobin value in newborn infants, infants and children]. *Beitr Infusionsther.* 1992;30:235-46; discussion 247-64. German. PMID: 1284712.
11. Siddappa AM, Rao R, Long JD, Widness JA, Georgieff MK. The assessment of newborn iron stores at birth: A review of the literature and standards for ferritin concentrations. *Neonatology.* 2007;92: 73–82.
12. Cheng C, Juul S. Iron balance in the neonate. *Neoreviews* 2011;12:148-158.
13. Rao R, Georgieff MK. Iron in fetal and neonatal nutrition. *Semin Fetal Neonatal Med.* 2007;12: 54–63.
14. WHO. Serum ferritin concentrations for the assessment of iron status and iron deficiency in populations. *Vitamin and Mineral Nutrition Information System.* Geneva, World Health Organization, (WHO/NMH/NHD/MNM/11.2); 2011. Available:http://www.who.int/vmnis/indicators/serum_ferritin pdf, accessed May 05, 2020
15. American Academy of Family Physician. Avoid using hemoglobin to evaluate patients for iron deficiency in susceptible populations. Instead, Use ferritin. Available:https://www.aafp.org/pubs/afp/collections/choosing-wisely/432.html Accessed: May 09, 2023.
16. Joy Y. Zhang, Jing Wang, Qinsheng Lu, Meizhen Tan, Ru Wei, Gendie E. Lash; Iron stores at birth in a full-term normal birth weight birth cohort with a low level of inflammation. *Biosci Rep.* 2020; 40(12): BSR20202853. Available:https://doi.org/10.1042/BSR20202853
17. Kim H, Park S, Lee E. Iron status in small for gestational age and appropriate for gestational age infants at birth. *Clin Exp Pediatr.* 2019;62(3):102-107. DOI: https://doi.org/10.3345/kjp.2018.06653
18. Sweet DG, Savage G, Tubman TRJ, Lappin TRJ, Halliday HL. Study of maternal influences on fetal iron status at term using cord blood transferrin receptors. *Arch Dis Child Fetal Neonatal Ed* 2001; 84: F40–F43.
19. Siddappa AM, Georgieff MK, Wewerka S, Worwa C, Nelson CA, deRegnier RA. Iron deficiency alters auditory recognition memory in newborn infants of diabetic mothers. *Pediatr Res.* 2004; 55:1034-41.
20. World Health Organization/Centers for Disease Control and Prevention. Assessing the iron status of populations: a report of a joint World Health Organization/Centers for Disease Control technical consultation on the assessment of iron status at the population level. Geneva; 2004 April. Available:http://whqlibdoc.who.int/publications/2004/9241593156_eng.pdf. Accessed January 4, 2008.
21. Chen X, Scholl TO, Stein TP. Association of elevated serum ferritin levels and the risk of gestational diabetes mellitus in pregnant women: The Camden study *Diabetes Care.* 2006;29: 1077–1082.
22. Whitley E, Ball J. Statistics review 4: Sample size calculations. *Critical Care.* 2002;6:335–341.
23. Carpani G, Marini F, Buscaglia M, Sinigaglia E, Moroni G. Red cell and plasma ferritin in normal fetuses at different ages of gestation. *Eur J Haematol.* 1992;49:260–262.
24. le Cessie S, Verhoeff FH, Mengistie G, Kazembe P, Broadhead R, Brabin B. Changes in haemoglobin levels in infants in Malawi: effect of low birth weight and

- fetal anaemia. *Arch Dis Child Fetal Neonatal Ed.* 2002;86:F182-F187.
25. Adediran A, Gbadegesin A, Adeyemo T, Akinbami A, Osunkalu V, Ogbenna A, et al. Cord blood haemoglobin and ferritin concentrations in newborns of anaemic and non-anaemic mothers in Lagos, Nigeria. *Nig Med J.* 2013; 54(1):22-26. DOI: 10.4103/0300-1652.108889.
 26. Tiruneh T, Shiferaw E, Enawgaw B. Prevalence and associated factors of anemia among full-term newborn babies at University of Gondar comprehensive specialized hospital, Northwest Ethiopia: a cross-sectional study. *Ital J Pediatr.* 2020; 46(1):1. DOI: 10.1186/s13052-019-0764-1
 27. Eslami Z, Ghilian R, Abbasi F. Evaluation of hemoglobin concentration of cord, capillary and venous sampling in neonates. *Iran J Ped Hematol Oncol.* 2012;2(4):159-63. Epub 2012 Sep 22.
 28. Kalteren W, Mebius M, Verhagen E, Tanis J, Kooi E, Bos A. Neonatal hemoglobin levels in preterm infants are associated with early neurological functioning. *Neonatology.* 2011;118(5): 593-9. Available:https://doi.org/10.1159/000518655
 29. Hasanbegovic E, Cengic N, Hasanbegovic S, Heljic J, Lutolli I, Begic E. Evaluation and treatment of anemia in premature infants. *Med Arch.* 2016;70(6): 408-412. DOI: 10.5455/medarh.2016.70.408-412
 30. Banerjee, J., Asamoah, F.K., Singhvi, D. et al. Haemoglobin level at birth is associated with short term outcomes and mortality in preterm infants. *BMC Med.* 2015;13:16. Available:https://doi.org/10.1186/s12916-014-0247-6
 31. Jopling J, Henry E, Wiedmeier SE, Christensen RD. Reference ranges for hematocrit and blood hemoglobin concentration during the neonatal period: data from a multihospital health care system. *Pediatrics.* 2009;123(2):e333-7. DOI: 10.1542/peds.2008-2654. PMID: 19171584
 32. Centers for Disease Control and Prevention (CDC). Recommendations to prevent and control iron deficiency in the United States. *Morb. Mortal. Wkly. Rep.* 1998;47:1–29.
 33. Uneke CJ, Duhlińska DD, Ujam TN. Effects of Maternal Plasmodium falciparum malaria and HIV infection on birth weight in southeastern Nigeria. *McGill J Med.* 2009;12(2):42. PMID: 21152333; PMCID: PMC2997242.
 34. West B, Arthur A, Aitafo J, Omunakwe H. Prevalence, Associated Factors and Outcome of Newborns with Anaemia as Seen in Rivers State University Teaching Hospital, Nigeria. *Arch Curr Res Int.* 2021;21(4):23-24. DOI: 10.9734/ACRI/2021/v21i430241
 35. Brabin BJ, Kalanda BF, Verhoeff FH, Chimsuku LH, Broadhead RL. Risk factors for fetal anaemia in a malarious area of Malawi. *Ann Trop Paediatr.* 2004;24: 311–21
 36. Lee S, Guillet R, Cooper EM, Westerman M, Orlando M, Kent T, et al. Prevalence of anemia and associations between neonatal iron status, hepcidin, and maternal iron status among neonates born to pregnant adolescents. *Pediatr Res.* 2016;79(1):42–48.
 37. Kalteran WS, Horst HJH, de Vetten L, Kooi EM, Bos AF. Perinatal anemia is associated with neonatal and neurodevelopmental outcomes in infants with moderate to severe perinatal asphyxia. *Neonatology.* 2018;114(4): 315-322. Available:https://doi.org/10.1159/000490369
 38. Augusta de Sá S, Willner E, Aguiar DuraesPereira T, Rosse de Souza V, TelesBoaventura G, Blondet de Azeredo V. Anemia in pregnancy: impact on weight and in the development of anemia in newborn. *Nutricion Hospitalaria.* 2015; 32(5):103-105. DOI: 10.3305/nh.2015.32.5.9186
 39. Al-Naseem A, Sallam A, Choudhury S, Thachil J. Iron deficiency without anaemia: a diagnosis that matters. *Clin Med J.* 2021;21(2):107-113. DOI: 10.7861/clinmed.2020-0582
 40. Collard KJ. Iron homeostasis in the neonate. *Pediatrics* 2009;4:1208-1216.
 41. Rao R, Georgieff MK. Iron therapy for preterms. *Clin Perinatol.* 2009;36:27-42.
 42. Siddappa AM, Georgieff MK, Wewerka S, Worwa C, Nelson CA, deRegnier RA. Iron deficiency alters auditory recognition memory in newborn infants of diabetic mothers. *Pediatr Res.* 2004;55:1034-41.
 43. Adediran A, Gbadegesin A, Adeyemo TA, Akinbami A, Osunkalu VO, Ogbenna AA, et al. Haemoglobin and ferritin concentrations in cord blood in a tertiary

- health centre in Nigeria. *Nig Q J Hosp Med.* 2011;21(4):284-9. PMID: 23175892.
44. Chew EC, Lam JC. Diagnosis and management of iron deficiency anaemia in children - A clinical update. *Proceedings of Singapore Healthcare.* 2012;21:278-295.
 45. Shao J, Rao R, Georgiff MK, Kaciroti N, Felt BT, Zhao Z, et al. Maternal serum ferritin concentration is positively associated with newborn iron stores in women with low ferritin status in late pregnancy. *J Nutr.* 2012;142:2004-2009.
 46. Baker RD, Frank R. The Committee on Nutrition. Diagnosis and prevention of iron deficiency and iron-deficiency anemia in infants and young children (0 –3 years of age). *Pediatrics.* 2010;126:1040 - 1050.
 47. Kawamura M, Nishida H. The usefulness of serial C-reactive protein measurements in managing neonatal infection. *Acta Pediatr.* 1995;84:10-13.
 48. Benitz W, Han M, Madan A, Ramachandra P. Serial serum C-reactive protein levels in the diagnosis of neonatal infection. *Pediatrics.* 1998;102:41.
 49. Kocabaş E, Sarıkçioğlu A, Aksaray N, Seydaoğlu G, Seyhun Y, Yaman A. Role of procalcitonin, C-reactive protein, interleukin-6, interleukin- 8 and tumor necrosis factor- α in the diagnosis of neonatal sepsis. *Turk J Pediatr.* 2007;49:7-20.

© 2023 Nwabueze; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/100035>